

Genetic Admixture Estimate in the Uruguayan Population Based on the Loci LDLR, GYPA, HBG, GC and D7S8

P. C. Hidalgo, M. Bengochea, D. Abilleira, A. Cabrera and I. Alvarez

*Laboratorio de Inmunogenética e Histocompatibilidad. Banco Nacional de Organos y Tejidos. Hospital de Clínicas "Manuel Quintela". CP. 11600. Montevideo. Uruguay
E-mail: bnot@hc.edu.uy*

KEYWORDS Polymarker; PM-PCR; population genetics; Latin American populations

ABSTRACT We have analyzed the allele and phenotype frequencies in five DNA loci: LDLR, GYPA, HBG, GC, and D7S8 in a sample from Uruguay. All the loci were in Hardy-Weinberg equilibrium (HWE) except the D7S8 locus. Our genetic admixture estimate showed evidence that the main genetic contribution comes from Europe with a small Amerindian and a minor African contribution with the admixture proportions: 84.1%, 10.4%, and 5.6% respectively. Genetic distances between the Uruguayan sample and several other Latin American populations revealed the closest genetic relationship with the Argentinean capital city, probably because its common history and demographic characteristics.

INTRODUCTION

Studies carried out in Latin American populations have shown that they have been originated by complex scenery of ethnic, cultural, and social relationships, moreover provide a valuable opportunity for population genetic studies (Salzano and Bortolini 2002). With the exception of some Amerindian isolate ethnic groups practically almost all Latin America populations have suffered an intense process of admixture all along in the last five centuries (Sans 2000).

Uruguay has been considered an immigrants country that, unlike other South American countries, has no Amerindian people and a small African contribution (Ribeiro 1969; Wagley, 1971). However, studies on HLA, blood groups, serum proteins, and red cell polymorphisms have documented the importance of the contribution of African and Amerindian people in the actual Uruguayan population (Alvarez et al. 1993; Sans 1997; Sans et al. 1993, 1997).

The Polymarker (PM) is detected by a polymerase chain reaction (PCR) multiplex amplification and typing kit of five specific

regions of the genetic loci: low density lipoprotein receptor (LDLR), glycophorin A (GYPA), hemoglobin G gamma globin (HBG), D7S8, and group specific component (GC); its use has proven highly efficient and informative for forensic and population genetics studies (Herin et al. 1994; Budowle et al. 1995; Chakraborty et al. 1995). A number of recent studies carried out in some American populations have provided additional information on polymarker distribution and consequently, useful to analyze relationships with different populations (Bertoni et al. 2003; Buentello-Malo et al. 2003).

Here, we analyzed the allele and genotype frequencies for the PM loci: LDLR, GYPA, HBG, GC, and D7S8 in a Uruguayan sample. The results are analyzed with two main goals: (1) to estimate the accumulated admixture contribution of the parental populations (i.e., European, African, and Amerindian) to the Uruguayan population; (2) to assess the relationship of the Uruguayan population with other Latin American populations using the available information.

MATERIALS AND METHODS

The sample was collected from 85 healthy, adult, and unrelated consenting individuals of both sexes, born in various regions of Uruguay, and sampled in Montevideo.

Genomic DNA was extracted from fresh whole blood using a previously described method comprising red cell lysis, proteinase K digestion,

Corresponding Autor: Dr. Pedro C. Hidalgo
Laboratorio de Inmunogenética e Histocompatibilidad.
Banco Nacional de Organos y Tejidos. Hospital de
Clínicas "Manuel Quintela". Ave. Italia s/n. Piso 4.
CP. 11600. Montevideo. Uruguay.
Phone: (5982) 487-2815; Fax: (5982) 487-7411;
E-mail: phidalgo@montevideo.com.uy

salt extraction, and ethanol precipitation (Miller et al. 1988) with slight modifications. PCR amplification and typing for the systems LDLR, GYPA, HBGG, GC and D7S8 were carried out using the AmpliType® PM PCR Amplification and Typing Kit (Polymarker Multiplex; Applied Biosystems; Foster City, CA) according to the manufacturer's protocol.

The allele frequencies were calculated by the gene count method (Li 1976). Departures from Hardy-Weinberg equilibrium were tested by the chi-square test and by the heterozygote deficiency test (Rousset and Raymond 1995) using the software Genepop (v.3.4; Raymond and Rousset 1995). Unbiased expected heterozygosity for single loci and average heterozygosity were computed according to Nei (1987).

Admixture estimates were calculated by the method described by Elston (1971), using a trihybrid model. The gene frequencies for the parental populations were calculated as an unweighted average from European, African, and Amerindian samples. The parental gene frequencies, and their sources, are listed in the Appendix.

Estimates of Nei's standard genetic distances (Nei 1972) were computed to compare the Uruguayan sample with other American populations in which the PM allele frequencies were available. The selected populations share common historical patterns, as well as comparable parental contributions regarding its origins (European, Amerindian, African). The available populations were: four African-derived: Costa Rican Afro-Caribbeans (Morales et al. 2001), Haití (Peterson et al. 2000), French Antilles (Peterson et al. 2000), and Brazilian Mulattoes (Soares-Viera et al. 2003); five heterogeneous samples of South American populations: Brazilian Whites (Soares-Viera et al. 2003), Chilean (Jorquera and Budowle 1998), Argentinean (Padula et al. 1996), Colombian (Castillo et al. 1996), and Uruguayan (this paper); three samples from North and Central America: Mexico City (Peterson et al. 2000), Nicaraguan (Morera et al. 2001), Costa Rican (Morales et al. 2001); three admixed Mexican Native Indian populations: Mixteca-Alta, Mixteca-Baja, and Nahuas-Xochimilco (Buentello-Malo et al. 2003). US Blacks (Peterson et al. 2000), and two samples of Hispanics from United States (West and East) (Bertoni et al. 2003) were also included as references. The neighbor-

joining method (NJ) was used to display the matrix of pairwise distance (Saitou and Nei 1987). The PHYLIP (v.3.5c; Felsenstein 1993) package of programs was employed to construct phylogenetic trees from the distance matrices. We used the programs GENDIST, SEQBOOT and then run NEIGHBORN. The consensus tree was built with CONSENSUS program.

RESULTS

Table 1 shows the phenotypes, allele frequencies, and unbiased expected heterozygosity of the studied loci. Departures from the Hardy-Weinberg genetic equilibrium at the single locus level were not statistically significant, with the only exception of locus D7S8 in which we also found a significant departure from Hardy-Weinberg expectation when the heterozygote deficiency test was applied ($p = 0.028$). When all the five loci were analyzed with the multi-locus test for the Hardy Weinberg expectation for heterozygote deficit (Rousset and Raymond 1995), no detectable deviation was observed ($p = 0.1237 \pm 0.0045$). Unbiased expected heterozygosity ranged from 0.4595 (D7S8) to 0.6155 (GC). Overall average heterozygosity showed a slightly high level with a value of 0.5125.

Table 2 presents the estimated admixture proportion of European, African and Amerindian genes in our sample. We obtained a strong global presence of 84.1% genes from European, followed by the Amerindian component (10.4%), and a minor African contribution (5.6%). We obtained an equivalent result by using Chakraborty's identity method (Chakraborty, 1985) (Table 2).

Nei's (1972) pairwise genetic distance matrix is presented in Table 3, showing the relationship of different American populations. The greatest separate the African derived and non-African derived American populations (Table 3). The distances among South American populations were consistently low. Brazilian whites and Chileans show the shortest distance, whereas Colombians and Uruguayans show the highest distance. To visualize the affinity between the studied populations the consensus NJ tree was drawn (Fig. 1). The tree shows two main clusters, whereas African derived populations are distinctly separated from the other samples. Uruguay and Argentine show single cluster and are distinctively different from other

Table 1: Phenotypes and allele frequencies at five loci.

<i>System</i>	<i>Phenotype</i>	<i>Observed frequency</i>	<i>Expected frequency</i>	<i>Chi-square</i>	<i>d.f.</i>	<i>P</i>	<i>HT¹</i>
LDLR							
N = 80	A-A	14	11.893	0.986	1	0.321	0.223
	A-B	34	38.214				
	B-B	32	29.893				
GYP A							
N = 85	A-A	32	32.929	0.185	1	0.667	0.745
	A-B	42	40.142				
	B-B	11	11.929				
HBGG							
N = 85	A-A	16	15.982	0.154	3	0.985	0.587
	A-B	39	38.970				
	A-C	3	3.065				
	B-B	23	23.172				
	B-C	4	3.686				
	C-C	0	.124				
D7SB							
N = 85	A-A	40	35.473	4.633	1	0.031*	0.028*
	A-B	30	39.053				
	B-B	15	10.473				
GC1							
N = 85	A-A	12	10.473	0.829	3	0.843	0.276
	A-B	10	9.586				
	A-C	26	29.467				
	B-B	2	2.077				
	B-C	13	13.260				
	C-C	22	20.136				

	<i>Locus</i>				
	<i>LDLR</i>	<i>GYP A</i>	<i>HBGG</i>	<i>D7SB</i>	<i>GC</i>
<i>Allele</i>					
A	0.3875	0.6245	0.4353	0.6471	0.3529
B	0.6135	0.3765	0.5235	0.3529	0.1588
C	0.0412	0.4883			
<i>Heterozygosity</i>					
Unbiased	0.4777	0.4723	0.5379	0.4595	0.6155
Observed	0.4250	0.4941	0.5412	0.3529	0.5765

¹HT: p values heterozygote deficiency test; *p < 0.05

Latin American populations. A second cluster consists in three subclusters: Chile and Brazilian whites integrated one of them, Nicaragua and Mixteca Alta (Mexico) integrated the second one, and the third one includes three Mexican samples, Colombians, admixed population of Costa Rica, and US Hispanics from the West of the country. Interestingly, US Hispanic from the West and from the East was always placed in separate cluster.

DISCUSSION

Our admixture estimates similar to the ones reported for Montevideo (92% European, 7% African, and 1% Amerindian contributions), and

Table 2: Admixture estimates in the Uruguayan sample.

<i>Method</i>	<i>Estimated proportion of parental contribution</i>		
	<i>European</i>	<i>African</i>	<i>Amerindian</i>
Elston (1971)	0.841	0.0536	0.104
Chakraborty (1986) ¹	0.870±0.026	0.027± 0.010	0.103±0.030

¹R² = 0.992

intermediate between them and the ones for the northeast of the country (65%, 15%, and 20% respectively) based on blood groups, serum proteins and red cell polymorphisms (Sans et al. 1997). However, African admixture is slightly lower in our estimation. The differences among

Table 3: Nei's genetic distances.

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)
HAITI	0.021																
AFUS	0.007	0.005															
AFROCR	0.036	0.099	0.068														
BRMUL	0.045	0.123	0.085	0.005													
MXA	0.093	0.180	0.140	0.027	0.030												
MXB	0.075	0.176	0.127	0.023	0.011	0.020											
NXO	0.129	0.252	0.196	0.044	0.036	0.013	0.012										
NICA	0.077	0.172	0.126	0.029	0.020	0.013	0.007	0.014									
HUSE	0.075	0.175	0.127	0.013	0.008	0.017	0.006	0.014	0.012								
HUSW	0.093	0.195	0.146	0.021	0.020	0.009	0.013	0.010	0.010	0.006							
MXC	0.111	0.220	0.164	0.034	0.033	0.020	0.025	0.021	0.020	0.012	0.013						
ADMCR	0.109	0.227	0.168	0.031	0.022	0.017	0.010	0.010	0.010	0.007	0.004	0.011					
CHI	0.132	0.267	0.200	0.047	0.028	0.034	0.012	0.015	0.017	0.012	0.017	0.018	0.005				
COL	0.145	0.279	0.215	0.052	0.038	0.022	0.014	0.007	0.015	0.018	0.011	0.026	0.005	0.007			
BRW	0.106	0.225	0.167	0.029	0.015	0.032	0.010	0.018	0.018	0.005	0.015	0.016	0.008	0.004	0.016		
URU	0.095	0.212	0.157	0.032	0.021	0.047	0.015	0.024	0.027	0.010	0.019	0.031	0.017	0.018	0.026	0.011	
ARG	0.100	0.213	0.162	0.025	0.017	0.026	0.010	0.010	0.020	0.005	0.011	0.023	0.012	0.015	0.017	0.008	0.005

1) ANTFR: French Antilles; 2) HAITI; 3) AFUS: US Blacks; 4) AFROCR: Costa Rica Afro-Caribbean; 5) BRMUL: Brazilian Mulattoes; 6) MXA: Mixteca-Alta; 7) MXB: Mixteca-Baja; 8) NXO:Nahuas-Xochimilco; 9) NICA: Nicaragua; 10) HUSE: US Hispanics Eastern; 11) HUSW:US Hispanics Western; 12) MXC: Mexico City; 13) ADMCR: Costa Rica admixed; 14) CHI: Chile; 15) COL: Colombia; 16) BRW: Brazilian Whites; 17) URU:Uruguay; 18)ARG:Argentine;




Fig.1. Consensus Neighbor-Joining tree from the Nei's distances.

AFROUS: US Blacks; ANTFR: French Antilles; AFROCR: Costa Rica Afro-Caribbean; BRMUL: Brazilian Mulattoes; HUSE: US Hispanics Eastern; NICA: Nicaragua; MXB: Mixteca-Baja; ADMCR: Costa Rica admixed; HUSW: US Hispanics Western; MXC: Mexico City; MXA: Mixteca Alta; COL: Colombia; NXO: Nahuas-Xochimilco; BRW: Brazilian Whites; CHI: Chile; ARG:Argentine; URU: Uruguay.

the samples, admixture estimation methods, parental selection, and genetic markers used, make clear that direct comparisons between these studies warrant caution. Our results suggest that the Uruguayan population present a moderate admixture level with a predominant European parental contribution and a more modest Amerindian and African parental contributions. Then, the data support an admixture trihybrid model modulated by the European migration waves occurred in the last century (Oddone 1966).

Some further comments should be made on the NJ tree in Figure 1. In this tree is evident the relationship between Uruguay and the sample from Argentine (Buenos Aires). Both regions have a common immigrant background with an important Spanish and Italian components (Oddone 1966) Moreover, our results are not surprising if we considering that, the population of the densely populated city of Buenos Aires and surrounding areas, show an African contribution between 3,5 and 7%, and Amerindian contribution between 14 and 33% (López-Camelo et al. 1996, Avena et al. 1999, 2001). In general, the Uruguayan population genetic make up is similar to other admixed urban Latin American populations, with a predominant European-derived component.

The Uruguayan-Argentinean cluster is separated to the other cluster that deserve our attention, formed by Chileans, Brazilian whites, Colombians, Costa Ricans, US Hispanics from the West, and 4 Mexican populations. This cluster denotes the Amerindian contribution, greater in those areas (Sans 2000). US Hispanics from the East is also in this cluster but a little more distant, probably indicating more African contribution (Bertoni et al. 2003) African-derived populations do not clusterize and are beyond the others. Moreover, the results of the frequencies of the alleles studied here clearly demonstrate a significant heterogeneity among the populations of Latin American in agreement with the different regional proportions of the European, Amerindian and African parental contributions to the admixture process for each American country. All the Latin American populations analyzed are far away of having a homogeneous ethnic composition, and none of them is free of gene flow and admixture effects. This study illustrate the genetic heterogeneity present in the Latin American populations, for

that reason, in the context of genetic epidemiology, future studies related to disease risk factors will need to address a research design that subdivides the population using genetic parameters (Chakraborty et al. 1999).

ACKNOWLEDGMENTS

Thanks are expressed to Mónica Sans for their invaluable commetaries.

REFERENCES

- Alvarez I, Sans M, Toledo R, Sosa M, Bengochea M, Salzano FM 1993. HLA gene and haplotype frequencies in Uruguay. *Int J Antropol*, **8**:163-168.
- Avena S, Goicochea A, Rey JA, Agosti JC, Carnese FR 1999. Análisis de la participación del componente indígena en una muestra hospitalaria de la ciudad de Buenos Aires. *Rev Argentina Antropol Biol*, **2**: 211-225.
- Avena S, Goicochea A, Dugoujon JM, Rey J, Slepoy A, Dejean C, Carnese FR 2001. Mezcla génica en la población de la región metropolitana de Buenos Aires. *Rev Argentina Antropol Biol*, **3**: 45.
- Bertoni B, Budowle B, Sans M, Barton SA, Chakraborty R 2003. Admixture in hispanic : Distribution of ancestral population contributions in the continental United States. *Hum Biol*, **75**:1-11.
- Budowle B, Lindsey JA, De Cou JA, Koons BW, Giusti AM, Comey CT 1995. Validation and population studies of the loci LDLR, GYPA, HBG, D7S8, and GC (PM loci), and HLA-DQ alpha using a multiplex amplification and typing procedure. *J Forensic Sci*, **40**:45-54.
- Buentello-Malo L, Peñalosa-Espinosa RI, Loeza F, Salamanca-Gómez F, Cerda-Flores RM 2003. Genetic structure of seven Mexican indigenous populations based on five Polymarker loci. *Am J Hum Biol*, **15**:23-28.
- Castillo MJ, Paredes M, Peñuela C, Bustos I, Jimenez M, Galindo, A 1996. Determination of the allele and genotype frequencies of loci HLA-DQA1, LDLR, GYPA, HBG, D7S8 and GC in Bogota - Colombia. *Adv Forensic Haemogenet*, **6**: 503-505.
- Chakraborty R 1985. Gene identity in racial hybrids and estimation of admixture rates. In: Y Ahuja and JV Neel (Eds.): *Genetics Microdifferentiation in Human and Other Animal Populations*. Delhi, India: Indian Anthropological Association, Delhi University Anthropology Department, pp. 171-180.
- Chakraborty R, Jin L, Zhong Y, Deka R 1995. Intra- and inter-population variation at VNRT short tandem repeat and polymarker loci and their implications in forensic and paternity analysis. In: *Proceedings of the 5th. Internationla Symposium on Human Identification 1994*. Promega Corporation: Madison, Wisconsin, pp. 29-41.
- Chakraborty B, Fernández-Esquer ME, Chakraborty R 1999. Is being Hispanic a risk factor for non-insulin dependent diabetes mellitus (NIDDM)? *Ethn. Dis*. **9**: 278-283.
- Elston RC 1971. The estimation of admixture in racial

- hybrids. *Ann Hum Genet*, **35**:9-17.
- Felsenstein J 1993. *PHYLIP (Phylogenetic Inference Package)*. Version 3.5c. Department of Genetics, University of Washington, Seattle, WA.
- Herrin G, Fildes N, Reynolds R 1994. Evaluation of the AmpliType® PM DNA test system on forensic case samples. *J Forensic Sci*, **39**:1247-1253.
- Jorquera H, Budowle B 1998. Chilean population data on ten PCR-based loci. *J Forensic Sci*, **43**: 171-173.
- Li CC 1976. *First Course in Population Genetics*. California: Pacific Grove .
- López-Camelo JS, Cabello PH, Dutra MG 1996. A simple model for the estimation of congenital malformation frequencies in racially mixed populations. *Braz J Genet*, **19**: 659-663.
- Miller SA, Dykes DD, Polesky HT 1988. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucl Acids Res*, **16**:1215
- Morales AI, Morera B, Jimenez-Arce G, Sanchez-Rivera G, Calafell F, Barrantes R 2001. Allele frequencies of markers LDLR, GYPA, D7S8, HBGG, GC, HLA-DQA1 and D1S80 in the general and minority populations of Costa Rica. *Forensic Sci Int*, **124**:1-4.
- Morera B, Sanchez-Rivera G, Jimenez-Arce G, Calafell F, Morales-Cordero AI 2001. Nicaraguan population data on LDLR, GYPA, D7S8, HBGG, GC and HLA-DQA1 loci. *Rev Biol Trop*, **49**:1253-1260.
- Nei M 1972. Genetic distance between populations. *Am Nat*, **106**:283-292.
- Nei M 1987. *Molecular Evolutionary Genetics*. New York: Columbia University Press.
- Odone JA 1966. *La Emigración Europea al Río de La Plata*. Montevideo: Ediciones de la Banda Oriental.
- Padula RA, Gangitano DA, Juvenal GJ., Budowle B 1999. Allele frequency in the population of Buenos Aires (Argentina) using AmpliType PM + DQA1. *J Forensic Sci*, **44**: 1320.
- Peterson BL, Su B, Chakraborty R, Budowle B, Gaensslen RE 2000. World population data for the HLA-DQA1, PM® and D1S80 loci with least and most common profile frequencies for combination of loci estimated following NRC II guidelines. *J Forensic Sci*, **45**:118-146.
- Raymond M, Rousset F 1995. GENEPOP (version 1.2): population exact tests and ecumenism. *J Heredity*, **86**:248-249.
- Ribeiro D 1969. *Las Américas y la Civilización. 3. Los Pueblos Trasplantados. Civilización y Desarrollo*. Buenos Aires: Centro Editorial de América Latina.
- Rousset F, Raymond M 1995. Testing heterozygote excess and deficiency. *Genetics*, **140**:1413-1419.
- Saitou N, Nei M 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol*, **4**: 406-425.
- Salzano FM, Bortolini MC 2002. *The Evolution and Genetics of Latin American Populations*. Cambridge: Cambridge University Press.
- Sans M 1997. Estudio de marcadores genéticos en la población uruguaya y su aplicación en el análisis del mestizaje. *Estudios en Antropología Biológica (UNAM, México)*, **6**:87-100.
- Sans M 2000. Admixture studies in Latin America: from the 20th to the 21st century. *Human Biology*, **72**:155-177.
- Sans M, Sosa M, Alvarez I, Toledo R, Bengochea M, Salzano FM 1993. Blood group frequencies and the question of race admixture in Uruguay. *Interciencia*, **18**:29-32.
- Sans M, Salzano FM, Chakraborty R 1997. Historical genetics in Uruguay: estimates of biological origins and their problems. *Hum Biol*, **69**:161-170.
- Scholl S, Budowle B, Radecki K, Salvo M 1996. Navajo, Pueblo, and Sioux population data on the loci HLA-DQA1, LDLR, GYPA, HBGG, D7S8, GC, and D1S80. *J Forensic Sci*, **41**:47-51.
- Soares-Vieira JA, Billerbeck AE, Iwamura ES, Otto PA 2003. Brazilian population data on the polymerase chain reaction-based loci LDLR, GYPA, HBGG, D7S8, and Gc. *Am J Forensic Med Pathol*, **24**:283-287.
- Wagley C 1971. The formation of the American people. In: FM Salzano (Ed.): *The Ongoing Evolution of Latin American Populations*. Springfield: Charles C. Tomas, pp. 19-39.
- Walkinshaw M, Strickland L, Hamilton H, Denning K, Gayley T 1996. DNA profiling in two Alaskan Native populations using HLA-DQA1, PM, and D1S80 loci. *J Forensic Sci*, **41**:478-484.

APPENDIX

Parental gene frequencies¹

Allele	European	African	Amerindian
LDLR*A	0.4634	0.1821	0.634
LDLR*B	0.5636	0.8179	0.366
GYPA*A	0.534	0.5087	0.8239
GYPA*B	0.466	0.4313	0.1761
HBGG*A	0.4845	0.3815	0.2772
HBGG*B	0.503	0.2197	0.7112
HBGG*C	0.0125	0.3988	0.0116
D7S8*A	0.5972	0.6734	0.5678
D7S8*B	0.4026	0.3266	0.4322
GC*A	0.3088	0.0867	0.1486
GC*B	0.1495	0.8439	0.2916
GC*C	0.5417	0.0694	0.5598

¹ Sources of parental populations:
European: Spain, Portugal, and Italy from the compilation of <http://www.uni-duelssendorf.de/WWW/MedFak/Serology/polymarker.html>.
African: Nigeria, and Zimbabwe from Peterson et al. (2000).
Amerindian: Alaskan Amerindians from Walkinshaw et al. (1996), North American Indians: Navajo, Pueblo, and Sioux from Scholl et al. (1995), Mexican: Nahuas-Guerrero, Tzeltales, Otomies, and Purepecha from Buentello-Malo et al. (2003), and Bribri-Cabecar from Costa Rica (Morales et al. 2001). No data available for PM from South American Indians.