

Familial Robertsonian Translocation 13;21 in a Down Syndrome Patient with XYY/XY Mosaicism

Cyril Cyrus¹, Teena K.², Solomon F.D.Paul², Chandra N.¹, Meena J.³, Anuradha D.³,
Ramesh A.¹, Gopinath P.M.⁴ and Marimuthu K.M.⁵

¹ *Department of Genetics, Dr. ALM. PG. Institute of Basic Medical Sciences, University of Madras, Taramani, Chennai 600 113*

² *Department of Human Genetics, Sri Ramachandra Medical College & Research Institute (Deemed University), Porur, Chennai 600 116*

³ *Department of Medical Genetics, Institute of Obstetrics and Gynecology, Madras Medical College, Government Hospital for Women and Children, Egmore, Chennai 600 008*

⁴ *KMC Life Sciences Center, Manipal Academy of Higher Education, Manipal-576104*
⁵ *26, I Main Road, Indira Nagar, Adyar, Chennai 600 020*

KEYWORDS Down syndrome; mosaicism; Robertsonian translocation; XYY males

ABSTRACT Double aneuploidy involving XYY and trisomy 21 is rare. XYY/XY mosaicism has been described in only a single Down syndrome patient. The Robertsonian translocation t(13;21) is also rare among these individuals. We report for the first time the occurrence of t(13;21) in a mosaic XYY Down male. Analysis of GTG-banded metaphases revealed the karyotype of the proband to be mos 47,XYY,der(13;21)(q10;q10),+21/46,XY,der(13;21),+21. Both his father and paternal grandfather were found to be carriers for the translocation. This 10-month-old child who presented with typical features of Down syndrome, developed leukemia and died at the age of 2¼ years.