Familial Robertsonian Translocation 13;21 in a Down Syndrome Patient with XYY/XY Mosaicism

Cyril Cyrus¹, Teena K.², Solomon F.D. Paul², Chandra N.¹, Meena J.³, Anuradha D.³,
Ramesh A.¹, Gopinath P.M.⁴ and Marimuthu K.M.⁵

¹ Department of Genetics, Dr. ALM. PG. Institute of Basic Medical Sciences, University of Madras, Taramani, Chennai 600 113

² Department of Human Genetics, Sri Ramachandra Medical College & Research Institute (Deemed University), Porur, Chennai 600 116

³ Department of Medical Genetics, Institute of Obstetrics and Gynecology, Madras Medical College, Government Hospital for Women and Children, Egmore, Chennai 600 008

⁴ KMC Life Sciences Center, Manipal Academy of Higher Education, Manipal-576104
⁵ 26, I Main Road, Indira Nagar, Adyar, Chennai 600 020

KEYWORDS Down syndrome; mosaicism; Robertsonian translocation; XYY males

ABSTRACT Double aneuploidy involving XYY and trisomy 21 is rare. XYY/XY mosaicism has been described in only a single Down syndrome patient. The Robertsonian translocation t(13;21) is also rare among these individuals. We report for the first time the occurrence of t(13;21) in a mosaic XYY Down male. Analysis of GTG-banded metaphases revealed the karyotype of the proband to be mos 47,XYY,der(13;21)(q10;q10),+21/46,XY,der(13;21),+21. Both his father and paternal grandfather were found to be carriers for the translocation. This 10-month-old child who presented with typical features of Down syndrome, developed leukemia and died at the age of 2½ years.

INTRODUCTION

Aneuploidy is the most frequently observed chromosomal abnormality in human liveborns, abortuses, blastomeres and oocytes. It is reported to occur in at least 3-4 percent of all clinically recognized pregnancies. XYY is a common sex chromosomal aneuploidy with an estimated incidence of 1 in 1000 male live births (Nussbaum et al. 2001). However, these cases usually go undetected during childhood due to inconsistent phenotypic features. Down syndrome (DS) or trisomy 21, with its characteristic clinical features is the most common autosomal aneuploidy with an incidence of 1:700. Double aneuploidies involving chromosome 21 and sex chromosomes have been occasionally reported. Parmar et al. (2002) found 17 published cases of 48,XYY,+21 in Medline 1965-2002 while describing an infant having the same karyotype. There was only one case of XYY Down syndrome male showing mosaicism for the Y chromosome (Schwanitz and

Hagner 1978). The clinical features of DS were apparent in these individuals. XYY males are generally fertile and bear normal children while males with DS are sterile. Thus, it would be reasonable to propose that DS males with XYY chromosome pattern may also be infertile (Parmar et al. 2002).

Robertsonian translocations are known to account for 4-5 percent of all DS cases. These translocations are common balanced chromosomal rearrangements with an estimated incidence of 1 in 1000 in the general population. They can occur *de novo* with a high mutation rate of 3.92×10^{-4} per gamete per generation (Jacobs 1981; Vogel and Motulsky 1997; Berend et al. 1998). The carriers of these translocations are usually phenotypically normal. This paper describes a child with Down syndrome who was found to be an XYY/XY mosaic and carried a Robertsonian translocation t(13;21) transmitted down from his paternal grandfather. This is the first report of an XYY Down syndrome individual with t(13;21).

MATERIALS AND METHODS

PHA-stimulated lymphocyte cultures were set up and metaphase preparations were obtained

Address for all correspondence: Dr. N. Chandra, Department of Genetics, Dr. ALM. PG. Institute of Basic Medical Sciences, University of Madras, Taramani, Chennai 600 113, Tamil Nadu, India
Telephone: 044-24480769, Fax: 044-24926709
E-mail: chandrarsn@yahoo.co.in

following the method of Hungerford (1965) from the patient, his parents and unaffected sister. Fifty GTG-banded (Seabright 1971) metaphases were analyzed from each individual. His grandparents and paternal uncle were also investigated. Chromosomal anomalies were designated as per the guidelines laid down by the nomenclature committee (ISCN 1995). Metaphases were photographed under oil immersion lens using NOVA film in Nikon photomicroscope (Labophot 2) for karyotype preparation.

CASE REPORT

A 10-month-old male, second-born of a 24-year-old mother and 35-year-old father, was referred for chromosomal analysis because of dysmorphic features suggestive of Down syndrome. The infant was delivered at term following an uneventful gestation. The proband cried at birth and weighed 2.75 kg. The proband exhibited delayed developmental milestones. The typical features observed were open fontanelle, brachycephaly, slanting palpebral fissures, epicanthal folds, low set ears, flat nasal root, open mouth, short broad hands, simian crease on the left hand, clinodactyly, hypotonia, gap between 1st and 2nd toes and plantar furrow. The child was reported to have suffered from frequent infections. On follow-up, he was found to have developed acute myeloid leukemia at the age of

2¼ years and died. The couple also has a healthy 3½-year-old daughter.

RESULTS

The patient was found to have Down syndrome with an inherited Robertsonian translocation and also, XYY/XY mosaicism – mos 47,XYY,der(13;21)(q10;q10),+21[64]/46,XY,der(13;21),+21[36] (Figs. 1 & 2). Fifty additional metaphases were analyzed to determine the frequencies of the two cell lines. Both his father and his paternal grandfather were found to be carriers for this translocation (Fig. 3). The chromosomal constitution of his mother and elder sister was normal. His paternal grandmother and uncle also showed a normal karyotype. However, his paternal aunt and her two children were not available for study (Fig. 4).

DISCUSSION

Cytogenetic surveys of neonates have revealed that approximately one in 500 males is born with an extra sex chromosome (Abramsky and Chapple 1997). The commonest indication for a 47,XYY male to be karyotyped was reported to be developmental delay and / or behaviour problems. However, most males having an extra Y chromosome will go through life without a cytogenetic investigation due to inconsistent phenotypic features. The occurrence of XYY and

Fig. 1. A GTG – banded karyotype of the proband exhibiting 47,XYY,der(13;21)(q10;q10),+21

Fig. 2. A GTG – banded karyotype of the proband exhibiting 46,XY,der(13;21)(q10;q10),+21

Fig. 3. A GTG – banded karyotype of the proband's father showing 45,XY,der(13;21)(q10;q10)

- demonstrates diversity in Robertsonian translocation formation. *Am J Hum Genet*, **80**: 252-259
- Hungerford A 1965. Leucocytes cultured from small inocula of whole blood and the preparation of metaphase chromosomes by treatment with hypotonic KCl. *Stain Technology*, **40**: 333-338
- Jacobs PA 1981. Mutation rates of structural chromosome rearrangements in man. *Am J Hum Genet*, **33**: 44-54
- Jacobs PA, Hassold TJ 1995. The origin of numerical chromosome abnormalities. *Adv Genet*, **33**: 101-33.
- Leary PM, Webb PM, Melzer CW, Close HG 1975. Down's syndrome with additional XYY aneuploidy. *Clin Genet*, **8**: 55-58.
- Mitelman F 1995. *An International System for Human Cytogenetic Nomenclature*. Basel: S.Karger.
- Nussbaum RL, McInnes RR, Willard HF 2001. *Thompson & Thompson Genetics in Medicine*. 6th Edition, Philadelphia: W.B.Saunders Company, pp.149
- Parmar RC, Muranjan MN, Swami S 2002. Trisomy 21 with XYY. *Ind J Pediatr*, **69**: 979-981
- Schwanitz G, Hagner M 1978. Double trisomy as a mosaic - Case history (48, XYY,+21 / 47,XY,+21) and survey of the literature of mixed autosomal-gonosomal trisomies. *Acta Genet Med Gemellol (Roma)*, **27**: 67-74
- Seabright M 1971. A rapid banding technique for human chromosomes. *Lancet*, **ii**: 971-972
- Vogel F, Motulsky AG 1997. *Human Genetics - Problems and Approaches*. Berlin: Springer-Verlag, pp. 396-397