Diagnostic Potential of Genomic and Proteomic Signatures in Oral Cancer

Ranju Ralhan

*Department of Biochemistry, All India Institute of Medical Sciences, Ansari Nagar,
New Delhi 110 029, India*

KEYWORDS Oral cancer; genomics; proteomics; molecular signatures; biomarkers

ABSTRACT Oral cancer is a public health problem with increasing incidence and mortality rates worldwide. Despite rapid advances in treatment, 5year survival rates have not improved significantly. Major thrust is being laid on diagnosis of the disease in early stages, which is hampered by non-availability of specific diagnostic markers. Advances in genomics and proteomics have made global assessment of expressed genes and proteins in clinical samples feasible. Gene and protein expression profiles derived from clinical specimens have been used to distinguish differences between normal and malignant oral tissues, which are not obvious by clinical, or histologic characteristics. This review focusses on comprehensive analyses of gene expression patterns and proteomic signatures of oral dysplastic lesions and squamous cell carcinomas that have shown considerable promise to improve the discovery of biomarkers for progression of premalignant lesions, prediction of clinical outcome, identification of novel targets for therapy and future directions of molecular signatures of oral cancer in disease diagnosis and therapy.

INTRODUCTION

Oral Cancer is a major global public health problem with 300,000 new cases diagnosed annually. It is the most common cancer in males and third commonest cancer in females in India. Despite rapid advances in multimodality therapy, the morbidity and mortality rates of this devastating disease have not improved in decades. Early detection of oral cancer is the most effective way to improve survival. The treatment planning of oral cancer is mainly based on the tumor, node and metastasis (TNM) classification and histopathological diagnosis. These methods are subjective and often lack sensitivity to detect the disease in early stages. Furthermore, these methods do not reflect the aggressiveness of tumors, prognosis and response to therapy. Therefore, there is an urgent need to develop biomarkers to: identify high risk individuals, improve cancer detection in early stages, predict disease outcome and response to therapy. Rapid advances in high throughput genomic and proteomic technologies have paved the way for better understanding the molecular pathogenesis of oral cancer and identify candidate biomarkers for oral cancer. This review will focus on ongoing

research on genomic and proteomic signatures of oral cancer, our current understanding of molecular basis of oral carcinogenesis, characterization and validation of predictive / diagnostic potential of candidate biomarkers identified from human oral squamous cell carcinoma (OSCC) tissues as well as body fluids such as serum and saliva from oral cancer patients.

MULTISTEP ORAL CARCINOGENESIS

The development of oral cancer is a tobacco related multistep and multifocal process involving field cancerization and intraepithelial, clonal spread. Oral leukoplakia is the most common oral premalignant lesion (OPL) and is often a precursor of oral squamous cell carcinoma (OSCC) (Scully and Porter 2000 a, b). Erythroplakia, oral lichen planus and oral submucous fibrosis constitute other OPLs and premalignant conditions with well defined risk of malignant transformation. Histopathologically, OSCC development is a progress through a series of stages, from hyperplasia to varying degrees of dysplasia and to carcinoma *in situ*, prior to the development of invasive squamous cell carcinoma (reviewed in Reibel 2003; Hunter et al. 2005). The average annual rate of malignant transformation of oral leukoplakia has been proposed to be 1% based on several studies (Hunter et al. 2005). However, a recent study on long term treatment outcome of OPLs showed 7

Address Correspondance to: Ranju Ralhan, Department of Biochemistry, All India Institute of Medical Sciences, Ansari Nagar, New Delhi 110 029, India
Telephone: +91-11-26593478;
E-mail: ralhanr@hotmail.com

fold higher risk of non-homogenous leukoplakia for malignant development as compared with homogenous leukoplakia (Holmstrup et al. 2006). The “field cancerization” theory proposes that an area of oral mucosa can sustain an initial injury from repeated carcinogen exposure and proliferate in a premalignant state. Accumulation of additional genetic insults results in individual subsites of the premalignant field and causes progression to frank carcinoma (Ha and Califano 2003). According to this theory, relatively large areas of mucosa may have a high propensity of eventually developing carcinomas, yet not exhibit all the classic histological markers of malignancy.

Molecular signatures are the qualitative and quantitative patterns of groups of biomolecules (mRNA, proteins, peptide or metabolites) in a cell, tissue, biological fluid, or an organism. To apply this concept to oral cancer biomarker discovery, the measurements should ideally be non-invasive and performed in a single readout. The genetic characterization of field cancerization by Braakhuis et al. (2005) proposes the following model for head and neck carcinogenesis. A stem cell located in the basal layer of the oral mucosa acquires a genetic alteration in the initial phase. The division of this altered stem cell into daughter cells gives rise to a clonal unit of cells that all share this DNA alteration and constitute a patch. Subsequently, accumulation of additional genetic alterations result in the change of a patch into an expanding field that pushes the normal epithelium aside. These fields though macroscopically invisible may appear as oral patches. Finally, clonal selection within this field of preneoplastic cells leads to the development of cancer.

Molecular Alterations in Oral Cancer

Several epigenetic, genetic and metabolic alterations resulting from exposure of oral mucosa to tobacco carcinogens, alcohol or human papilloma virus have been identified in the multistep process of oral carcinogenesis. These have been extensively reviewed by us and others and thus will not be described here (Kaur and Ralhan 2003; Reibel 2003; Nagpal and Das 2003; Schliephake 2003; Bettendorf et al. 2004; Chimenos-Kustner et al. 2004; Warnakulasuriya and Ralhan 2006).

Comparative genomic hybridization (CGH), fluorescence *in situ* hybridization (FISH) and cytogenetic techniques have been used to detect

chromosomal rearrangements and genomic instability in OSCCs (Baldwin et al. 2005; Snijders et al. 2005). Extensive studies have reported the clinical significance of tumor suppressor genes and oncogenes in smoking associated OSCC in South East Asian populations including India. Some of these studies underscore the need to carry out multicentric studies in different populations with well defined tobacco habits for effective translation of biomarkers from bench to the clinic in different population and not translate the panel of biomarkers identified in Caucasians to the Asian population without large scale validations. Though several genes listed above have been associated with clinical and histopathological staging of OSCC (p16, p53, cyclin D1, p14^{ARF}, Ets-1), these markers provide little predictive or prognostic information for translation into a clinical setting (Arora et al. 2005a; Soni et al. 2005). In view of the vast heterogeneity in OSCCs it is unlikely that a single biomarker will be useful for early detection of all OSCCs. Therefore, there is a need for identification of a panel of biomarkers for diagnosis of OSCC.

Genomics and Proteomics for Biomarker Discovery in Oral Cancer

The advent of array technologies have provided a high throughput approach to monitor RNA for mRNA expression and DNA for genomic polymorphisms. In addition, epigenetic changes such as global hypomethylation and promoter hypermethylation of CpG rich areas have also been investigated. Rapid advances in proteomic technologies and mass spectrometry have provided the tools for analysis of protein profiles of OPLs and OSCCs. Furthermore, major thrust is being laid on analysis of DNA, RNA and protein profiles of biological fluids such as serum, plasma, saliva and urine for identification of biomarkers using non-invasive methods for multiple analyses, not only for early detection of oral cancer, but also for monitoring prognosis of the disease and response to therapy. Studies carried out using these techniques confirm the involvement of multiple pathways and cross-talks between pathways in oral tumorigenesis.

Genomics Signatures of OSCC

DNA microarrays are powerful tools for measuring gene expression profiles on a tissue-

wide level. Microarrays consist of millions of individual nucleotide sequences affixed to a solid substrate in a defined configuration. These nucleotide sequences (called “probes”) are complementary to the sequences of known and putative genes. Messenger RNA (termed “target”) that is extracted from a tissue of interest can be labeled with a fluorescent tag and hybridized to the microarray. By measuring the intensity of fluorescence at the precise locations of each probe on the microarray, the abundance of the target mRNA can be determined. The highly parallel nature of microarray technologies allows for the simultaneous measurement of thousands of genes in a relatively rapid manner. A variety of experimental platforms, probes, hybridization and signal detection methods, computational and bioinformatic analyses programs are being developed for global gene expression profiling both at micro- and macro- levels. Our group has previously identified differentially expressed gene patterns between OSCC and normal oral mucosa (Arora et al. 2005b). DNA microarrays have extensively been used to compare gene expression profiles of OSCCs with normal healthy oral tissues and reviewed (Todd and Wong 2002; Kuo et al. 2003; Otao-Rey et al. 2004; Brigitta and Wong 2006). These studies have led to identification of clusters of genes (also known as molecular signatures) that are associated with OSCC, thereby providing insight into the molecular mechanisms involved in oral tumorigenesis and identify candidate novel tumor derived biomarkers (Alevizos et al. 2001; Mendez et al. 2002; Conrads et al. 2003; Hwang et al. 2003; Chimenos-Kustner et al. 2004; Diamandis 2004; Carinci et al. 2005; Cheng et al. 2005; Dallas et al. 2005; Drake et al. 2005). Strikingly, there is a vast heterogeneity in the findings reported in several different studies. Kuo et al. (2002 a) suggested the possibility of classifying OSCC on the basis of gene expression patterns. Using a microarray of 4000 genes, they identified 210 genes that may be related to oral cancer. Some of these genes (CKS1, TSPY, CBK, TLEY and BCHE) have previously been reported in other cancers, but not in oral cancer. In addition, these authors also listed a set of genes whose expression is correlated with other classic prognostic factors, such as p53, MST1, HLA-DBQ1 and UBA52. Alevizos et al. (2001) and Mendez et al. (2002) also provided verifiable molecular signatures of

OSCC using DNA microarrays ; these data have been reviewed by Otao-Rey et al. (2004).

Another interesting study by Kuo et al. (2002 b) comparing gene expression profiles of oral normal, dysplastic and malignant cells using DNA microarrays and laser capture micro dissection found several genes and expressed sequence tags (ESTs) to be associated with cancer namely DDB2, a damage-specific DNA protein, to be a potential biomarker for the progression of OSCC. Intriguingly, expression profiles of dysplasia samples did not cluster together as a group, but were found in either ‘normal’ or ‘cancer’ clusters, suggesting that subtle changes occur in the progression of OSCC that are not captured by histopathology.

Odanti et al. (2006) compared gene expression profiles of oral leukoplakia and OSCC using Affymetrix Gene Chip system and suggested that gene abnormalities in cytoskeleton network components might be responsible for the development and progression of oral leukoplakia. Though the small sample size (4 leukoplakia and 2 OSCC) is a major limitation of this study. Nevertheless, it demonstrated that some of the 18 alternatively expressed genes were markedly down-regulated in OSCC compared with leukoplakia, further confirming the complexity of molecular alterations in the progression of OSCC proposed by the earlier study of Kuo et al. (2002 b, 2003a).

A major limitation of most studies using microarrays is that known OSCCs have been analysed and compared with normal counterparts. The major challenge is to identify molecular signatures that can accurately predict OSCC in unknown tissue specimens. In this context, Whipple et al. (2004) showed that principal component analysis (PCA) can be used with genomic microarray data to correctly predict the presence of OSCC in unknown tissue samples. Leethankul and co-workers (2000) have identified several differentiation and growth related genes in HNSCC using normal and malignant keratinocytes obtained from biopsy specimens by LCM using DNA microarray containing 588 cancer related genes . Oral cancer molecular signatures have also been investigated using high density oligonucleotide arrays. Ohyama et al. (2000) demonstrated the feasibility of procuring high amounts of target sample using LCM generated tissues for hybridizing high density oligo arrays. Using this optimized technology Alevizos et al. (2001), Mendez et al.

(2002) and others carried out gene expression profiling of OSCCs.

Giri et al. (2006) determined the molecular signatures associated with clinical outcome in patients with high risk head and neck cancer squamous cell carcinoma (HNSCC) treated with surgery and radiation. Patients who developed distant metastases has 205 genes differentially expressed as compared to patients with no recurrence. Genes associated with cell growth and proliferation; cell adhesion, DNA replication, recombination and repair, antiapoptotic pathways and angiogenesis were identified in tumors of patients with distant metastases. In contrast, tumors of patients with no recurrences showed discriminatory genes associated with the onset of apoptosis.

Liu et al. (2006) used array-CGH to detect genome wide changes in microdissected primary and metastatic OSCCs. Genomic alterations (TGFb2, cellular retinal binding protein 1 gene (CRBP1), PIK3CA, HTR1B, HRAS, ERBB3 and STK6) differed significantly between primary OSCC and neck lymph node metastatic tissues. An intriguing finding was the significant difference in PRKCZ, ABL1 and FGF4 in patients who died compared with those who survived. These findings after validation in larger cohorts could provide a panel of biomarkers for prognosis of OSCC.

Using array CGH O'Regan et al. (2006) demonstrated less genomic instability in young nonsmokers with OSCC than found in typical patients (smokers) with oral cancer. Bremmer et al. (2005) developed a non-invasive genetic screening test for detection of OPLs based on multiplex ligation-dependent probe amplification (MLPA). MLPA enabled measurement of gains and losses at 40 different chromosomal locations in a single PCR reaction using only 150ng DNA obtained by brushing the oral mucosa. Validation of this method on large sample size could be a promising method for screening of OPLs. Transcriptional profiling in OPLs is crucial for revealing molecular and biological changes resulting from oral epithelial transformation. Some studies such as the reports of Banerjee et al. (2003, 2005) highlighted dysregulation of several genes related to arachidonic acid metabolism and related inflammatory cascades suggesting that this pathway may be involved in the irreversible transition to progressive stages of oral cancer. While the other studies by Warner et al. (2004)

identified distinct patterns of gene expression associated with clinical parameters in OSCCs.

Irie' T et al. (2004) showed that genes which are important for histogenesis and acquisition of invasion and proliferation capability are commonly aberrant in OSCCs. Noteworthy studies by Roepman et al. (2005) and O'Donnell et al. (2005) independently identified and validated primary tumor tissue expression signatures that can detect the presence of local lymph node metastasis reliably. In contrast, in the recent study by Braakhuis et al. (2006) no specific metastasis signature was identified in HNSCC including OSCC. These conflicting reports may reflect heterogeneity of HNSCC with respect to metastatic behavior or may be attributed to the relatively small patient groups used in this study. In conclusion though signatures of differentially expressed genes have been identified it is perhaps too early to state whether a useful signature exists. The analysis of much larger numbers of HNSCCs will be required to validate if a specific molecular signature exists for distant metastasis. DNA hybridization arrays have been used for identifying novel gene clusters predictive of radiation responsiveness of OSCC (Hanna et al. 2001). Using DNA microarrays, Higo et al. (2005) identified a panel of 7 candidate radioresistant genes (CK18, DTNBP1, ASNA1, Tep20, Cyclophilin F, K1Aa0218 and HBp17) that could be useful in identifying radioresistant OSCCs.

Oncogene expressing human papilloma virus type 16 (HPV16) is found in a subset of HNSCC including OSCC. Smeets et al. (2006) used microarray CGH to compare genome wide DNA copy number alterations in HNSCC with or without HPV. Four regions (losses at 3p11.2-26.3, 5q11.2-35.2, and 9p21.1-2y and gain at 11q12.1-13.4) showed alterations in HPV negative tumors that were absent in HPV positive tumors. However, 7 regions were altered at high frequency (>33%) in both groups. This study suggests that HNSCC including OSCC arising by environmental carcinogens are characterized by genetic alterations that differ from those observed in HPV-16 induced tumors and most likely occur in early stages of carcinogenesis, while numerous genetic changes that are shared in both tumor groups may be considered crucial in the later stages of tumor progression.

Intense efforts are being directed towards analysis of DNA and RNA in body fluids such as

serum, plasma, saliva or urine, using array technologies. Human DNA biomarkers have been identified in saliva and used for oral cancer detection (Liao et al. 2000; El Naggar et al. 2001). An excellent example is the noteworthy study by Li et al. (2004) on salivary transcriptome analysis for oral cancer detection. Microarray analysis showed that 1679 genes exhibited significantly different expression level in saliva between OSCC patients and controls. Seven salivary RNAs combined were able to predict the presence of oral cancer with 91% specificity and sensitivity. These potential salivary RNA biomarkers are transcripts of IL8, IL1B, DUS, HA3, OAZ1, S100P and SAT. The utility of salivary transcriptome diagnostics for oral cancer detection as a non invasive tool with sufficient predictive power has been demonstrated in this study. This novel clinical approach could be exploited as a high throughput tool for early cancer detection if validated in other independent studies. Some of these have been confirmed in a later study in 272 OSCC patients (Brinkman and Wong 2006).

High density oligonucleotide microarrays have recently been used for the first time by Li et al. (2006) for global transcriptome profiling from serum for OSCC detection. Human serum circulating mRNAs were amplified by RTPCR and microarray identified 335 serum RNAs to be significantly differentially expressed in OSCC patients as compared to the controls. Five cancer related gene transcripts (H3F3A, TPT1, FTH1, NCOA4 and ARCR) were validated giving a sensitivity of 91% and specificity of 71% in distinguishing OSCC.

These studies illustrate that expression profiling using microarrays is a means to refine conventional assessment of OSCCs allowing for a more accurate prediction of disease prognosis. There is a need for integrated microarray database the would allow comparison of microarray data generated from different laboratories. Molecular signatures of OSCC will ultimate translate into early detection, more predictable response to therapy and development of novel molecular therapeutics.

The ultimate goal of oral cancer proteomics is to adapt proteomic technologies for routine use in clinical laboratories for diagnostic and prognostic use, as well as in evaluating response to therapy including drug toxicity and efficacy. In addition, proteome analysis also enables evaluation of regulation of protein function by

proteolysis, recycling and isolation of subcellular compartments that effect gene products. Finally, protein-protein interactions and the molecular composition of cellular structures can also be determined only at the protein level. However, the dynamic nature of the proteome and the immense biological variability among the patient samples are major challenges for deducing diagnostic patterns unique to OSCC. The current biomarker identification strategies for HNSCC, different combinations of MS platforms, LCM and 2D gel electrophoresis procedures applied to readily available clinical specimens (tissue, blood and saliva) have been recently reviewed by Drake et al. (2005). These authors have also discussed issues related to assay reproducibility, management of large data sets and future improvements in clinical proteomics. The same considerations hold for OSCC, therefore these issues will not be further addressed in this review.

Proteomics Signatures in Oral Cancer

In comparison with gene expression profiling of OSCCs using microarrays, the studies on proteome analysis of oral cancer are limiting. The first proteomic analysis of oral tongue carcinoma was reported by He et al. (2004). Protein expression profiles of 10 tumors and their matched normal mucosa resection margins were analyzed by two dimensional gel electrophoresis and matrix-assisted laser desorption / ionization-time of flight mass spectrometry (MALDI-MS). Several tumor associated proteins including HSP60, HSP27, alpha-B-crystalline, ATP synthase beta, calgranulin B, myosin, tropomyosin and galectin 1 were found to be significantly altered in tongue carcinoma tissues compared with their normal mucosa, providing a proteomic signature of tongue carcinoma. Using the same technique, his laboratory also provided the proteomic signatures of buccal squamous cell carcinoma (Chen et al. 2004). These authors observed increased expression of glycolytic enzymes, heat shock proteins, tumor antigens, cytoskeletal proteins, enzymes involved in detoxification and anti-oxidation systems, and proteins involved in mitochondrial and intracellular signaling pathways. These findings indicate that multiple cellular pathways are involved in oral tumorigenesis. Hence multiple proteins need to be simultaneously targeted for rational design of diagnostic and therapeutic strategies. The panel

of candidates for targeted proteins included SCC antigen, G protein, glutathione S-transferase, manganese superoxide dismutase, annexins, voltage dependent anion channel, cyclophilin A, stratifin and galectin 7, although no specific biomarker for buccal SCC was found in this study.

To gain insight into the molecular mechanisms involved in oral carcinogenesis Koike et al. (2005) used fluorescent two dimensional differences in gel electrophoresis and MALDI TOF-MS for proteomic profiling of human normal oral keratinocytes and OSCC derived cell lines. Nine proteins were up-regulated and 13 were down-regulated in OSCC cells. These included annexin A1, HSP27, laminin A/C, interleukin/ receptor antagonist, serine proteinase inhibitor clade B5, stathmin 1 and superoxide dismutase 2. Maddola et al. (2005) have developed MALDI Imaging MS proteome analysis technique to be used on thin histological samples and a specific statistical method of analysis to enable data processing in the absence of internal standards by defining similarity scores. However utilization of this technique in clinical specimens remains to be tested.

Differentially expressed metastasis-associated proteins in adenoid cystic carcinoma cell lines of human salivary gland have been identified using 2D-MS. Some of the proteins associated with tumor metastasis included transketolase, Dim1p, v-Ha-ras oncogene, type I collagen protein alpha, tumor necrosis factor (ligand) superfamily member 4 and pirin. However, the functions of these proteins in tumor metastasis remain to be determined (An et al., 2004).

Nakashima et al. (2006) identified maspin and stathmin as potential biomarkers and also markers of biological behaviour of adenoid cystic carcinoma of the salivary glands using fluorescent-2-dimensional differential in gel electrophoresis (2D-DIGE)-MALDI-TOF-MS. 2D-DIGE is an emerging technique for comparative proteomics that improves the reproducibility and reliability of differential protein expression analysis between the samples. Fluorescent 2-D differential in-gel electrophoresis (2-D-DIGE) allows multiplex analysis of 3 sample proteomes on the same gel. (Tonge et al. 2001; Skynner et al. 2002; Zhou et al. 2002). The protein extracts being compared are covalently labeled with different fluorescent Cy Dyes, which are N-hydroxysuccinimidyl ester derivatives of Cy2, Cy3 and Cy5. Fluorescent imaging of the gel at the wavelengths specific to each Cy Dye generates images that can be

overlaid directly by the DeCyder software to identify any differentially expressed proteins between the samples.

Cheng et al. (2005) developed a bead based affinity fractionated proteomic method to discover a novel plasma marker for oral cancer. Affinity purification of heparinized plasma with copper chelated beads and MALDI-TOFMS analysis were used to screen potential oral cancer markers from 57 OSCC patients and 29 healthy controls. The spectra were analysed using flex analysis™ and Clin-Prot™ bioinformatic softwares. Six markers that differentiated between cancer and control spectra were found with mean molecular masses of 2664, 2850, 3250, 7735, 7927 and 9240 Da. The marker identified as a fragment of the fibrinogen a-chain had the highest sensitivity (100%) and specificity (97%) for cancer, suggesting that it may be a clinically useful tumor marker. However, clinical studies on a larger sample size are needed to determine whether this biomarker can be used for diagnostic purposes.

Affinity bead-based purification has been developed to reduce costs and make proteomic procedures suitable for general MS analysis. This method uses different chemical chromatographic surfaces on an outer layer of magnetic beads to selectively purify certain subsets of proteins, allowing unbound impurities to be removed by washing with buffers. Protein bound to the magnetic beads are then eluted, diluted, and directly analyzed by MALDI-TOF MS.

Serum proteomics and peptidomics are gaining popularity among oral oncologists in their quest for cancer biomarkers with high diagnostic accuracy. This is based on the premise that sera from oral cancer patients contain small proteins and peptides detectable by mass spectrometry (MS) that reflects the presence and biology of a tumor.

There are very limited studies comparing transcriptome and proteome analysis. In this context the noteworthy study by Roesch Ely and coworkers (2005) combined cDNA microarrays, quantitative reverse transcriptase-PCR (qRT-PCR) and surface enhanced laser desorption ionization (SELDI)-TOF MS to study differential gene expression in HNSCC. Calgranulins A and B and Annexins 1 and 2 were found to be down-regulated in HNSCC compared with normal mucosa, at both the mRNA and protein level, and these findings were confirmed by validation in HNSCCs. Hypoxia decreases therapeutic

efficacy in solid tumors. Therefore identification of hypoxia markers may influence the choice of therapeutic modality. With this rationale Chen et al. (2004) using 2 D-gel electrophoresis and Power Blot (antibody- based array), identified a group of 20 proteins upregulated during hypoxia. These proteins included I Kappa B kinase beta (IKK beta), MKK3b, highly expressed in cancer (HEC), density regulated protein 1, P150 (glued), nuclear transport factor 2, binder of ARL 2, Poxillin, and transcription termination factor 1. Correlation of IKK beta expression in HNSCC specimens with tumor oxygenation suggested that IKK beta may be a novel endogenous marker of tumor hypoxia and may represent a novel target for therapy. Melle et al. (2004) using a technical tirade of tissue microdissection (LCM), protein Chip arrays and immunohistochemistry for proteomic identification of HNSCC showed calgranulin A (S100A8) and calgranulin B (S100A9) to be candidate biomarkers for HNSCC. Wu et al. (2002) identified metastasis associated proteins in HNSCC cell lines using SELDI Protein chip technology – MS. Enolase-alpha, annexin I and annexin II were found to be upregulated and may be important in head and neck cancer invasion and metastasis.

Clinical Validation of Biomarkers

Despite of large amount of data obtained from several expression profiling studies worldwide, there is a gap between the knowledge accrued in the laboratory and its translation into clinical practice. Validation of the data poses a major challenge. Most studies have been carried out on a small sample sizes and often involved cancers from different anatomic sites. Therefore, there is a need for large scale validation in prospective, well designed clinical studies in multicentric settings. The Early Detection Research Network by National Cancer Institute aims to rapidly bring molecular markers to the clinic. Similar initiatives by other countries would considerably aid the quest for validation of candidate novel biomarkers for translation from bench to bedside.

REFERENCES

- An J, Sun JY, Yuan Q, Tian HY, Qiu WL, Guo W, Zhao FK 2004. Proteomics analysis of differentially expressed metastasis-associated proteins in adenoid cystic carcinoma cell lines of human salivary gland. *Oral Oncol*, **40**: 400-408.
- Anderson NL, Anderson NG 2002. The human plasma proteome: history, character, and diagnostic prospects. *Mol Cell Proteomics*, **1**: 845-867. Erratum in: *Mol Cell Proteomics*. 2003 Jan; **2**(1): 50.
- Arora S, Kaur J, Sharma C, Mathur M, Bahadur S, Shukla NK, Deo SV, Ralhan R. 2005. Stromelysin 3, Ets-1, and vascular endothelial growth factor expression in oral precancerous and cancerous lesions: correlation with microvessel density, progression, and prognosis. *Clin Cancer Res*, **11**: 2272-2284.
- Arora S, Matta A, Shukla NK, Deo SV, Ralhan R 2005. Identification of differentially expressed genes in oral squamous cell carcinoma. *Mol Carcinog*, **42**: 97-108.
- Baldwin C, Garnis C, Zhang L, Rosin MP, Lam WL 2005. Multiple microalterations detected at high frequency in oral cancer. *Cancer Res*, **65**: 7561-7567.
- Banerjee AG, Bhattacharyya I, Vishwanatha JK 2005. Identification of genes and molecular pathways involved in the progression of premalignant oral epithelia. *Mol Cancer Ther*, **4**: 865-875.
- Banks RE, Dunn MJ, Hochstrasser DF, Sanchez JC, Blackstock W, Pappin DJ, Selby PJ 2000. Proteomics: new perspectives, new biomedical opportunities. *Lancet*, **356**: 1749-1756.
- Barker PE 2003. Cancer biomarker validation: Standards and process: Roles for the National Institute of Standards and Technology (NIST). *Ann N Y Acad Sci*, **983**: 142-150.
- Belbin TJ, Singh B, Smith RV, Socci ND, Wreesmann VB, Sanchez-Carbajo M, Masterson J, Patel S, Cordon-Cardo C, Prystowsky MB, Childs G 2005. Molecular profiling of tumor progression in head and neck cancer. *Arch Otolaryngol Head Neck Surg*, **131**: 10-18.
- Ben-Izhak O, Kablan F, Laster Z, Nagler RM 2005. Oropharyngeal cancer pathogenesis: ubiquitin proteolytic, apoptotic and epidermal growth factor related pathways act in concert—first report. *Oral Oncol*, **41**: 851-860.
- Bettendorf O, Piffko J, Bankfalvi A 2004. Prognostic and predictive factors in oral squamous cell cancer: important tools for planning individual therapy? *Oral Oncol*, **40**: 110-9.
- Braakhuis BJ, Brakenhoff RH, Leemans CR 2005. Head and neck cancer: molecular carcinogenesis. *Ann Oncol*, **16 Suppl 2:ii**: 249-250.
- Braakhuis BJ, Senft A, de Bree R, de Vries J, Ylstra B, Cloos J, Kuik DJ, Leemans CR, Brakenhoff RH 2006. Expression profiling and prediction of distant metastases in head and neck squamous cell carcinoma. *J Clin Pathol*, [Epub ahead of print]
- Bremmer JF, Braakhuis BJ, Ruijter-Schippers HJ, Brink A, Duarte HM, Kuik DJ, Bloemena E, Leemans CR, van der Waal I, Brakenhoff RH 2005. A noninvasive genetic screening test to detect oral preneoplastic lesions. *Lab Invest*, **85**: 1481-1488.
- Brinkman BM, Wong DT 2006. Disease mechanism and biomarkers of oral squamous cell carcinoma. *Curr Opin Oncol*, **18**: 228-233.
- Bunn PJ Jr. 2003. Early detection of lung cancer using serum RNA or DNA markers: Ready for “prime time” or for validation? *J Clin Oncol*, **21**: 3891-93.
- Carinci F, Lo Muzio L, Piattelli A, Rubini C, Chiesa F,

- Ionna F, Palmieri A, Maiorano E, Pastore A, Laino G, Dolci M, Pezzetti F 2005. Potential markers of tongue tumor progression selected by cDNA microarray. *Int J Immunopathol Pharmacol*, **18**: 513-524.
- Chapman K 2002. The ProteinChip Biomarker System from Ciphergen Biosystems: a novel proteomics platform for rapid biomarker discovery and validation. *Biochem Soc Trans*, **30**: 82-87.
- Chen J, He QY, Yuen AP, Chiu JF 2004. Proteomics of buccal squamous cell carcinoma: the involvement of multiple pathways in tumorigenesis. *Proteomics*, **4**: 2465-2475.
- Chen Y, Shi G, Xia W, Kong C, Zhao S, Gaw AF, Chen EY, Yang GP, Giaccia AJ, Le QT, Koong AC 2004. Identification of hypoxia-regulated proteins in head and neck cancer by proteomic and tissue array profiling. *Cancer Res*, **64**: 7302-7310.
- Cheng AJ, Chen LC, Chien KY, Chen YJ, Chang JT, Wang HM, Liao CT, Chen IH 2005. Oral cancer plasma tumor marker identified with bead-based affinity-fractionated proteomic technology. *Clin Chem*, **51**: 2236-2244.
- Chimenos-Kustner E, Font-Costa I, Lopez-Lopez J 2004. Oral cancer risk and molecular markers. *Med Oral Patol Oral Cir Bucal*, **9**: 381-4; 377-80.
- Conrads TP, Zhou M, Petricoin EF 3rd, Liotta L, Veenstra TD 2003. Cancer diagnosis using proteomic patterns. *Expert Rev Mol Diagn*, **3**: 411-20.
- Dallas PB, Gottardo NG, Firth MJ, Beesley AH, Hoffmann K, Terry PA, Freitas JR, Boag JM, Cummings AJ, Kees UR 2005. Gene expression levels assessed by oligonucleotide microarray analysis and quantitative real-time RT-PCR-how well do they correlate? *BMC Genomics*, **6**: 59.
- Diamandis EP 2004. Mass spectrometry as a diagnostic and a cancer biomarker discovery tool: opportunities and potential limitations. *Mol Cell Proteomics*, **3**: 367-378.
- Drake RR, Cazare LH, Semmes OJ, Wadsworth JT 2005. Serum, salivary and tissue proteomics for discovery of biomarkers for head and neck cancers. *Expert Rev Mol Diagn*, **5**: 93-100.
- El-Naggar AK, Mao L, Staerkel G, Coombes MM, Tucker SL, Luna MA, Clayman GL, Lippman S, Goepfert H 2001. Genetic heterogeneity in saliva from patients with oral squamous carcinomas: implications in molecular diagnosis and screening. *J Mol Diagn*, **3**: 164-170.
- Giri U, Ashorn CL, Ramdas L, Stivers DN, Coombes K, El-Naggar AK, Ang KK, Story MD 2006. Molecular signatures associated with clinical outcome in patients with high-risk head-and-neck squamous cell carcinoma treated by surgery and radiation. *Int J Radiat Oncol Biol Phys*, **64**: 670-677.
- Ha PK, Califano JA 2003. The molecular biology of mucosal field cancerization of the head and neck. *Crit Rev Oral Bio Med*, **14**: 363-369.
- Hanna E, Shrieve DC, Ratanatharathorn V, Xia X, Breau R, Suen J, Li S 2001. A novel alternative approach for prediction of radiation response of squamous cell carcinoma of head and neck. *Cancer Res*, **61**: 2376-2380.
- He QY, Chen J, Kung HF, Yuen AP, Chiu JF 2004. Identification of tumor-associated proteins in oral tongue squamous cell carcinoma by proteomics. *Proteomics*, **4**: 271-278.
- Holmstrup P, Vedtofte P, Reibel J, Stoltze K 2006. Long-term treatment outcome of oral premalignant lesions. *Oral Oncol*, **42**: 461-474.
- Hunter KD, Parkinson EK, Harrison PR 2005. Profiling early head and neck cancer. *Nat Rev Cancer*, **5**: 127-135.
- Hwang D, Alevizos I, Schmitt WA, Misra J, Ohyama H, Todd R, Mahadevappa M, Warrington JA, Stephanopoulos G, Wong DT, Stephanopoulos G 2003. Genomic dissection for characterization of cancerous oral epithelium tissues using transcription profiling. *Oral Oncol*, **39**: 259-268.
- Irie T, Aida T, Tachikawa T 2004. Gene expression profiling of oral squamous cell carcinoma using laser microdissection and cDNA microarray. *Med Electron Microsc*, **37**: 89-96.
- Issaq HJ, Veenstra TD, Conrads TP, Felschow D 2002. The SELDI-TOF MS approach to proteomics: protein profiling and biomarker identification. *Biochem Biophys Res Commun*, **292**: 587-592.
- Kornberg LJ, Villaret D, Popp M, Lui L, McLaren R, Brown H, Cohen D, Yun J, McFadden M 2005. Gene expression profiling in squamous cell carcinoma of the oral cavity shows abnormalities in several signaling pathways. *Laryngoscope*, **115**: 690-698.
- Kozak KR, Amneus MW, Pusey SM, Su F, Luong MN, Luong SA, Reddy ST, Farias-Eisner R 2003. Identification of biomarkers for ovarian cancer using strong anion-exchange ProteinChips: potential use in diagnosis and prognosis. *Proc Natl Acad Sci U S A*, **100**: 12343-12348.
- Kuo WP 2003. Overview of bioinformatics and its application to oral genomics. *Adv Dent Res*, **17**: 89-94.
- Kwong RA, Kalish LH, Nguyen TV, Kench JG, Bova RJ, Cole IE, Musgrove EA, Sutherland RL 2005. p14ARF protein expression is a predictor of both relapse and survival in squamous cell carcinoma of the anterior tongue. *Clin Cancer Res*, **11**: 4107-4116.
- Lewczuk P, Esselmann H, Groemer TW, Bibl M, Maler JM, Steinacker P, Otto M, Kornhuber J, Wiltfang J 2004. Amyloid beta peptides in cerebrospinal fluid as profiled with surface enhanced laser desorption/ionization time-of-flight mass spectrometry: evidence of novel biomarkers in Alzheimer's disease. *Biol Psychiatry*, **55**: 524-530.
- Li J, Zhang Z, Rosenzweig J, Wang YY, Chan DW 2002. Proteomics and bioinformatics approaches for identification of serum biomarkers to detect breast cancer. *Clin Chem*, **48**: 1296-1304.
- Li Y, Elashoff D, Oh M, Sinha U, St John MA, Zhou X, Abemayor E, Wong DT 2006. Serum circulating human mRNA profiling and its utility for oral cancer detection. *J Clin Oncol*, **24**: 1754-1760.
- Li Y, St John MA, Zhou X, Kim Y, Sinha U, Jordan RC, Eisele D, Abemayor E, Elashoff D, Park NH, Wong DT 2004. Salivary transcriptome diagnostics for oral cancer detection. *Clin Cancer Res*, **10**: 8442-8450.
- Li Y, Zhou X, St John MA, Wong DT 2004. RNA profiling of cell-free saliva using microarray technology. *J Dent Res*, **83**: 199-203.
- Liao PH, Chang YC, Huang MF, Tai KW, Chou MY

2000. Mutation of p53 gene codon 63 in saliva as a molecular marker for oral squamous cell carcinomas. *Oral Oncol*, **36**: 272-276.
- Lippman SM, Sudbo J, Hong WK 2005. Oral cancer prevention and the evolution of molecular-targeted drug development. *J Clin Oncol*, **23**: 346-56.
- Liu CJ, Lin SC, Chen YJ, Chang KM, Chang KW 2006. Array-comparative genomic hybridization to detect genomewide changes in microdissected primary and metastatic oral squamous cell carcinomas. *Mol Carcinog*, **45**: 721-731.
- Maddalo G, Petrucci F, Iezzi M, Pannellini T, Del Boccio P, Ciavardelli D, Biroccio A, Forli F, Di Ilio C, Ballone E, Urbani A, Federici G 2005. Analytical assessment of MALDI-TOF Imaging Mass Spectrometry on thin histological samples. An insight in proteome investigation. *Clin Chim Acta*, **357**: 210-218.
- Mao L, Hong WK, Papadimitrakopoulou VA 2004. Focus on head and neck cancer. *Cancer Cell*, **5**: 311-316.
- Melle C, Ernst G, Schimmel B, Bleul A, Koscielny S, Wiesner A, Bogumil R, Moller U, Osterloh D, Halbhuber KJ, von Eggeling F 2004. A technical triade for proteomic identification and characterization of cancer biomarkers. *Cancer Res*, **64**: 4099-4104.
- Merchant M, Weinberger SR 2000. Recent advancements in surface-enhanced laser desorption/ionization-time of flight-mass spectrometry. *Electrophoresis*, **21**: 1164-1177.
- Nagpal JK, Das BR 2003. Oral cancer: reviewing the present understanding of its molecular mechanism and exploring the future directions for its effective management. *Oral Oncol*, **39**: 213-21.
- Nakashima D, Uzawa K, Kasamatsu A, Koike H, Endo Y, Saito K, Hashitani S, Numata T, Urade M, Tanzawa H 2006. Protein expression profiling identifies maspin and stathmin as potential biomarkers of adenoid cystic carcinoma of the salivary glands. *Int J Cancer*, **118**: 704-713.
- Ng EK, Tsui NB, Lam NY, Chiu RW, Yu SC, Wong SC, Lo ES, Rainer TH, Johnson PJ, Lo YM 2002. Presence of filterable and nonfilterable mRNA in the plasma of cancer patients and healthy individuals. *Clin Chem*, **48**: 1212-1217.
- Odani T, Ito D, Li MH, Kawamata A, Isobe T, Iwase M, Nagumo M. 2006. Gene expression profiles of oral leukoplakia and carcinoma: genome-wide comparison analysis using oligonucleotide microarray technology. *Int J Oncol*, **28**(3): 619-24.
- O'Donnell RK, Kupferman M, Wei SJ, Singhal S, Weber R, O'Malley B, Cheng Y, Putt M, Feldman M, Ziober B, Muschel RJ 2005. Gene expression signature predicts lymphatic metastasis in squamous cell carcinoma of the oral cavity. *Oncogene*, **24**: 1244-1251.
- O'Farrell PH 1975. High resolution two-dimensional electrophoresis of proteins. *J Biol Chem*, **250**: 4007-4021.
- O'Regan EM, Toner ME, Smyth PC, Finn SP, Timon C, Cahill S, Flavin R, O'Leary JJ, Sheils O 2006. Distinct array comparative genomic hybridization profiles in oral squamous cell carcinoma occurring in young patients. *Head Neck*, **28**: 330-338.
- Otero-Rey E, Garcia-Garcia A, Barros-Angueira F, Torres-Espanol M, Gandara-Rey JM, Somoza-Martin M 2004. DNA microarrays in oral cancer. *Med Oral*, **9**: 288-292.
- Petricoin EF, Ardekani AM, Hitt BA, Levine PJ, Fusaro VA, Steinberg SM, Mills GB, Simone C, Fishman DA, Kohn EC, Liotta LA 2002. Use of proteomic patterns in serum to identify ovarian cancer. *Lancet*, **359**: 572-577.
- Ralhan R, Chakravarti N, Kaur J, Sharma C, Kumar A, Mathur M, Bahadur S, Shukla NK, Deo SV 2006. Clinical significance of altered expression of retinoid receptors in oral precancerous and cancerous lesions: relationship with cell cycle regulators. *Int J Cancer*, **118**: 1077-1089.
- Reibel J 2003. Prognosis of oral pre-malignant lesions: significance of clinical, histopathological, and molecular biological characteristics. *Crit Rev Oral Biol Med*, **14**: 47-62.
- Roepman P, Wessels LF, Kettelarij N, Kemmeren P, Miles AJ, Lijnzaad P, Tilanus MG, Kooze R, Hordijk GJ, van der Vliet PC, Reinders MJ, Slootweg PJ, Holstege FC 2005. An expression profile for diagnosis of lymph node metastases from primary head and neck squamous cell carcinomas. *Nat Genet*, **37**: 182-6.
- Roesch Ely M, Nees M, Karsai S, Magele I, Bogumil R, Vorderwulbecke S, Ruess A, Dietz A, Schnolzer M, Bosch FX 2005. Transcript and proteome analysis reveals reduced expression of calgranulins in head and neck squamous cell carcinoma. *Eur J Cell Biol*, **84**: 431-444.
- Rogers MA, Clarke P, Noble J, Munro NP, Paul A, Selby PJ, Banks RE 2003. Proteomic profiling of urinary proteins in renal cancer by surface enhanced laser desorption ionization and neural-network analysis: identification of key issues affecting potential clinical utility. *Cancer Res*, **63**: 6971-6983.
- Schliephake H 2003. Prognostic relevance of molecular markers of oral cancer—a review. *Int J Oral Maxillofac Surg*, **32**: 233-245.
- Scully C, Porter S 2000a. ABC of oral health. Oral cancer. *BMJ*, **321**: 97-100.
- Scully C, Porter S 2000b. ABC of oral health. Swellings and red, white, and pigmented lesions. *BMJ*, **321**: 225-228.
- Seliger B, Kellner R 2002. Design of proteome-based studies in combination with serology for the identification of biomarkers and novel targets. *Proteomics*, **2**: 1641-1651.
- Sidransky D 1997. Nucleic acid-based methods for the detection of cancer. *Science*, **278**: 1054-1059.
- Skyner HA, Rosahl TW, Knowles MR, Salim K, Reid L, Cuthliff R, McAllister G, Guest PC 2002. Alterations of stress related proteins in genetically altered mice revealed by two-dimensional differential in-gel electrophoresis analysis. *Proteomics*, **2**: 1018-1025.
- Snijders AM, Schmidt BL, Fridlyand J, Dekker N, Pinkel D, Jordan RC, Albertson DG 2005. Rare amplicons implicate frequent deregulation of cell fate specification pathways in oral squamous cell carcinoma. *Oncogene*, **24**: 4232-4242.
- Soni S, Kaur J, Kumar A, Chakravarti N, Mathur M, Bahadur S, Shukla NK, Deo SV, Ralhan R 2005. Alterations of rb pathway components are frequent events in patients with oral epithelial dysplasia and

- predict clinical outcome in patients with squamous cell carcinoma. *Oncology*, **68**: 314-325.
- St John MA, Li Y, Zhou X, Denny P, Ho CM, Montemagno C, Shi W, Qi F, Wu B, Sinha U, Jordan R, Wolinsky L, Park NH, Liu H, Abemayor E, Wong DT 2004. Interleukin 6 and interleukin 8 as potential biomarkers for oral cavity and oropharyngeal squamous cell carcinoma. *Arch Otolaryngol Head Neck Surg*, **130**: 929-935.
- Steel LF, Shumpert D, Trotter M, Seeholzer SH, Evans AA, London WT, Dwek R, Block TM 2003. A strategy for the comparative analysis of serum proteomes for the discovery of biomarkers for hepatocellular carcinoma. *Proteomics*, **3**: 601-609.
- Tonge R, Shaw J, Middleton B, Rowlinson R, Rayner S, Young J, Pognan F, Hawkins E, Currie I, Davison M 2001. Validation and development of fluorescence two-dimensional differential gel electrophoresis proteomics technology. *Proteomics*, **1**: 377-396.
- Vigneswaran N, Beckers S, Waigel S, Mensah J, Wu J, Mo J, Fleisher KE, Bouquot J, Sacks PG, Zacharias W 2006. Increased EMMPRIN (CD 147) expression during oral carcinogenesis. *Exp Mol Pathol*, **80**: 147-159.
- Villanueva J, Philip J, Chaparro CA, Li Y, Toledo-Crow R, DeNoyer L, Fleisher M, Robbins RJ, Tempst P 2005. Correcting common errors in identifying cancer-specific serum peptide signatures. *J Proteome Res*, **4**: 1060-1072.
- Wang BG, Huang HY, Chen YC, Bristow RE, Kassaei K, Cheng CC, Roden R, Sokoll LJ, Chan DW, Shih IeM 2003. Increased plasma DNA integrity in cancer patients. *Cancer Res*, **63**: 3966-3968.
- Warnakulasuriya KAAS, Ralhan R 2006. Clinical, pathological, cellular and molecular lesions caused by oral smokeless tobacco. *J Oral Pathol Med*, REVIEW, In Press.
- Warner GC, Reis PP, Jurisica I, Sultan M, Arora S, Macmillan C, Makitie AA, Grenman R, Reid N, Sukhai M, Freeman J, Gullane P, Irish J, Kamel-Reid S 2004. Molecular classification of oral cancer by cDNA microarrays identifies overexpressed genes correlated with nodal metastasis. *Int J Cancer*, **110**: 857-868.
- Wu W, Tang X, Hu W, Lotan R, Hong WK, Mao L 2002. Identification and validation of metastasis-associated proteins in head and neck cancer cell lines by two-dimensional electrophoresis and mass spectrometry. *Clin Exp Metastasis*, **19**: 319-326.
- Zerhouni EA 2005. Translational and clinical science-time for a new vision. *N Engl J Med*, **353**: 1621-1623.
- Zhou G, Li H, DeCamp D, Chen S, Shu H, Gong Y, Flaig M, Gillespie JW, Hu N, Taylor PR, Emmert-Buck MR, Liotta LA, Petricoin EF 3rd, Zhao Y 2002. 2D differential in-gel electrophoresis for the identification of esophageal cancer cell cancer-specific protein markers. *Mol Cell Proteomics*, **1**: 117-124.