Distribution of ABO and Rh(D) Blood Groups among Koari (Backward Caste) Population of Jaunpur District

Pradeep Kumar¹, Sarita Maurya² and Vandana Rai¹

¹Department of Biotechnology, ²Department of Applied Biochemistry, V B S Purvanchal University, Jaunpur 222 001, Uttar Pradesh, India

KEYWORDS Blood Groups. Koari. Variation.

ABSTRACT The distribution of ABO blood groups and Rh (D) factor has been studied among the Koari caste group population of Jaunpur (U.P.). The O, A, B, and AB blood group percentage were recorded as 34.5%, 24.6 %, 33% and 7.9% respectively. The allele frequencies of *O*, *A* and *B* were found to be 0.5898, 0.1787, and 0.2315 respectively and Rh (D) allelic frequency was 0.901.

INTRODUCTION

ABO blood group genes are mapped at 9q34.2. ABO and Rh(D) blood groups varies among different geographical and ethnic populations. Since its discovery in 1900 by Landsteiner, ABO typing is the single most important test performed in blood banking services to avoid mortality. During the last few decades, several studies have been carried out on the distribution of blood groups among several endogamous populations in India (Bhasin et al. 1992, 1994). In this paper an attempt is made to study the distribution of ABO and Rh(D) blood group systems among the Koari population of Jaunpur district.

MATERIALS AND METHODS

The blood samples were drawn from randomly selected 203 unrelated individuals of Koari caste (of both sexes) from Jaunpur district of Uttar

Pradesh. The samples were analyzed for the ABO and Rh (D) blood groups using standard antisera (Bhasin and Chahal 1996). Gene frequencies were calculated according to the method of Maurant et al. (1976).

RESULTS AND DISCUSSION

The frequency distribution of ABO phenotypes with gene frequencies in Koari population in Jaunpur district is presented in table 1. Phenotype O has the highest frequency (34.5%) followed by B (33.0%), A (24.6%) and AB (7.9%). The overall picture of phenotypic frequencies of ABO blood groups is O>B>A>AB. The decreasing order of allele frequency in Koari is *O* (0.5898)> *B* (0.2315) and *A* (0.1787). Out of the 203 subjects tested, 201(99.01%) were found to be Rh positive and 2(0.99%) were Rh negative. The frequency of D and d alleles is 0.901 and 0.099 respectively (Table 2).

Table 1: Distribution of the ABO blood group and their allele frequencies among the Koari.

Phenotype	Observed Number	%	Expected Number	Allele Frequency
O	70	34.5	70.6	A= 0.1787
A	50	24.6	49.3	B= 0.2315
B	67	33.0	66.3	O= 0.5898
AB	16	7.9	16.8	
Total	203	100.0	203.0	1.0000

Corresponding Author:

Pradeep Kumar, Lecturer
Department of Biotechnology,
VBS Purvanchal University, Jaunpur-222001 (U.P.)
Telephone: 05452252538(O), 09451713485(M)
E-mail: pradipk14@yahoo.co.in
pradeepk14@rediffmail.com

Table 2: Distribution of the Rh(D) blood group and their allele frequencies among the Koari.

Phenotype	Observed Number	%	Allele Frequency
Rh(D)Positive	201	99.01%	0.901
Rh(d)Negative	2	0.99%	0.099
Total	203	100.00	1.0000

ACKNOWLEDGEMENT

The authors would like to thank Dr. V. S. Upadhyay and his pathologist Mr. Pradeep for their generous help in blood grouping. We are very much grateful to Koari community of Jaunpur district because without their help and support the present study was not possible.

REFERENCES

- Bhasin MK, Chahal SMS 1996. *A Laboratory Manual for Human Blood Analysis*. Delhi: Kamla-Raj Enterprises.
- Bhasin MK, Walter H, Danker-Hopfe H 1992. *The Distribution of Genetical Morphological and Behavioural Traits among the Peoples of Indian Region*. Delhi: Kamla-Raj Enterprises.
- Bhasin MK, Walter H, Danker-Hopfe H 1994. *People of India: An Investigation of Biological Variability in Ecological, Ethno-economic and Linguistic Groups*. Delhi: Kamla-Raj Enterprises.
- Mourant A E Kopec AC Domaniewska-Sobczak K 1976. *The Distribution of the Human Blood Groups and Other Polymorphisms*. 2nd Edition, London: Oxford University Press.