



© *Kamla-Raj* 2004

PRINT: ISSN 0972-3757 ONLINE: 2456-6360

Int J Hum Genet, 4(4): 257-260 (2004)

DOI: 10.31901/24566330.2004/04.04.06

**Comparative Study of Exfoliated Oral Mucosal Cell Micronuclei
Frequency in Normal, Precancerous and Malignant Epithelium**

**A. Halder¹, T. Chakraborty¹, K. Mandal², P. K. Gure², S. Das³,
R. Raychowdhury², A. K. Ghosh³ and M. De¹**

*1. Department of Genetics, Vivekananda Institute of Medical Sciences, Ramakrishna
Mission Seva Pratisthan, 99, Sarat Bose Road, Kolkata 700 026, West Bengal, India*

2. ENT, Department RKMS, VIMS

3. Department of Oral and Maxillo facial Surgery

KEYWORDS Micronuclei; precancerous lesions; buccal smear

ABSTRACT Analysis of exfoliated buccal cell micronuclei (MN) is a sensitive method of monitoring genetic damage in the human population. In the present study 50 patients with precancerous or malignant oral epithelial lesions from the Departments of Facio-maxillary and ENT Surgery of RKMS Hospital, Kolkata were compared with 50 age and sex matched healthy controls without any oral lesions. The MN frequency was increased in preoperative cancer cases and decreased in postoperative cases, while in pre cancerous cases it was higher than in the controls.

[Home](#)

[Back](#)
