

Genetics of Castes and Tribes of India: Somatometry

M. K. Bhasin

Department of Anthropology, University of Delhi, Delhi 110 007, India

KEYWORDS Anthropometry; measurements and indices; biological variation; ethnic groups

ABSTRACT The discipline of human population biology incorporates study of biology and environmental factors, as well as the forces of micro-evolution leading to macro-evolution which ultimately influence the biological structure of human populations. The unit of study in understanding variations in man is a 'breeding population' some times also referred to as 'Mendelian population'. India is inhabited by people of great diversity, different creeds and customs forming what may be designated as multiple (or plural) society. There are about 3000 castes in India, some have genesis in tribal stock while others are occupational, linguistic, religious and territorial entities. Each caste is a social unit or what may be called 'monopolistic guild' in itself. All these groups are not entirely independent; usually people belong to two or more of such groups at the same time. India is a country with distinct geographical entity and is marked off from rest of Asia by both mountains and sea. Indian sub-continent may be divided into four natural regions: (1) The Himalayan Mountain Complex, (2) The Indus-Ganga-Brahmaputra Plains, (3) The Peninsular Plateau and (4) The Islands. The climate of India has many regional variations determined by locations, altitude, distance from the sea or the mountains and the general relief. India is divided into eight climatic regions based on the monthly value of temperature and precipitation. India is a Union comprising 25 States and 7 Union Territories and these may be categorised into six zones (North, West, East, Central, South India and Islands). Himalayan region may be divided into three divisions (Western, Central and Eastern Himalayan Regions). The present paper aims at investigating first the variation in the people of India in *regional* (Natural Regions of India, Climatic Regions of India, Political Divisions of India), *ethnic* (castes, scheduled castes, scheduled tribes, communities), *traditional occupational* and *linguistic groups* and *families* with the help somatometry measurement and indices. Second, to study the variation with the help of biostatistics methods in the region, ethnic groups and linguistic groups. The basic data were collected from the literature and it was categorized in regional, ethnic, occupational and linguistic groups and coded accordingly for the analysis on computer.

INTRODUCTION

Biological relationships and distances between human individuals and groups can be assessed by the use of anthropometric data at least as successfully as this can be done by the use of serological traits with known modes of inheritance (Spielman and Smouse 1976). The obvious reason is that anthropometric dimensions are also genetically determined even if the polygenic nature of the genetic control and the environmental effect on the development of their phenotypic manifestation cannot be spelled out in detail (Brace and Hunt 1990; Brace et al. 1991).

Although the effects of physical environment such as temperature, rainfall, humidity, altitude etc. upon human physique have been widely discussed, from the available literature yet it is difficult to offer definitive explanations in terms of the causations of variations in body dimensions. As it appears, the variations may be caused by complex interactions of physical environmental factors with others *e.g.* biological

(diseases—degenerative and contagious), nutritional, economic and cultural (behavioural norms and practices relating to habitation, food, diseases etc.) factors.

The environmental changes brought about by the factor of 'migration' of the original migrants and their progeny may be responsible for measurable variations that have been observed. The findings of many investigations are available in which attempt has been made to isolate the effect of dietary differences and climatic changes upon human physique.

Shapiro (1939) found that the Hawaiian-born children of Japanese-born immigrants are considerably taller than their parents, but their heads are significantly shorter and broader and thus they have greater stature with pronounced brachycephaly.

Mills from his studies on the effects of climate on body growth and development revealed that temperate climate is more favourable for growth and development than the tropics (Mills 1937) and that dietary modifications and altered vitamin intake in the tropics affect adult dimensions (Mills 1942).

Kaplan (1954) has reviewed 25 studies and

observed that most of the investigators assume that bodily changes which they report are not evolutionary in nature, although several of them refer to long-range secular changes in size. She pointed out that in fact the changes are not evolutionary, but merely individual responses to the particular environment and it should be possible to find the precise facet or facets of the environment responsible for the changes observed. She found that dietary changes are most frequently mentioned as important and concluded one wonders, to what extent eating and living habits, improvements in housing and sanitary conveniences, greater attention to personal hygiene, shorter school and working days, or even psychological factors play their part.

Tobias (1985) discussed the concept of secular trend *i.e.* long-term systematic or non-random changes in a wide variety of traits in successive generations of a population living in the same territories (Wolanski 1966, 1967). He added that it is probably analytically more useful to distinguish between secular changes with a territorial restriction, and the effects of migration in general and of urbanization as a special case. He argued that the direction of the secular processes should always be specified *i.e.* first upward or positive secular trend, second negative secular trend and third absence of secular changes, positive or negative. On the basis of secular trend data, Tobias suggested a four-fold subdivision of the world's populace taking socio-economic status into consideration:

1. Have most (affluent) with absence of secular trend at present as they have reached upper phenotypic limit set by their genetic potential.
2. Have ample (less affluent) with positive secular trend at present as they are moving from middle to upper reaches of genetically determined growth rate and body size range.
3. Have little (pastoral/agriculture), absent or negative secular trend as they are moving from middle to lower reaches of genetically determined range of growth rates and body sizes.
4. Have least (hunters and gatherers) positive secular trend as they are rising from the lowest reaches of genetic potential towards the middle of the range of growth rates and body sizes.

Tobias pointed out that the biggest challenge facing planners is how to solve the problem of transforming the negative secular trend of the "has little" sector to the positive secular trend of the "have ample" sectors.

The anthropometric studies on the Indian populations started in 1868. However systematic surveys were carried on All India level by Risley (1915), v. Eickstedt (1934) and Guha (1935). The main emphasis of these studies was to classify the peoples of India into various 'Racial Types'. They recognized the existence of more than one 'Racial Type' and a great deal of anthropometric heterogeneity among the peoples of India. Later, a number of surveys were conducted to study the regional variations for example among the people of Gujarat (Majumdar and Sen 1949), Uttar Pradesh (Mahalanobis et al. 1949), Maharashtra (Karve and Dandekar 1951), Bengal (Majumdar and Rao 1960), Tamil Nadu (Malhotra et al. 1981), among others. Besides these a good number of other studies are available, and about 1200 population groups reported by different authors are listed by Bhasin et al. (1992).

There are a rather high number of anthropometric studies on populations of India. Unfortunately, technical differences exist, and therefore sometimes the results are not comparable. It has been observed that some of the measurements on face taken by different authors show variation in dimensions which obviously reflect technical deviations, probably in the determination of nasion and other landmarks. Moreover, the samples are not distributed homogeneously. Only a few studies are available from Central India, whereas rests of the zones are better represented. For the techniques, methods and analysis of the data, readers are referred to Martin and Saller (1957), Wiener and Lourie (1969, 1982), Lohman et al. (1988), Knussmann (1988) and Singh and Bhasin (1989, 2004).

The aim of the study is to have a satisfactory knowledge of micro-evolutionary processes as they are reflected in genetic and morphological traits in human populations. This variability has been studied in terms of natural regions, climatic factors, climatic regions, political division of India, ethnic groups, traditional occupations and linguistic groups (For details see Bhasin 1988; Bhasin et al. 1992, 1994; Bhasin and Walter 2001, Bhasin 2006a).

The studies on Genetics of Castes and Tribes have been reported on Serum Protein Polymorphisms KM and GM Systems (Bhasin and Walter 2002); Glucose-6-Phosphate Dehydrogenase Deficiency and Haemoglobin variants (Bhasin 2006b); Red and Green Colour

Vision Deficiency (Bhasin, 2006c); Taste Sensitivity (Bhasin 2006d) among the people of India and factors responsible to operate in favour of these traits.

IDENTIFY AND DISTINGUISH THE PEOPLE

For the biogenetical study of the people of India, researchers have generally used the following criteria to identify and distinguish the people: 1. Regional Groups, 2. Ethnic Groups, 3. Linguistic Groups, and 4. Religious Groups (For details see Bhasin et al., 1994; Bhasin and Walter 2001; Bhasin 2006)

It should, however, be kept in mind that these are the convenient units of study, although there are significant levels of overlapping between them. For example, an occupational group pursuing traditional job inhabits a region, shares religion with other categories, belongs to one or the other language group and has an aggregation of ethnic properties. But in the human population genetic studies, out of these criteria one is chosen (Bhasin 1988).

In the present study an attempt has been made to analyse the above mentioned biogenetical traits into 1. Regional Groups, 2. Ethnic Groups, 3. Traditional Occupational Groups and 4. Linguistic Groups.

I. Regional Groups

These can be divided into the following groups:

1. Natural Regions of India
2. Climatological Factors and Climatic Regions of India
3. Political Division of India

Each region has its own characteristics and a brief description of each one will give an idea of what it constitutes of.

1. Natural Regions of India

The natural regions have broad uniformity in their characteristics, such as relief, geomorphological history, drainage, climate, soil, natural vegetation and wild life. Broadly speaking the Indian sub-continent may be divided in the following natural regions:

1. The Himalayan Mountain Complex
2. The Indus-Ganga-Brahmaputra Plain

3. The Peninsular Plateau and
4. The Islands

2. Climatological Factors and Climatic Regions of India

Various climatological factors (Rainfall, Humidity, Temperature) and Altitude have been considered to study correlations with different biological traits. The values for the climatological factors are after "Climatological Tables of Observations in 1931-1960" Meteorological Department, Government of India, New Delhi.

A climatic region generally possesses a broad uniformity in climatic conditions produced by combined effects of climatic factors. India can be divided into the following climatic regions after Köppen's method, based on the monthly values of temperature and precipitation:

- (1) Tropical Savannah Type, (2) Monsoon Type with Short Dry Season, (3) Monsoon Type with Dry Season in High Sun Period, (4) Semiarid and Steppe Climate, (5) Hot Desert Type, (6) Monsoon Type with Dry Winters, (7) Cold Humid Winters Type with Shorter Summer, and (8) Polar Type.

3. Political Division of India

India is a Union of States. Comprising 25 States and 7 Union Territories, according to the Census 1991, there are 4689 towns and 587,226 inhabited and 47,095 uninhabited villages in the country. The country had 466 districts in 1991.

India-Political and Ethnic Zones: The weighted mean values of various biological traits have been classified into 25 States and 7 Union Territories (U.T.) which have been categorised as follows (after Bhasin, 1988):

I. North India, II. West India, III. East India, IV. Central India, V. South India, and VI. Islands. Himalayan Region may be divided into three divisions, i.e., A) Western Himalaya, B) Central Himalaya, and C) Eastern Himalaya as follows:

I. North India:

(A) *Western Himalaya* (S. No. 1, 2):

- (1) Jammu and Kashmir, (2) Himachal Pradesh, (3) Punjab, (4) Chandigarh (U.T.), (5) Haryana, (6) Delhi (U.T.), (7) Uttar Pradesh

(B) *Central Himalaya* (S.No.7, Eight Districts of Uttar Pradesh)

- [(i) Almora, (ii) Chamoli, (iii) Dehra Dun, (iv) Garhwal (Pauri), (v) Naini Tal, (vi) Pithoragarh, (vii) Tehri Garhwal, and (viii) Uttarkashi.] and 8. Rajasthan

II. West India:

(1) Gujarat, (2) Maharashtra, (3) Goa², (4) Daman and Diu² (U.T.) and (5) Dadra and Nagar Haveli (U.T.)

III. East India:

C) Eastern Himalaya: (S. No.1 to 8 and Darjeeling District of West Bengal)

(1) Arunachal Pradesh, (2) Assam, (3) Nagaland, (4) Manipur, (5) Mizoram, (6) Tripura, (7) Meghalaya, (8) Sikkim, (9) West Bengal, (10) Bihar, and (11) Orissa

IV. Central India:

(1) Madhya Pradesh

V. South India:

(1) Karnataka, (2) Andhra Pradesh, (3) Tamil Nadu, (4) Kerala and (5) Pondicherry (U.T.).

VI. Islands:

(1) Lakshadweep (U.T.) and (2) Andaman and Nicobar Islands (U.T.).

II. Ethnic Groups

An aggregation of biological and socio-cultural characteristics constitutes an ethnic group. Within the category of Ethnic Group, we include Castes, Scheduled Castes, Scheduled Tribes and Communities (the names of Scheduled Castes and Scheduled Tribes after Manual of Election Law 1982, Government of India, New Delhi). Biological anthropological studies of such ethnic groups as well as "Communities" have been reported in India. By Community we generally refer to a group of people who may have occupational, linguistic, religious or regional characteristics (Bhasin 1988).

III. Traditional Occupational Groups

In the traditional society, there were occupational guilds. The *Chaturvarna* system with its division into *Brahman* (priestly caste), *Kshatriya* (warrior caste), *Vaishya* (land owners and traders) and *Sudra* (labouring caste) was based on occupational differentiation. The occupations are grade manual labour is looked down upon, and those dealing with swine-herding, scavenging, butchery, removal of night soil are regarded as polluting (Bhasin 1988). The caste based division of occupation is 1. Priesthood, 2. Warfare, 3. Trade and Commerce, 4. Agriculture, 5. Animal Husbandry, 6. Artisan and 7. Menial Workers.

IV. Linguistic Groups

Although the Schedule VIII recognizes fifteen languages in India, there are innumerable dialects which change after few scores of kilometers. Linguistic diversity is an important factor in the formation of regional groups, and it also reflects the regional differentiation. The four-fold regional division can be seen for the major languages, i.e., (i) the Dravidian region of the south; (ii) the Indo-Aryan regions of the north and north-west; (iii) the Mon Khmer and the Tibeto-Burman region of the north-east and the Himalayan region; and (iv) the Austric region of the Aravalli-Vindhya-Chota Nagpur complex. These languages are again divided into sub-families and groups as follows:

Language Classification—Indian Languages

The family, branch, group and language of India are as follows:

I. AUSTRO-ASIATIC FAMILY

Mon-Khmer Group (Mon-Khmer Branch)

Munda Group (Munda Branch)

II. TIBETO-CHINESE FAMILY

SIAMESE-CHINESE SUB-FAMILY

Tai Group

TIBETO-BURMAN SUB-FAMILY

Tibeto-Himalayan Branch

Bhotia Group (Tibetan Group)

Himalayan Group (Pronominalized/Non-Pronominalized Himalayan Group)

North-East Frontier Group (North Assam Branch)

Assam-Burmese Branch

Bodo Group (Bara or Bodo Group)

Naga Group

Kachin Group

Kuki-Chin Group

Burma Group

III. DRAVIDIAN FAMILY

South Dravidian Group

Central Dravidian Group

North Dravidian Group

IV. INDO-EUROPEAN FAMILY ARYAN SUB-FAMILY

Dradic (or Pisacha) Branch

Kafir Group

Khovar Group

Dard Group

Indo-Aryan Branch:

Outer Sub-Branch

North-Western Group

Southern Group
 Eastern Group
 Bihari Group (Bhojpuri, Maithili & Magahi
 Sub-Group)
Mediate Sub-Branch/Inner Sub-Branch:
 Mediate Group/Central Group/Pahari Group
 Mediate Group/Central Group
 Pahari Group
 Eastern Pahari
 Central Pahari
 Western Pahari

Mean Weighted Values

To discern the pattern of regional groups, ethnic groups, traditional occupational groups and linguistic groups using the frequency data, the mean weighted values of the biological traits have been calculated and estimates for the various groups are presented.

1. MEASUREMENT

1.1. Stature

The mean value of stature among Indian populations is 163.06 cm (medium), which varies from 143.50 cm - very short (among Naya Kurumba of Tamil Nadu) to 181.90 cm - very tall (in Mohammedans from West Punjab). The value of stature is low (159.75 cm - short) among scheduled tribes as compared to other ethnic groups. In the natural regions, populations from Himalayan mountain complex are showing lower values (162.13 cm) as compared to other regions except from Islands from where different scheduled tribes are reported (Table 1), (Bhasin et al. 1994; Bhasin and Walter 2001). The dimension at Himalayan mountain complex is in conformity with Bergmann's and Allen's rule, which states that in warm-blooded animals body size, and consequently surface area/weight, as also the length of the extremities, are smaller in colder than in warmer climates to minimize heat loss. Similar observations have been reported by Harrison et al. (1969) in their Ethiopian study. Attention should also be given to the ideas of Roberts (Roberts 1953, 1973) who asserts that these significant relationships between the temperature of biome and body size have both direct (response to temperature) and/or indirect (acting through food, genetics etc.) relationship. However, from the available literature, it may not be possible to

offer definitive explanations in terms of causations by altitudinal variations (as they reflect temperature values) as it appears that these variations may be caused by complex interactions of physical environment factors with other biological, socio-economic and cultural factors (e.g. Walter 1982).

The mean value of stature is highest from semi arid steppe type climatic region (165.39 cm) and lowest from monsoon type with short dry season region (156.80 cm), followed by cold humid winters with short summers (159.05 cm) whereas from monsoon type with dry season, tropical savannah type and monsoon type with dry winters region the values show less differences (163.07, 163.06 and 163.66 cm, respectively). Takahashi (1971) also reported geographical gradients of stature with respect to climatic environmental variables and observed that taller statures are found in the arid belt of the northern and north-western sectors of the subcontinent while shorter statures occur in the humid regions of the eastern and southern zones. There is a gradient of stature distribution from north-west to east and south positions of the Indian subcontinent.

The inhabitants of Punjab, Delhi and Rajasthan are in general of taller stature (168.85, 168.57, 168.69 cm, respectively) than the populations of other regions of India, a feature especially marked among castes as compared to other ethnic groups (Table 1, Fig. 1). From the states of Jammu and Kashmir and Himachal Pradesh of Western Himalayan region and from Central Himalayan region the mean values are almost similar (163.38, 163.13 cm, respectively). In the state of Jammu and Kashmir, Kashmiri Pandits and Muslims are taller (medium) as compared to Ladakhis (from high altitude) with Mongoloid affinities (lower medium, 160 - 163.9 cm), and also populations from Himachal Pradesh. The populations studied from Lahaul-Spiti and Kinnaur districts (middle to high altitude) with Mongoloid affinities are showing low medium height as compared to other populations and similar results have been observed from Central Himalayan region from where also populations with Mongoloid affinities living at middle to high altitude are showing lower medium stature (different groups of Bhotias). From the Eastern Himalayan region, the mean values of stature are quite low from all the states [Arunachal Pradesh (159.05 cm), Assam (162.04 cm), Nagaland (161.10

cm), Manipur (160.02 cm), Mizoram (163.26 cm), Tripura (160.17 cm). Meghalaya (158.50 cm), Sikkim (159.97 cm)] inhabited by populations with Mongoloid admixture of varying degrees at different altitudes (Table 1, Fig. 1). Among the different ethnic groups from the Himalayan region the mean value of stature is less than 164 (lower medium) with little differences, except from Eastern region where scheduled tribes are having low value (160.42 cm) as compared to other ethnic groups. From Nepal also the various populations having Mongoloid admixture in varying degrees and living in middle to low altitudes (Rai, Tamang, Sunwar, Newars, Magars, Gurungs, Gurkhas among others) show short to low medium stature.

As compared to the North Indian populations with taller stature (medium category) the values of height among populations inhabiting different states of West India, Orissa, Madhya Pradesh and States and Union Territories of South India

fall in the category lower medium whereas Islanders have a short stature. Takahashi (1971) attributes this cline to differential diet noting that the western and northern inhabitants of South Asia include many herding groups who consume high quantities of dairy food. Milk products are high in protein relative to grains that make up diet of Indians living in eastern and southern habitats. Among the different ethnic groups, peoples belonging to castes are taller followed by communities as compared to scheduled castes (lower medium) and scheduled tribes (short stature). Caste groups from North India and East India zones are taller as compared to other zones, whereas tribes from South India and East India zones are short in stature (157.69 and 159.62 cm, respectively) as compared to other zones, and they are falling in lower medium category. Mahalanobis (1927) and Olivier (1963) also reported that caste status was positively




Fig. 1. Mean values of stature (in cm) in different regions of India (after Bhasin and Walter 2001)

correlated to stature.

From islands, among isolates very short stature (130.0 to 149.9 cm) is observed among Onge and Andamanese tribals and short to lower medium stature in Nicobarese tribals (Table 1, Fig. 1). The Andamanese and Onges are considered to have Negrito affinities and Nicobarese indicate their Malayan ancestry (Dutta 1978).

As observed, caste populations are on an average taller as compared to scheduled castes. This may be due to genetic differences between these groups since they belong to different racial affinities—Caucasoid (Aryan and/or Dravidian) in caste populations and Caucasoid and/or Australoid (Dravidian and/or Pre-Dravidian) among scheduled caste people. Further there is a difference between socio-economic conditions between the two groups. Therefore the taller stature may be largely but not solely attributable to genetic differences. The tribals who are considered to be autochthonous have had a longer stay in their habitat as compared to other ethnic groups. The short stature among tribal groups suggests in particular that the basic substratum of ethnic elements have been in favour of short individuals. In view of the advantages of the smaller body size in both 'undernourished' and 'hot climate' the smaller stature among the tribals might have developed in the process of adaptation to these conditions.

The mean values of stature correlations with various climatic factors and altitude by different ethnic groups although showing significant differences are not high except in scheduled tribes, where there are negative correlations of stature with climatic factors and altitude (Table 6).

For assessment of the environment - body dimension relationship, Basu et al. (1980) looked at the correlations between stature, cephalic index and nasal index on one hand and environmental variables like rainfall and temperature on the other. They found that none of them is significant except the correlation between stature and rainfall in the all India sample, and also that this correlation disappears when they restrict analysis to a relatively less heterogeneous group *i.e.* Brahmans. From this finding they pointed out that the relevant environmental factors are not the climatic ones they have considered but others *e.g.* social (*i.e.* castes/class), economic, nutritional etc.

To study the correlations between stature and caste status, Mahalanobis (1927) ranked castes

in a hierarchical order and showed positive correlation between caste status and stature and sought to interpret these relations in terms of successively greater intrusion of Caucasoid genes into the higher castes through admixture. Olivier (1963) observed similar results on data from Tamil Nadu. Basu et al. (1980) stated that the immensely elaborate and rigid systems of social stratification, in which each strata was virtually closed to fresh intrusion at an unknown antiquity, seem to be a plausible enough determinant of anthropometric variations in India. They added that for this hypothesis to assume general validity, findings such as of Mahalanobis (1927) and Olivier (1963) should be repeated from diverse Indian populations, which is however not the case. They argued that for instance while an anthropometric survey conducted in the state of Uttar Pradesh (Mahalanobis et al. 1949) shows that castes of comparable hierarchical status tend to cluster together, a similar survey in Bengal shows that the clusters are not necessarily composed of castes of similar social status (Majumdar and Rao 1960). Further, the nature of biological similarity/differences among castes/subcastes of similar and different hierarchical status lead Karve and Malhotra (1968) to conclude that hierarchical affinity does not often correspond to biological similarity. Also Rakshit's (1966) study showing wide diversity among the Brahmans also points in the same direction. They concluded that it seems that both the hypotheses (a) of fission of an ancestral social group into castes/sub-castes, each of the latter having more biological similarities with the other than with sub-castes of other castes and (b) of coming together of social groups of disparate origins without fusion of their marital boundaries to constitute caste clusters/castes, are probable but not sufficient from the point of view of explaining the immense biological variations among the Indian population. Further in the few regional studies in which clusters have been worked out or at least distance matrices have been computed (Mahalanobis et al. 1949; Majumdar and Rao 1960; Karve and Malhotra 1968), the similarities/differences do not seem to follow any definite pattern. Mahalanobis et al.'s (1949) study does show some clusters which correspond to traditional caste affinities, but even then there are frequent exceptions. Further, they considered the tribal populations only and observed that variations among conventional racial categories

Table 1: Stature (in cm)

Particulars	Subjects studied		No. of studies	Stature		
				Mean	Min	Max
1. NATURAL REGION						
Himalayan Mountain Complex	8446	122		162.13	154.90	169.80
Indus-Ganga-Brahmaputra Plains	27932	300		164.15	154.80	181.90
Peninsular Plateau	49124	623		162.77	143.50	175.00
Islands	1124	18		155.50	146.80	161.93
2. CLIMATIC REGION						
Monsoon Type with Short Dry Season	5026	83		156.80	146.80	172.50
Monsoon Type with Dry Season	2727	59		163.07	154.91	174.00
Tropical Savannah Type	36813	379		163.06	147.52	174.00
Semi Arid Steppe Type	5015	110		165.39	143.50	175.00
Hot Desert Type		0				
Monsoon Type with Dry Winters	35986	414		163.66	154.80	181.90
Cold Humid Winters with Short Summers	338	5		159.05	157.71	160.86
Polar Type	723	13		162.49	159.45	164.54
3. POLITICAL DIVISION OF INDIA						
I. NORTH INDIA						
<i>A. Western Himalaya (S. No. 1, 2)</i>						
1. Jammu and Kashmir	989	17		163.84	159.96	166.65
2. Himachal Pradesh	2695	28		163.21	159.45	165.40
3. Punjab	1077	15		168.85	159.78	181.90
4. Chandigarh UT	-	0		-	-	-
5. Haryana	-	0		-	-	-
6. Delhi UT	742	10		168.57	162.59	175.50
<i>B. Central Himalaya (S. No. 7, Eight Districts of Uttar Pradesh)</i>						
7. Uttar Pradesh	14833	161		164.07	156.21	171.91
8. Rajasthan	1698	11		168.69	158.80	174.80
II. WEST INDIA						
9. Gujarat	4956	49		163.05	159.02	169.51
10. Maharashtra	7696	130		163.11	158.40	174.00
11. Goa, Daman and Diu UT	-	0		-	-	-
12. Dadra and Nagar Haveli UT	-	0		-	-	-
III. EAST INDIA						
<i>C. Eastern Himalaya (S. No. 13 to 20 and Darjeeling District of West Bengal)</i>						
13. Arunachal Pradesh	338	5		159.05	157.71	160.86
14. Assam	4888	70		162.04	154.80	165.08
15. Nagaland	1021	18		161.10	155.87	165.84
16. Manipur	297	6		160.02	154.90	167.40
17. Mizoram	40	1		163.26	163.26	163.26
18. Tripura	311	5		160.97	155.76	160.90
19. Meghalaya	487	8		158.50	156.91	161.26
20. Sikkim	797	16		159.97	155.76	164.45
21. West Bengal	7822	63		164.11	157.00	169.80

Particulars	Subjects studied		No. of studies	Stature		
				Mean	Min	Max
22. Bihar	17048	84		164.07	147.52	171.38
23. Orissa	3884	54		160.28	153.64	166.30
IV. CENTRAL INDIA						
24. Madhya Pradesh	2578	37		160.32	156.10	164.36
V. SOUTH INDIA						
25. Karnataka	2193	49		163.68	153.13	168.70
26. Andhra Pradesh	1421	34		162.98	159.10	165.80
27. Tamil Nadu	2437	93		163.73	143.50	174.00
28. Kerala	4106	71		157.38	151.50	175.00
29. Pondicherry UT	1150	10		162.80	159.95	165.37
VI. ISLANDS						
30. Lakshadweep UT	-	0		-	-	-
31. Andman Islands and Nicobar Islands UT	400	11		148.45	146.80	151.80
	724	7		159.39	156.70	161.93
3A. ZONES OF INDIA						
I. North India						
	22034	242		164.70	156.21	181.90
II. West India						
	12652	179		163.09	158.40	174.00
III. East India						
	36933	330		163.05	147.52	171.38
IV. Central India						
	2578	37		160.32	156.10	164.36
V. South India						
	11307	257		161.23	143.50	175.00
VI. Islands						
	1124	18		155.50	146.80	161.93
INDIA (TOTAL)						
	86628	1063		163.06	143.50	181.90
3B. REGIONS OF HIMALAYA						
A. Western Himalaya						
	3684	45		163.38	159.45	166.65
B. Central Himalaya						
	1394	17		163.13	156.21	166.53
C. Eastern Himalaya						
	8745	138		161.20	154.80	169.80
HIMALAYA (TOTAL)						
	13823	200		161.97	154.80	169.80
NON-HIMALAYAN REGIONS						
	72805	863		163.27	143.50	181.90
4. ETHNIC GROUP						
A. ZONES OF INDIA						
I. NORTH INDIA						
Caste	6801	82		165.69	159.71	174.80
Scheduled Caste	3563	48		163.30	158.17	167.47
Scheduled Tribe	2799	29		163.15	156.21	171.37
Community	8871	83		164.99	159.78	181.90
II. WEST INDIA						
Caste	5401	86		163.48	159.86	167.95
Scheduled Caste	562	9		161.64	158.42	163.68
Scheduled Tribe	2340	28		161.28	158.40	168.22

Table 1: Contd.....

Particulars	Subjects studied	No. of studies	Mean	Stature Min	Stature Max
Community	4349	56	164.76	159.63	174.00
III. EAST INDIA					
Caste	16671	75	165.97	156.20	171.38
Scheduled Caste	3982	42	161.55	157.70	165.05
Scheduled Tribe	12776	168	147.52	150.10	167.40
Community	3504	45	163.42	155.76	169.80
IV. CENTRAL INDIA					
Caste	104	2	162.57	160.91	164.36
Scheduled Caste	-	0	-	-	-
Scheduled Tribe	2424	34	160.27	156.10	163.64
Community	50	1	158.17	158.17	158.17
V. SOUTH INDIA					
Caste	3591	80	163.79	155.60	170.10
Scheduled Caste	1474	34	160.91	152.53	166.55
Scheduled Tribe	3827	80	157.69	143.50	175.00
Community	2415	63	163.21	153.13	174.00
VI. ISLANDS					
Caste	-	0	-	-	-
Scheduled Caste	-	0	-	-	-
Scheduled Tribe	1124	18	155.50	146.80	161.93
Community	-	0	-	-	-
INDIA					
Caste	32568	325	165.25	155.60	174.80
Scheduled Caste	9581	133	162.11	152.53	167.47
Scheduled Tribe	25290	357	159.75	143.50	175.00
Community	19189	248	164.18	153.13	181.90
B. REGIONS OF HIMALAYA					
A. Western Himalaya					
Caste	839	9	163.62	160.74	165.40
Scheduled Caste	75	2	162.31	162.11	165.08
Scheduled Tribe	1470	14	163.30	159.45	164.75
Community	1300	20	163.38	159.96	166.65
B. Central Himalaya					
Caste	60	2	160.85	160.80	160.89
Scheduled Caste	213	2	166.05	165.50	166.53
Scheduled Tribe	536	6	161.35	156.21	163.33
Community	585	7	163.93	161.61	164.90
C. Eastern Himalaya					
Caste	243	5	163.21	161.61	164.61
Scheduled Caste	1221	14	162.77	158.67	165.05
Scheduled Tribe	5785	92	160.42	154.80	167.40
Community	1496	27	162.61	155.76	169.80
5. TRADITIONAL OCCUPATION					
Priesthood	15726	78	165.64	159.86	168.50
Particulars	Subjects studied	No. of studies	Mean	Stature Min	Stature Max
Warfare	3452	31	166.96	160.44	174.80
Trade and Commerce	2383	29	165.47	161.20	169.48
Agriculture	1957	40	163.52	159.10	169.72
Animal Husbandry	880	12	167.17	161.37	175.50
Artisans	1463	20	162.79	159.90	166.37
Menial Workers	6571	72	161.66	152.53	166.33
No Information	54196	781	162.05	143.50	181.90
6. LANGUAGE GROUP					
I. AUSTRAL-ASIATIC FAMILY					
Mon Khmer Group	1185	13	158.78	156.70	161.93
Munda Group	6379	65	158.91	147.52	164.40
II. TIBETO-CHINESE FAMILY					
(i) Siamese-Chinese Sub-Family					
Tai Group	85	1	159.90	159.90	159.90
(ii) Tibeto-Burman Sub-Family					
Bhotia Group	2285	27	162.90	156.21	164.90
Himalayan Group	1039	20	159.69	156.36	169.80
North East Frontier Group	1141	20	160.80	157.52	165.03
Bodo Group	1985	25	161.48	154.80	164.30
Naga Group	1228	24	160.79	155.70	167.40
Kachin Group	13	1	160.50	160.50	160.50
Kuki Chin Group	403	9	160.12	154.90	163.60
III. DRAVIDIAN FAMILY					
South Dravidian Group	11074	253	161.14	143.50	175.00
Central Dravidian Group	2215	34	160.69	156.67	162.89
North Dravidian Group	3008	11	162.58	157.52	163.40
IV. INDO-EUROPEAN FAMILY					
Dard Group	572	6	164.08	162.11	166.00
North Western Group	30	1	165.81	165.81	165.81
Southern Group	7818	124	163.04	158.40	174.00
Eastern Group	2025	36	160.92	155.14	166.30
Bihari	21075	111	165.35	156.20	171.38
Central Group	21246	249	164.67	158.17	181.90
Pahari Group	955	16	163.11	155.76	165.40
Unspecified	256	2	162.84	161.40	163.33
Other Languages	638	15	154.69	146.80	168.20
No Information	-	0	-	-	-
6A. LANGUAGE FAMILY					
I. Austro Asiatic Family	7564	78	158.89	147.52	164.40
II. Tibeto Chinese Family	8152	127	161.37	154.80	169.80
III. Dravidian Family	16297	298	161.34	143.50	175.00
IV. Indo European Family	53977	545	164.52	155.14	181.90
Other Languages	638	15	154.69	146.80	168.20
No Information	-	0	-	-	-

i.e., Australoid, Mongoloid, Mundari and Negroid are not significant and concluded that neither caste as such nor race, therefore, seem to be determinant of inter group anthropometric variability.

The correlation between nutritional factors and body size showed positive correlation between calorie intake and stature, fat intake and height and fat intake and weight, although no such correlation between protein intake and height or weight seems to exist (Basu et al., 1980). The protein-deprived individuals fail to attain their full ontogenetic growth potential and under these conditions of chronic deficiency stress smaller body size (of which stature is a critical component) may be highly adaptive for survival. Natural selection for reduction of body size and mass may continue in a population existing under these nutritional conditions which may persist over many generations. However the genotype is not altered and realization of full ontogenetic development may be attained in high proportion of the population when adequate protein is introduced into the diet.

Malhotra (1966) also demonstrated the effect of nutrition on height and weight. Considering that nutritional level is strongly influenced by economic condition, the source of much of the variations that are traditionally ascribed to racial properties can be traced to much less abstruse socio-economic variables.

A number of small scale studies from all over the world suggest that growth and maturation of children and adult body dimensions *e.g.*, height, weight etc. are positively related to urbanization, social, economic and professional status and family size. Basu et al. (1980) reported that Ganguly conducted a study in 1974 taking these factors into consideration and observed that rural/urban habitat does not have any effect on any of the anthropometric traits he considered and the well to do section of the population differs significantly from the poorer section almost in every character. They pointed that the weakness of Ganguly's (1979) study lies in the fact that the probable effects of occupational and sexual selection and nutrition on body dimensions have not been tested by any rigorous method (Basu et al. 1980). The effect of nutrition on height, weight and/or other measures of child growth and adult body dimensions have been more amply demonstrated by Pachauri et al. (1971), Gopalan et al. (1973), Easwaran et al. (1974), Rao and

Satyanarayana (1974), Rao et al. (1975), Vijayalakshmi and Devaki (1976), Devdas et al. (1977) among others.

Among the higher occupational groups—priesthood, warfare, trade and commerce, the populations are taller (165.64, 166.96 and 165.47 cm, respectively) as compared to lower groups—artisans and menial groups (162.79 and 161.66 cm, respectively). Mahalanobis (1927) and Olivier (1963) reported positive correlations between stature and caste status. Whereas among pastoralists the value is maximum as compared to other groups (167.17 cm), which may be explained due to selection as suggested by Wolpoff (1980). He suggests that tallness was adaptive for prehistoric people, who were nomadic, travelling long distances in pursuit of game which migrated seasonally to different ecozones. The physical demands of the hunting-gathering life style which involved tracking of large and dangerous game placed a selective advantage on high muscular-skeletal robusticity and large body size, particularly among males who were involved in procurement of flesh foods under conditions of high risk and strenuous demands upon physical strength and stamina. Selection for large and robust body form was relaxed among sedentary food producing populations where the physical demands and risks are of a very different sort (in the present study stature mean value is 163.52 among agriculture group). Pastoralism may involve selection for efficiency of long-distance walking, and the tallest modern South Asians are from the northwest sector of the subcontinent where herding is a common way of life.

Among the speakers of Mon Khmer group (Austro-Asiatic family) and Tibeto-Chinese languages the mean value of stature is low (158.78 and 161.37 cm, respectively) as compared to Indo-European language groups (Dard group - 164.08 and Pahari group - 163.11 cm) from the Himalayan regions among whom most have Mongoloid affinities. Most of the tribals reported in the present study from peninsular region speak Munda group (Austro-Asiatic family) and Dravidian language and among them also the mean values of stature are low, whereas speakers of Indo-European languages (mostly caste groups) show taller stature as compared to the speakers of other languages (164.52 cm) (Table 1, Fig. 1) (Bhasin et al. 1994; Bhasin and Walter, 2001).

Indian populations are on an average medium in stature (163.06 cm) among whom caste people

are taller as compared to scheduled castes whereas tribals are short in stature, suggesting that the basic substratum of ethnic elements has been in favour of short individuals. In view of the advantages of the smaller body size in both 'under-nourished' and 'hot climate' the smaller stature among the tribals might have developed in the process of adaptation to these conditions. The inhabitants of Punjab, Delhi and Rajasthan are in general taller than the populations of other regions, which show a gradient of stature distribution from northwest to east and south positions of Indian subcontinent. Among occupational groups, the taller stature is found among pastoralists who indicate that pastoralism may involve selection for efficiency of a long distance walking. The speakers of Austro-Asiatic, Tibeto-Chinese and Dravidian languages are shorter in stature as compared to Indo-European language speakers.

2. INDICES

2.1. Cephalic Index

The mean value of cephalic index among Indian populations is 76.06, which is falling into the mesocephaly category, and it varies from 67.91 (hyperdolichocephalic from Kerala, South India) to 90.88 (hyperbrachycephalic in West Bengal, East India). The mean value of cephalic index is low among scheduled tribes (75.81 - dolichocephalic) as compared to other ethnic groups. From the natural regions, the population groups of Indus-Ganga-Brahmaputra plains with lower average mean value (75.67 - mesocephalic index) as compared to populations of other natural regions. The cephalic index from monsoon type with short dry season and cold humid winters with short summers climatic region are dolichocephalic as compared to other climatic regions, where mesocephalic index is observed (Table 2) (Bhasin et al. 1994; Bhasin and Walter 2001).

There are 244 studies available from North India and all the populations are showing mean values less than 76 which falls in the dolichocephaly category, except a few populations with Mongoloid affinities from the Himalayan regions—Ladakhi, Changpa, Kashmiri from the state of Jammu and Kashmir; Kanets Lahoul, Keonthali of Mahasi, Bunan and Lahauli of Lahaul, Spiti, Tibetans and Nyamslat of Kinaur

of Himachal Pradesh of Western Himalayan region; Bhotia groups, Tharus, Kewat who are mesocephalics and Burman Mongoloid (83.10 brachycephalic) of Central Himalayan region. The value of index is high among scheduled tribes of North India, who are mostly from Himalayan region with Mongoloid affinities as compared to other ethnic groups from North India. From Nepal, the Old Nepalese and Gurkha are having meso-to brachycephalic index and there is tendency towards higher indices in the East. Among the population groups from Eastern Himalayan region the mean value of cephalic index is high (78.09) as compared to Western and Central Himalayan regions (74.00 and 76.71, respectively). From the states of Arunachal Pradesh, Assam, Nagaland, Manipur, Mizoram, Tripura, Meghalaya and Sikkim, most of the populations studied show mean average value between 76.0 and 80.9 (mesocephaly) except Garo and Abor groups, Kuki Chin, Rengma and Sema Nagas, Kaipeng, Synteng, Pnar Khasi who are falling in dolichocephalic category and Ahom, Assamese Brahmin, Ao Naga, Bhutias, Chhetris (Sikkim), Rais and Limboos who are brachycephalics. From Darjeeling district of West Bengal also brachycephalics are observed among Lepchas, Limbus, Khambus, Gurungs, Bhutias population groups with Mongoloid affinities. From West Bengal in most of the populations the mean value of index is falling in mesocephalic category except a few namely Mal Pharia, Santal, Mala (tribal groups) and Rajbanshi, Barui, Mashiya, Rishi, Mahato (scheduled castes and caste groups) who are dolichocephalics. From West Bengal incidence of mesocephalic is predominant among populations belonging to different ethnic groups and this may be due to Mongoloid racial element present among them in varying degrees. From the states of Bihar and Orissa most of the tribal groups are dolichocephalics as compared to other ethnic groups who are mesocephalics (For details see Bhasin et al. 1992).

From the states of Gujarat and Maharashtra of West India, the mean values of cephalic index are quite high (78.89 and 77.79, respectively). The values are high among castes (78.69) and communities (78.68) as compared to scheduled tribes (76.55). The brachycephalic index is reported among Brahmin Nagar, Parsis, Ahir, Kunbis, Kshatriyas, Prabhu Pathare whereas dolichocephalic index is observed among Dhanuk, different groups of Bhils, Halbi, Govari, Koli,

Katkari, all scheduled tribes.

The mean value of cephalic index is 74.96 among the populations of Central India—who is mostly tribals and speaks Indo-European (Central group), Austro-Asiatic (Munda group) and Dravidian (Central Dravidian) languages and is falling in dolichocephalic category. Similarly from the states of Bihar and Orissa among the tribals the mean value of cephalic index is less than 76 (dolichocephalic) and they belong mainly to the speakers of Austro-Asiatic (Munda group) and Indo-European (Eastern group) in Bihar and Austro-Asiatic (Munda group) and Dravidian (Central Dravidian) from Orissa.

The value of cephalic index is quite high among the populations from the states of Karnataka (78.71), Andhra Pradesh (77.79) as compared to Tamil Nadu (75.71) and Kerala (74.21), from where the mean value is low among

scheduled tribe (74.21 dolichocephaly, varies from 69.44 to 78.90) as compared to other ethnic groups who are mesocephalics and are having almost similar values. The brachycephaly (81.0) is observed among the populations namely—Gauda, Kodagu, Kunchitiga, Brahmin Havig, Pattasali, Gangadikara Vokkaliga from Karnataka state, Badaru and Kamti from Andhra Pradesh, Sukum Sale and Suka Sale from Tamil Nadu.

Onges and Andamanese of Andaman Islands are brachycephalics whereas among Nicobarese the differences are quite wide as observed from dolichocephalic Car Nicobarese to mesocephalic Nicobarese of Choura and Terressa Islands to hyperbrachycephalic Southern Nicobarese (Ganguly 1976). It appears that dolichocephalics are predominant in North, East and towards Southern tip of India, whereas mesocephalics are frequent in West and parts of South India and




Fig. 2. Frequencies of cephalic index in different regions of India (after Bhasin and Walter 2001)

Eastern Himalayan region (Table 2, Fig. 2).

The correlations of mean value of cephalic index with various climatic factors and altitude by different ethnic groups though showing significant differences are not high (Table 6).

The brachycephalic element is observed among the population groups with Mongoloid affinities from the Himalayan region, particularly from Eastern Himalayan region. From West India, brachycephalics are observed among caste groups and a migrated community of Parsis and from the states of West Bengal, Karnataka, Andhra Pradesh and Tamil Nadu also this element is observed among the caste groups. From the Islands, among Onges, Andamanese and Nicobarese-Southern, brachycephaly is predominant. It is interesting to note that brachycephalic element, which is in general quite low among populations of India is observed among populations with Mongoloid

affinities and caste groups of West and South India and state of West Bengal.

Guha in Census of India 1931 has made a coloured map of the distributions of the head form in India to which Keith (1936) has referred as follows:

“Guha’s map shows brachycephaly (red) sweeping southwards rounds both ends of Himalayas. From Pamir it descends through Afghanistan, Baluchistan and Sind and then extends continuously along the west, broadening out from Bombay so far as the whole of Deccan is included. The red band sweeps across the peninsular so as to include the South Madras. Only a small area in South is left as blue (dolichocephalic), it is along the Malabar Coast. From the eastern end of the Himalaya the brachycephalic (red) area passes from Bhutan and Tibet southward through Assam to spread over




Fig. 3. Distribution of type of cephalic index in different regions of India (after Sarkar 1958)

Table 2: Cephalic Index

Particulars	Subjects studied	No. of studies	Cephalic Index			Particulars	Subjects studied	No. of studies	Cephalic Index								
			Mean	Min	Max				Mean	Min	Max						
1. NATURAL REGION																	
Himalayan Mountain Complex	8860	140	76.38	69.29	86.51	22. Bihar	16028	76	75.11	72.40	78.20						
Indus-Ganga-Brahmaputra Plains	27517	294	75.87	70.80	90.88	23. Orissa	3700	53	75.23	72.22	78.40						
Peninsular Plateau	49096	597	76.07	67.91	87.00	IV. CENTRAL INDIA											
Islands	762	14	78.64	74.33	86.11	24. Madhya Pradesh	2525	36	74.96	71.80	79.80						
2. CLIMATIC REGION																	
Monsoon Type with Short Dry Season	4223	72	75.00	67.91	86.11	V. SOUTH INDIA											
Monsoon Type with Dry Season	3000	61	76.16	70.00	80.87	25. Karnataka	2187	48	78.71	73.10	83.00						
Tropical Savannah Type	34622	362	76.23	71.80	87.00	26. Andhra Pradesh	1426	34	77.79	72.89	87.00						
Semi Arid Steppe Type	4945	102	76.00	69.06	83.00	27. Tamil Nadu	2635	90	75.71	69.16	82.20						
Hot Desert Type	-	0	-	-	-	28. Kerala	3665	64	74.01	67.91	78.40						
Monsoon Type with Dry Winters	36574	431	76.01	69.29	90.88	29. Pondicherry UT	1161	8	76.26	72.44	80.87						
Cold Humid Winters with Short Summers	338	5	75.79	73.23	78.06	VI. ISLANDS											
Polar Type	533	12	76.24	73.44	79.56	30. Lakshadweep UT	-	0	-	-	-						
3. POLITICAL DIVISION OF INDIA																	
I. NORTH INDIA																	
A. Western Himalaya (S. No. 1, 2)																	
1. Jammu and Kashmir	1139	24	74.15	69.29	79.56	31. Andman Islands and Nicobar Islands UT	187	8	82.65	81.18	84.80						
2. Himachal Pradesh	2294	24	73.92	70.70	78.54	3A. ZONES OF INDIA											
3. Punjab	1077	15	73.60	72.30	74.40	I. North India	21763	244	73.65	69.29	83.10						
4. Chandigragh UT	-	0	-	-	-	II. West India	11713	172	78.24	72.40	83.76						
5. Haryana	-	0	-	-	-	III. East India	36398	335	76.82	71.75	90.88						
6. Delhi UT	742	10	74.14	72.35	76.00	IV. Central India	2525	36	74.76	71.80	79.80						
B. Central Himalaya (S. No. 7, Eight Districts of Uttar Pradesh)												VI. Islands	762	14	78.64	74.33	86.11
7. Uttar Pradesh	14813	160	73.54	70.80	83.10	INDIA (TOTAL)						84235	1045	76.06	67.91	90.88	
8. Rajasthan	1698	11	73.85	72.40	76.50	3B. REGIONS OF HIMALAYA											
II. WEST INDIA												A. Western Himalaya	3433	48	74.00	69.29	79.56
9. Gujarat	4749	50	78.89	74.90	83.76	B. Central Himalaya	1394	17	76.71	71.13	83.10						
10. Maharashtra	6964	122	77.79	72.40	83.15	C. Eastern Himalaya	9766	153	78.09	73.23	86.51						
11. Goa, Daman and Diu UT	-	0	-	-	-	HIMALAYA (TOTAL)						14593	218	77.00	69.29	86.51	
12. Dadra and Nagar Haveli UT	-	0	-	-	-	NON-HIMALAYAN REGIONS						69642	827	75.86	67.91	90.88	
III. EAST INDIA																	
C. Eastern Himalaya (S. No. 13 to 20 and Darjeeling District of West Bengal)																	
13. Arunachal Pradesh	338	5	75.79	73.23	78.06	4. ETHNIC GROUP											
14. Assam	4846	66	78.11	73.47	82.83	A. ZONES OF INDIA											
15. Nagaland	1700	35	77.83	75.69	81.87	I. NORTH INDIA											
16. Manipur	280	4	75.87	74.19	77.25	Caste	6803	82	73.40	69.88	77.50						
17. Mizoram	40	1	75.15	75.15	75.15	Scheduled Caste	3558	47	73.26	71.00	75.30						
18. Tripura	312	5	76.56	74.98	76.93	Scheduled Tribe	2428	27	75.01	71.14	79.89						
19. Meghalaya	887	12	77.04	73.40	77.81	Community	8974	88	73.62	69.29	83.10						
20. Sikkim	797	16	80.06	76.46	86.51	II. WEST INDIA											
21. West Bengal	7470	62	79.93	71.75	90.88	Caste	4656	78	78.69	74.50	83.76						
						Scheduled Caste	561	9	77.42	76.51	78.64						
						Scheduled Tribe	2107	27	76.55	72.40	80.27						

Table 2: Contd.....

Particulars	Subjects studied	No. of studies	Cephalic Index	
			Mean	Max
Community	4389	58	78.68	83.15
III. EAST INDIA				
Caste	16721	76	76.51	90.88
Scheduled Caste	3927	42	78.03	80.48
Scheduled Tribe	12598	175	76.25	86.51
Community	3152	42	79.23	84.30
IV. CENTRAL INDIA				
Caste	104	2	74.83	75.53
Scheduled Caste	-	0	-	-
Scheduled Tribe	2371	33	74.96	79.80
Community	50	1	75.17	75.17
V. SOUTH INDIA				
Caste	3295	82	76.98	87.00
Scheduled Caste	1337	32	76.50	83.00
Scheduled Tribe	3316	65	74.21	78.90
Community	3126	65	76.89	81.70
VI. ISLANDS				
Caste	-	0	-	-
Scheduled Caste	-	0	-	-
Scheduled Tribe	762	14	78.64	86.11
Community	-	0	-	-
INDIA				
Caste	31579	320	76.20	90.88
Scheduled Caste	9383	130	75.96	83.00
Scheduled Tribe	23582	341	75.81	86.51
Community	19691	254	76.17	84.30
B. REGIONS OF HIMALAYA				
A. Western Himalaya				
Caste	858	10	73.78	77.50
Scheduled Caste	70	1	71.87	71.87
Scheduled Tribe	1098	12	74.43	78.54
Community	1407	25	73.90	79.56
B. Central Himalaya				
Caste	60	2	75.66	76.84
Scheduled Caste	213	2	74.21	73.69
Scheduled Tribe	536	6	77.33	79.89
Community	585	7	77.16	83.10
C. Eastern Himalaya				
Caste	243	5	79.08	81.41
Scheduled Caste	1221	14	78.22	80.12
Scheduled Tribe	6811	108	77.57	86.51
Community	1491	26	80.19	84.30
5. TRADITIONAL OCCUPATION				
Priesthood	15528	76	75.60	81.56
Particulars	Subjects studied	No. of studies	Cephalic Index	
			Mean	Max
Warfare	3491	33	74.45	69.88
Trade and Commerce	2383	29	77.18	71.30
Agriculture	1950	39	75.34	81.40
Animal Husbandry	880	12	73.85	71.87
Artisans	710	11	77.33	80.12
Menial Workers	6409	71	76.27	72.40
No Information	52884	774	76.27	81.28
6. LANGUAGE GROUP				
I. AUSTRAL-ASIATIC FAMILY				
Mon Khmer Group	1436	16	77.34	74.33
Munda Group	5441	58	75.18	86.11
II. TIBETO-CHINESE FAMILY				
(i) Siamese-Chinese Sub-Family	85	1	78.48	71.75
Tai Group				80.48
(ii) Tibeto-Burman Sub-Family				
Bhotia Group	2098	27	77.34	78.48
Himalayan Group	1039	20	79.66	70.38
North East Frontier Group	1111	19	78.11	86.51
Bodo Group	1983	24	76.89	72.98
Naga Group	1901	39	77.90	85.04
Kachin Group	13	1	76.06	73.23
Kuki Chin Group	388	8	75.55	82.83
III. DRAVIDIAN FAMILY				
South Dravidian Group	10836	240	76.06	70.38
Central Dravidian Group	2031	33	75.41	85.04
North Dravidian Group	3009	11	73.67	73.23
IV. INDO-EUROPEAN FAMILY				
Dard Group	652	10	74.05	72.30
North Western Group	30	1	82.91	76.20
Southern Group	7064	115	77.93	82.91
Eastern Group	1892	34	76.20	81.81
Bihari	20718	109	76.94	72.40
Central Group	20853	249	74.49	78.40
Pahari Group	969	16	74.10	72.99
Unspecified	256	2	76.46	90.88
Other Languages	430	12	79.29	70.80
No Information	-	0	-	84.80
6A. LANGUAGE FAMILY				
I. Austro Asiatic Family	6877	74	75.63	71.75
II. Tibeto Chinese Family	8616	139	77.67	86.11
III. Dravidian Family	15876	284	75.52	70.38
IV. Indo European Family	52434	536	75.99	86.51
Other Languages	430	12	79.29	67.91
No Information	-	0	-	87.00

Bengal and to end in Orissa. The actual data on which Guha based his map has not been published. So Guha's map cannot be studied or checked by other researchers".

Sarkar et al. (1955) made an attempt to find out whether brachycephaly is restricted to certain zones or it has sweeping distribution as shown in Guha's map. In their paper, they placed the data into four tables and found the following zones with highest concentration of brachycephaly (Fig. 3).

Zone A: North West Frontier Provinces (NWFR), Punjab, Rajputana, and Southern Extension

Zone B: Himalayan foot hill

Zone C: Chittagao Hill Tracks, Bengal and Assam

Others excluding zones A, B, and C in which states are Uttar Pradesh, Bihar and then from Melghat Orissa, Nilgiri Hills, Travancore, Tinnevalley.

It has been observed that brachycephaly confines more or less to certain zones and these zones are not interrelated. It does not appear as clear sweeping continued and evenly distributed strain as reported by Guha (1931).

In *Zone A*, the highest value is 76.8 per cent of Northwest Frontier Provinces to about 60 per cent in Punjab and Baluchistan and then it falls by 1 to 2 per cent in certain groups. The southward extension of the above is justified because of the absence of the high percentage of brachycephaly similar to N.W.F.P., only in Prabhu highest is 52 per cent which falls to 46 in Nagar Brahmin and 40 per cent in Vадnagar Nagar Brahmin. The other groups of these areas show very low percentage of brachycephaly. They suggested that Pamir have always been recognized as centre of brachycephaly and the same is true for Hindukush region. It would therefore be not out of place to assume that N.W.F.P. brachycephaly has its source in the above region.

In *Zone B*, high brachycephaly has been with Limbu of Nepal (82 per cent). This appears to be of Mongoloid origin.

Zone C has the highest brachycephaly among Chakama (76 per cent). It therefore appears that whatever brachycephaly is found in this area has its origin in the east and this is all the more evident in the absence of any influx of brachycephaly from the *Zone B*.

The latter argument of Sarkar et al. (1955) is born out of the fact that the brachycephaly

appears to decrease appreciably and significantly in the region further from the nucleus. The Kayasthas (Uttar Radhi) of Bengal have only 10 per cent of brachycephaly and it has been found to be very low in the western region of Bengal. They observed that the source of this brachycephaly is Southeast Asia, which is predominantly brachycephalic and closer to Eastern India.

Sarkar et al. (1955) have not grouped in any category the people of Tinnevalley in the southern tip of peninsula among whom they found fairly high brachycephaly (Parwars and Shanars, who show 34.0 ± 6.7 and 51 ± 4.9 per cent of brachycephaly, respectively). Hornwell (1923) who has studied them in course of his research on the origin and ethnological significance of Indian Boat was of the opinion that there have been separate waves migrations from Southeast Asia and the first wave was of Polynesian and second of Malayanesians. He emphasized the similarities between the Sharnas and Malayanesian people of Java. It is therefore not improbable that Malayan element has been responsible for the brachycephaly in East India to a certain extent. (Basu et al. 1980).

Among the higher occupational groups—priesthood and warfare, the mean value of cephalic index is low (75.60 and 74.45) and are dolichocephalics as compared to the artisans and menial workers who are mesocephalics. However among all these occupational groups the differences in the mean values are not high and the values are either dolichocephalic or near the lower range of mesocephaly (76.0 to 80.9), which indicates gene flow among them, as also observed for other biological traits (Table 2) (Bhasin et al. 1994; Bhasin and Walter 2001).

Among the speakers of Munda group of Austro-Asiatic and North and Central Dravidian groups of Dravidian languages, the predominant element present is dolichocephaly and they are mostly tribals. The mean value of cephalic index is high among speakers of Mon Khmer group (Austro-Asiatic family), Bhotia group, Himalayan group, North East Frontier group, Naga group (Tibeto-Chinese family) and Northwestern group (Indo-European family) languages who are having Mongoloid affinities. The speakers of Central group are dolichocephalics as compared to Bihari, Eastern (East India) and Southern groups (Maharashtra) who are mesocephalics (Table 2) (Bhasin et al. 1994; Bhasin and Walter 2001).

Zone	Incidence		
	High	Medium	Low
North India	Dolichocephaly	-	Meso-and Brachycephaly
Central India	Dolichocephaly	Mesocephaly	-
West India	Mesocephaly	-	Brachy-and Dolichocephaly
East India	Mesocephaly	Dolichocephaly	Brachycephaly
South India	Mesocephaly	Dolichocephaly	Brachycephaly
Islands	Brachycephaly	-	Meso-and Dolichocephaly

In India, cephalic index value varies from dolichocephaly to mesocephaly barring a few population groups from West, East, South and Islands among whom brachycephaly has been observed; the mean value of the index is 76.06 which are mesocephaly. In various zones, the distribution of cephalic index is mostly dolichocephaly in North and Central India and brachycephaly is absent; West, East and South India are predominantly mesocephalic with low frequency of brachycephaly, while the brachycephaly element is prominent from Islands. In the Hiamalayan region the low mean value of cephalic index is observed from Western region which is dolichocephalic than in Central and Eastern regions which fall in mesocephaly class. The value is low in the scheduled tribe ethnic groups as compared to others in all zones and India except in North India, where it is higher than other groups. No correlation is observed with climatic factors and altitude with different ethnic groups. Among the different occupational groups value of cephalic index is low among priesthood and warfare groups as compared to artisans and menial workers groups but the differences are not significant. Among the language families, high value of the index is observed among speakers of Tibeto-Chinese languages who are having Mongoloid affinities whereas low value is recorded in Central and North Dravidian (Dravidian family) and Munda group (Austro-Asiatic family) speakers who are mostly tribals.

2.2. Nasal Index

The nose, like the dentition, is more directly regulated by the forces of selection than is true for other aspects of cranial morphology (Brace 1964; Brace and Hunt 1990; Brace et al. 1991).

Global variation in the human nasal index is widely viewed as the result of an adaptive response to climatic variability. Thomson and Buxton (1923) first, and many later investigators (Davies 1932; Weiner 1954; Crognier 1981; Beals

et al. 1984) found that nasal indices are positively associated with both temperature and humidity. Nasal indices are also related to latitude, which underlies variation in temperature and humidity (Newman 1953; Kelso 1970). In addition, nasal height and breadth, the components of the index, have been studied with respect to climate (Wolpoff 1968; Hiernaux and Froment 1976; Corgnier 1981). Nasal height is more strongly associated with temperature, covarying negatively, whereas nasal breadth is most strongly associated with humidity, covarying positively. However, the nasal index shows a higher correlation with climatic gradients than do either nasal height or nasal breadth alone. These patterns have been explained as evolutionary adaptations optimizing respiratory heat and moisture exchange in nasal mucosa, primarily to prevent ciliary and lung alveoli damage, as well as to moderate body water loss and maintain thermal equilibrium (*e.g.*, Thomson and Buxton 1923; Davies 1932; Weiner 1954).

Despite the associations described above, the nasal index is not universally accepted as a reflection of climatic adaptation (Hoyme 1965; St. Hoyme and Iscan 1989).

The mean value of nasal index is 75.53 among Indian populations (varies from 51.40 - hyperleptorhinae in Jain Pancham of Maharashtra to 96.75 - chamaerhinae in Garo or Mande of Assam).

The value is high (79.40 - mesorhinae) among scheduled tribes as compared to other ethnic groups (scheduled caste - 75.61; caste - 74.78 and community - 71.57). The value of index is low in Himalayan mountain complex region (70.68) followed by Indus-Ganga-Brahmaputra plains (72.13) and then gradually increases towards peninsular plateau (78.01) and islands region (79.98). From the climatic regions, the value is lowest from polar region (66.38) and starts increasing towards cold humid winters with short summers (71.31) and monsoon type with dry winters (72.11) regions and reach quite high values in tropical savannah type (78.05) to maximum in monsoon type with short dry season

Table 3: Nasal Index

Particulars	Subjects studied	No. of studies	Mean	Min	Max
1. NATURAL REGION					
Himalayan Mountain Complex	8408	129	70.68	60.83	92.86
Indus-Ganga-Brahmaputra Plains	22844	249	72.13	50.90	96.75
Peninsular Plateau	46480	563	78.01	51.40	95.90
Islands	751	13	79.98	69.58	93.01
2. CLIMATIC REGION					
Monsoon Type with Short Dry Season	4179	68	80.62	65.70	93.01
Monsoon Type with Dry Season	2827	48	76.75	67.92	95.10
Tropical Savannah Type	34389	354	78.05	51.40	95.90
Semi Arid Steppe Type	4768	92	76.10	64.52	92.90
Hot Desert Type	-	0	-	-	-
Monsoon Type with Dry Winters	31449	375	72.11	59.00	96.75
Cold Humid Winters with Short Summers	338	5	71.31	70.34	72.90
Polar Type	533	12	66.38	62.18	68.46
3. POLITICAL DIVISION OF INDIA					
I. NORTH INDIA					
A. Western Himalaya (S. No. 1, 2)					
1. Jammu and Kashmir	926	15	64.74	61.75	75.54
2. Himachal Pradesh	2294	24	65.81	60.83	74.10
3. Punjab	1035	13	69.78	64.71	75.20
4. Chandigarh UT	-	0	-	-	-
5. Haryana	-	0	-	-	-
6. Delhi UT	740	10	67.96	65.92	71.17
B. Central Himalaya (S. No. 7, Eight Districts of Uttar Pradesh)					
7. Uttar Pradesh	12232	123	71.01	59.00	86.10
8. Rajasthan	1698	11	73.63	70.03	84.10
II. WEST INDIA					
9. Gujarat	4750	50	73.53	65.84	80.80
10. Maharashtra	6935	120	77.45	51.40	94.80
11. Goa, Daman and Diu UT	-	0	-	-	-
12. Dadra and Nagar Haveli UT	-	0	-	-	-
III. EAST INDIA					
C. Eastern Himalaya (S. No. 13 to 20 and Darjeeling District of West Bengal)					
13. Arunachal Pradesh	338	5	71.31	70.34	72.90
14. Assam	4847	66	76.05	65.69	96.75
15. Nagaland	1699	35	77.46	64.42	92.86
16. Manipur	280	4	75.93	70.84	80.95
17. Mizoram	40	1	69.98	69.98	69.98
18. Tripura	311	5	72.92	70.63	75.31
19. Meghalaya	887	12	79.16	73.58	87.63
20. Sikkim	797	16	70.13	65.15	78.20
21. West Bengal	5378	55	72.18	65.40	94.70
Particulars					
22. Bihar					
23. Orissa					
IV. CENTRAL INDIA					
24. Madhya Pradesh					
V. SOUTH INDIA					
25. Karnataka					
26. Andhra Pradesh					
27. Tamil Nadu					
28. Kerala					
29. Pondicherry UT					
VI. ISLANDS					
30. Lakshadweep UT					
31. Andaman Islands and Nicobar Islands UT					
3A. ZONES OF INDIA					
I. North India					
II. West India					
III. East India					
IV. Central India					
V. South India					
VI. Islands					
INDIA (TOTAL)					
3B. REGIONS OF HIMALAYA					
A. Western Himalaya					
B. Central Himalaya					
C. Eastern Himalaya					
HIMALAYA (TOTAL)					
NON-HIMALAYAN REGIONS					
4. ETHNIC GROUP					
A. ZONES OF INDIA					
I. NORTH INDIA					
Caste					
Scheduled Caste					
Scheduled Tribe					
Community					
II. WEST INDIA					
Caste					
Scheduled Caste					
Scheduled Tribe					

Table 3: Contd....

Particulars	Subjects studied	No. of studies	Nasal Index		Particulars	Subjects studied	No. of studies	Nasal Index	
			Mean	Max				Mean	Max
Community	4359	56	74.39	51.40 87.56	Warfare	3114	26	69.01	62.11 82.48
III. EAST INDIA					Tade and Commerce	1937	26	73.16	63.00 79.60
Caste	14694	69	76.34	58.80 90.16	Agriculture	1846	33	72.74	64.00 83.70
Scheduled Caste	3684	40	74.44	66.39 88.70	Animal Husbandry	478	10	71.32	65.74 78.95
Scheduled Tribe	12551	172	79.95	64.42 96.75	Artisans	708	11	71.94	63.00 81.71
Community	2998	41	71.87	65.40 82.06	Mental Workers	5359	64	75.98	64.00 88.70
IV. CENTRAL INDIA					No Information	50001	714	76.08	51.40 96.75
Caste	104	2	79.26	72.92 85.14	6. LANGUAGE GROUP				
Scheduled Caste	-	0	-	-	I. AUSTRAL-ASIATIC FAMILY				
Scheduled Tribe	2350	32	82.30	75.90 92.20	Mon Khmer Group	1436	16	79.57	69.58 87.63
Community	50	1	77.91	77.91 77.91	Munda Group	5395	55	82.28	69.00 95.00
V. SOUTH INDIA					II. TIBETO-CHINESE FAMILY				
Caste	3199	75	76.03	65.70 85.50	(i) Siamese-Chinese Sub-Family				
Scheduled Caste	1304	30	78.45	69.70 82.31	Tai Group	85	1	77.26	77.26 77.26
Scheduled Tribe	3183	62	81.34	64.52 95.10	(ii) Tibeto-Burman Sub-family				
Community	3005	51	76.33	62.50 92.62	Bhotia Group	1774	23	66.05	60.83 75.54
VI ISLANDS					Himalaya Group	1039	20	71.33	65.15 78.50
Caste	-	0	-	-	North East Frontier Group	1111	19	74.05	69.66 90.58
Scheduled Caste	-	0	-	-	Bodo Group	1982	24	77.64	68.54 96.75
Scheduled Tribe	751	13	79.98	69.58 93.01	Naga Group	1901	39	77.96	64.42 92.86
Community	-	0	-	-	Kachin Group	13	1	81.26	81.26 81.26
INDIA					Kuki Chin Group	388	8	75.69	69.98 82.67
Caste	28943	292	74.78	58.80 90.16	III. DRAVIDIAN FAMILY				
Scheduled Caste	8188	117	75.61	63.00 92.31	South Dravidian Group	10454	215	78.04	64.52 95.10
Scheduled Tribe	23332	332	79.49	62.18 96.75	Central Dravidian Group	2010	32	82.06	73.67 91.81
Community	18020	213	71.57	51.40 92.62	North Dravidian Group	3009	11	72.82	67.00 89.12
B. REGIONS OF HIMALAYA					IV. INDO-EUROPEAN FAMILY				
A. Western Himalaya					Dard Group	572	6	63.34	62.11 66.50
Caste	818	8	64.90	62.11 74.10	North Western Group	30	1	73.58	73.58 73.58
Scheduled Caste	70	1	65.39	65.39 65.39	Southern Group	7057	114	76.77	51.70 88.40
Scheduled Tribe	1059	11	66.48	62.18 75.54	Eastern Group	1892	34	81.30	70.46 93.60
Community	1273	19	65.09	60.83 69.58	Bihari	18294	99	75.29	58.80 88.70
B. Central Himalaya					Central Group	18438	210	72.13	59.00 94.80
Caste	60	2	69.17	68.54 70.06	Pahari Group	929	14	67.43	62.42 77.93
Scheduled Caste	213	2	79.98	75.98 83.00	Unspecified	256	2	77.05	69.10 79.50
Scheduled Tribe	536	6	72.09	65.78 79.50	Other Languages	418	10	79.95	- 93.01
Community	348	5	70.73	66.30 73.14	No Information	-	0	-	-
C. Eastern Himalaya					6A. LANGUAGE FAMILY				
Caste	243	5	69.08	65.69 70.17	I. Austro Asiatic Family	6831	71	81.72	69.00 95.90
Scheduled Caste	1221	14	73.34	66.39 83.70	II. Tibeto Chinese Family	8293	135	73.87	60.83 96.75
Scheduled Tribe	9810	108	76.69	64.42 96.75	III. Dravidian Family	15473	258	77.55	64.52 95.10
Community	1491	26	73.27	65.75 82.06	IV. Indo European Family	47468	480	74.23	51.40 94.80
5. TRADITIONAL OCCUPATION					Other Languages	418	10	79.95	69.10 93.01
Priesthood	14940	70	75.89	58.80 82.81	No Information	-	0	-	-

(80.62) region. The value of index increases as one moves from cold climate towards hot climate (Table 3), (Bhasin et al. 1994; Bhasin and Walter 2001).

The value of nasal index is low from the North India zone (70.13) as compared to other zones. The nasal index is leptorhinae among the populations from the states of Jammu and Kashmir (64.74), Himachal Pradesh (65.81) of Western Himalayan region, and also from Punjab and Delhi (69.78 and 67.96, respectively) as compared to the peoples from any other State or Union Territory of India, where the nasal index is mesorhinae type (Table 3, Fig. 4). From Central Himalayan region among the population groups with Mongoloid affinities the value of index is less than 70.0 (different groups of Bhotias, Jaunsari) as also among caste groups compared to scheduled castes in which the value is quite

high (79.28). From Rajasthan also the value of the index is high among scheduled tribes (different groups of Bhils) as compared to the caste groups.

From East India, from the states of Assam, Nagaland, Manipur, Meghalaya of Eastern Himalayan region the values of index are quite high (76.05, 77.46, 75.93, 79.16, respectively) among the population groups with Mongoloid affinities as compared to the population groups from Western and Central Himalayan regions and Sikkim state (70.13). This may be due to racial elements present among the peoples from Eastern Himalayan region—Australoid (Pre-Dravidian), Caucasoid (Dravidian, and/or Aryan) and Mongoloid in varying degrees. From Nepal, among Sherpas, Chhetris, Rais, Thakurs, Tamangs, Newars, Sunwars, Limbus the nasal index is leptorhinae as also observed among




Fig. 4. Frequencies of nasal index in different regions of India (after Bhasin and Walter 2001)

different populations from Sikkim among whom Nepalis are in majority.

From West India the value of the index is 75.86 (mesorhinae) and quite high value is found among the populations from Maharashtra (77.45) as compared to Gujarat (73.53). The value is highest among the scheduled tribes (80.04 - mesorhinae) followed by scheduled castes (78.86) as compared to other ethnic groups. From West India, however, the value of index below 70 (leptorhinae) is observed among a few castes (Brahmin, Rajput, Khatri, Jain) and Parsis - a migrant community from Southwest Asia.

Among the populations from Central India (Madhya Pradesh) the value of index is quite high (82.09) followed by the states of Orissa (81.53) and Bihar (78.46) from which the studies are on scheduled tribes. From these states, as well as from West Bengal, the values of index are quite high (above 85) among tribals (*e.g.* Santal, Munda, Oraon, Paharia, Gond, Muria) and low (less than 70) among caste groups (mostly Brahmin).

The mean value of nasal index is again high from the states of South India (77.99) and in this zone also, the value is highest among scheduled tribes (81.34) followed by scheduled castes (78.45), as also observed from West India zone. The low values (less than 70) are observed among a few populations—Curgs, Lambadi-nomads migrated from North India, Toda, Kaku and Mohammadans. Whereas quite high values (more than 85) have been reported among Kuruba, Yeruvas, Sholega, Siddi (a community with black ancestry), Kadar, Malasar, Paniyan, Malavetan, Ulladan, Muthuvan, Urali, Nayadi, Malser, Malapulayan, Malayarayan. From this zone, the studies available on different caste populations are falling in mesorhinae class.

From Islands, among Andamanese from North and South the nasal index is quite high (more than 85), whereas others are falling in mesorhinae class except Nicobarese Southern who are having leptorhinae nasal index.

In general, it has been observed that the value of nasal index is low among population groups from the Himalayan region, however, differences are observed among the peoples from western and central Himalayan regions who are having Caucasoid (Aryans) affinities and Caucasoid (Aryan) and Mongoloid admixture in varying degrees in which value of nasal index is low while that of Eastern Himalayan region who are having admixture of Australoid (Pre-Dravidian),

Caucasoid (Dravidian and/or Aryans) and Mongoloid racial elements in varying degrees have high value.

As one moves towards Indus-Ganga-Brahmaputra plains region, the value of nasal index increases a little whereas among castes and communities who are Caucasoid (Aryans) the value of index is low as compared to scheduled castes and scheduled tribes and they are having Caucasoid (Aryan, Dravidian) and Australoid (Pre-Dravidian) racial elements in varying degrees.

In the peninsular region, the value of the index increases sharply and it has been observed that in West India, Central India and state of Orissa, the mean values are quite high among scheduled tribes and scheduled castes who are having mostly Australoid and/or Caucasoid (Dravidian) admixture, albeit higher caste groups from this region are having leptorhinae type of nose. In South India, among Dravidians, the value of nasal index is quite high among all the ethnic groups with highest in scheduled tribes who are having Australoid (Pre-Dravidian) and/or Caucasoid (Dravidian) racial element (s).

The nasal index shows a correlation with climatic gradients, but as it appears from the above discussion, the differences observed are also due to different racial elements present among them and due to various ethnic groups from the same natural regions.

Further when nasal index correlations are calculated with various climatic factors and altitude by different ethnic groups significant differences are observed but correlations are not high (Table 6). Therefore it appears difficult to evaluate to what extent there is an association of the nasal index with climatic adaptation.

The value of nasal index among different occupational groups is low in warfare (69.01 - leptorhinae), animal husbandry (71.32) and artisans (71.94) followed by agriculture (72.74) and trade and commerce (73.16) group as compared to priesthood and menial workers among whom the values are quite high (75.89 and 75.98, respectively) (Table 3), (Bhasin et al. 1994; Bhasin and Walter 2001). Most of the studies reported on priesthood group are from West Bengal, West India and South India among whom the value of nasal index is observed high as compared to other groups, particularly warfare group. The social stratification is most clearly shown in the nasal index—with increasing social rank the nasal index become slower *i.e.* nose becomes narrower and

more pronounced (Risley 1915; v. Eickstedt 1952; Büchi 1968). However in present study it is not so with the highest rank *i.e.* priesthood.

The value of the index is high among the speakers of Mon Khmer group (79.57) of Austro Asiatic; Bodo group (77.84), Naga group (77.96), Kachin group (81.26), Kuki Chin group (75.69), Tai group (77.26) and North East Frontier group (74.05) of Tibeto-Chinese languages from Eastern Himalayan region who are having Mongoloid affinities as compared to Dard group (63.34), Pahari group (67.43) of Indo-European and Bhotia group (66.05), Himalayan group (71.33) of Tibeto-Chinese languages from Western and Central Himalayan regions and state of Sikkim. Among the speakers of Eastern group of Indo-European, Munda group of Austro-Asiatic and Central Dravidian group of Dravidian languages who belong mostly to tribals the values are quite high (81.30, 82.28 and 82.06). In general, the values of nasal index are low among the speakers of Tibeto-Chinese (73.87) and Indo-European (74.23) languages as compared to others (Table 3) (Bhasin et al. 1994; Bhasin and Walter 2001).

In India, the nasal index is observed of mesorhinae type (75.53) among various population groups of India. The mean value of the index is observed high in Central India and Islands from where mostly scheduled tribe groups are reported, followed by South, East, West and North India. The value of nasal index is high in Eastern region (mesorhinae) as compared to Western and Central regions of Himalaya. In general, the high value of nasal index is observed among scheduled tribe groups followed by scheduled caste as compared to caste and community from various zones and regions. The value is high in Austro-Asiatic followed by Dravidian language speakers as compared to Indo-European and Tibeto-Chinese language speakers. No correlations are observed with various climatic factors and altitude by different ethnic groups. Leptorhinae type of nose is observed mostly from North India and in few caste groups from West and East India, although a few cases have also been reported from South India among inbreeding groups (Muhammadans and Curg), small ethnic groups (Kapu and Toda) and a group which is originally not from South India (Lambadi) and chamaerhinae type of nose is reported generally among scheduled tribe and scheduled caste groups from West, East, South India and Islands zones.

2.3. Facial Index

The mean value of facial index is 86.34 (mesoprosopic) among Indian populations, which varies from 75.00 (hypereuryprosopic type among Naga Sundan of Nagaland) to 122.80 (hyperleptoprosopic in Bhil Khandesh of Maharashtra). Among all the ethnic groups the index falls in mesoprosopic class (84.0 - 87.9) with low value in scheduled caste (85.79) and high in community (86.58). The facial index is leptoprosopic among population groups from Himalayan mountain complex natural region (88.33) and as one moves towards south the value starts decreasing gradually as observed in Indus-Ganga-Brahmaputra plains (87.00 - mesoprosopic) and peninsular plateau (85.84 - mesoprosopic) whereas in Islands region, the value is quite low among Andamanese who are having Negrito affinities (83.48 - euryprosopic) and high among Nicobarese (86.10 - mesoprosopic) among whom Mongoloid element is present. From the climatic regions—polar type, monsoon type with short dry season and monsoon type with dry season, the facial index is above 88.0 (leptoprosopic) followed by monsoon type with dry winters (87.21) as compared to semi arid steppe type and tropical savannah type regions where the values are low (84.78 and 85.34, respectively) (Table 4), (Bhasin et al. 1994; Bhasin and Walter 2001).

The facial index is leptoprosopic among the populations of North India zone as compared to other zones from where it is falling in mesoprosopic class. The values are quite high among the populations from the states of Jammu and Kashmir (90.71 - leptoprosopic) and Himachal Pradesh (89.50 - leptoprosopic) of Western Himalayan region (89.92 - leptoprosopic) as compared to Central Himalayan region (87.71 - mesoprosopic). From East India in the Eastern Himalayan region the value of index is low (85.47 - mesoprosopic) as compared to Western and Central Himalayan regions. However, a few exceptions have been observed from Manipur, Mizoram and Tripura states where the value of index is above 88 (leptoprosopic) which may be due to smaller number of studies reported from these states. From West Bengal the value of index is high (88.43) as most of the studies are among the caste and community populations as compared to Bihar and Orissa states from where the tribals are studied in majority (86.74 and 84.88, respectively). Among the scheduled tribes from

Central India the value of index is high (87.31) as compared to that observed from East and West India tribals (85.29 and 86.69, respectively).

Among the scheduled tribes from the West and South India zones, the mean values of facial index are high (86.69 and 89.49, respectively) as compared to other ethnic groups from these zones. From West India, the value of index is low (84.52) as compared to other zones and the index is euryprosopic among caste populations in this zone. In different states of South India zones the mean value varies from 83.99 (Karnataka) to 90.01 (Kerala); the value of index is quite high among the tribals of Kerala (in most cases above 88) as compared from other states.

In general the value of index is high in Northern India and decreases towards peninsular region (Table 4, Fig. 5).

The facial index correlations with various climatic factors and altitude by different ethnic groups though showing significant differences are not high (Table 6).

The values of facial index among priesthood, agriculture, artisans and menial workers traditional occupational groups are almost similar (86.03, 86.21, 85.30 and 86.04, respectively) as compared to warfare, trade and commerce and animal husbandry groups which have high values (87.93, 87.51 and 86.76, respectively). However, the differences observed are not significant which indicates gene flow among these groups as also observed for other traits (Bhasin et al., 1994; Bhasin and Walter 2001).

Among the speakers of Dard group, Pahari group of Indo-European and Bhotia group and Himalayan group of Tibeto-Chinese languages,




Fig. 5. Frequencies of facial index in different regions of India (after Bhasin and Walter 2001)

Table 4: Facial Index

Particulars	Subjects studied	No. of studies	Facial Index		
			Mean	Min	Max
1. NATURAL REGION					
Himalayan Mountain Complex	4907	71	88.33	75.00	99.00
Indus-Ganga-Brahmaputra Plains	10526	106	87.00	79.65	93.90
Peninsular Plateau	23433	320	85.64	79.97	122.80
Islands	596	8	86.10	82.08	88.91
2. CLIMATIC REGION					
Monsoon Type with Short Dry Season	1455	20	88.48	82.08	110.99
Monsoon Type with Dry Season	1466	11	88.47	84.50	90.75
Tropical Savannah Type	17486	238	85.34	79.97	122.80
Semi Arid Steppe Type	2662	47	84.78	80.24	87.71
Hot Desert Type	-	0	-	-	-
Monsoon Type with Dry Winters	15552	173	87.21	75.00	99.00
Cold Humid Winters with Short Summers	338	5	87.13	84.35	90.10
Polar Type	503	11	90.03	88.25	92.00
3. POLITICAL DIVISION OF INDIA					
I. NORTH INDIA					
<i>A. Western Himalaya (S. No. 1, 2)</i>					
1. Jammu and Kashmir	895	14	90.71	88.25	99.00
2. Himachal Pradesh	1679	19	89.50	88.01	94.15
3. Punjab	199	4	90.69	87.30	93.90
4. Chandigragh UT	-	0	-	-	-
5. Haryana	-	0	-	-	-
6. Delhi UT	100	1	88.95	88.95	88.95
<i>B. Central Himalaya (S. No. 7, Eight Districts of Uttar Pradesh)</i>					
7. Uttar Pradesh	3186	39	87.27	80.80	92.83
8. Rajasthan	722	5	86.57	86.34	87.40
II. WEST INDIA					
9. Gujarat	3160	30	83.99	81.40	86.18
10. Maharashtra	5440	101	84.83	79.97	122.80
11. Goa, Daman and Diu UT	-	0	-	-	-
12. Dadra and Nagar Haveli UT	-	0	-	-	-
III. EAST INDIA					
<i>C. Eastern Himalaya (S. No. 13 to 20 and Darjeeling District of West Bengal)</i>					
13. Arunachal Pradesh	338	5	87.13	84.35	90.10
14. Assam	4104	43	85.29	79.65	91.23
15. Nagaland	704	14	83.89	75.00	97.37
16. Manipur	100	1	88.15	88.15	88.15
17. Mizoram	40	1	90.50	90.50	90.50
18. Tripura	312	5	88.73	87.32	90.72
19. Meghalaya	860	11	84.99	82.00	88.89
20. Sikkim	80	1	87.75	87.75	87.75
21. West Bengal	3653	30	88.43	82.98	91.68

Particulars	Subjects studied	No. of studies	Facial Index		
			Mean	Min	Max
22. Bihar	3369	33	86.74	81.59	92.10
23. Orissa	3081	39	84.88	80.59	95.30
IV. CENTRAL INDIA					
24. Madhya Pradesh	2189	29	87.45	80.45	90.58
V. SOUTH INDIA					
25. Karnataka	1840	39	83.99	80.24	87.60
26. Andhra Pradesh	490	8	84.39	82.76	87.30
27. Tamil Nadu	158	4	87.63	83.50	89.94
28. Kerala	901	13	90.01	82.10	110.99
29. Pondicherry UT	1266	8	88.74	86.99	90.75
VI. ISLANDS					
30. Lakshadweep UT	-	0	-	-	-
31. Andman Islands and Nicobar Islands UT	22	2	83.48	83.38	83.54
	574	6	86.20	82.08	88.91
3A. ZONES OF INDIA					
I. North India	6781	82	88.33	80.80	99.00
II. West India	8600	131	84.52	79.97	122.80
III. East India	16641	183	86.27	75.00	97.37
IV. Central India	2189	29	87.45	80.45	90.58
V. South India	4655	72	86.61	80.24	110.99
VI. Islands	596	8	86.10	82.08	88.91
INDIA (TOTAL)	39462	505	86.34	75.00	122.80
3B. REGIONS OF HIMALAYA					
A. Western Himalaya	2574	33	89.92	88.01	99.00
B. Central Himalaya	659	10	87.71	86.26	89.15
C. Eastern Himalaya	6638	82	85.47	75.00	97.37
HIMALAYA (TOTAL)	9871	125	86.78	75.00	99.00
NON-HIMALAYAN REGIONS					
	29591	380	86.20	79.97	122.80
4. ETHNIC GROUP					
A. ZONES OF INDIA					
I. NORTH INDIA					
Caste	2781	28	87.29	80.80	90.63
Scheduled Caste	207	4	88.53	87.86	92.83
Scheduled Tribe	1344	18	88.53	86.26	92.29
Community	2449	32	89.37	86.32	99.00
II. WEST INDIA					
Caste	2664	53	83.66	79.97	92.41
Scheduled Caste	462	8	81.74	80.71	84.29
Scheduled Tribe	1436	20	86.69	80.24	122.80

Table 4: Contd.....

Particulars	Subjects studied	No. of studies	Facial Index		Particulars	Subjects studied	No. of studies	Facial Index	
			Mean	Max				Mean	Max
Community	4038	50	84.63	80.18 120.60	Warfare	2030	19	87.93	83.02 91.54
III. EAST INDIA					Trade and Commerce	709	11	87.51	81.37 90.56
Caste	3504	41	87.95	82.41 91.68	Agriculture	781	13	86.21	82.40 88.95
Scheduled Caste	2679	24	86.80	80.75 91.41	Animal Husbandry	193	4	86.76	82.73 93.90
Scheduled Tribe	8835	101	85.29	75.00 97.37	Artisans	385	4	85.30	82.52 87.84
Community	1623	17	87.05	83.95 91.43	Menial Workers	2899	28	86.04	80.71 91.41
IV. CENTRAL INDIA					No Information	28982	375	86.29	75.00 122.80
Caste	104	2	88.98	88.30 89.61	6. LANGUAGE GROUP				
Scheduled Caste	-	0	-	-	I. AUSTRALO-ASIATIC FAMILY				
Scheduled Tribe	2035	26	87.31	80.45 90.58	Mon Khmer Group	1354	15	85.33	82.00 88.91
Community	50	1	89.83	89.83 89.83	Munda Group	4056	35	85.79	80.59 95.30
V. SOUTH INDIA					II. TIBETO-CHINESE FAMILY				
Caste	1509	30	86.17	80.24 90.75	(i) Siamese-Chinese Sub-Family				
Scheduled Caste	543	10	83.21	80.38 87.66	Tai Group	85	1	85.81	85.81 85.81
Scheduled Tribe	854	13	89.49	83.50 110.99	(ii) Tibeto-Burman Sub-Family				
Community	1749	19	86.65	80.65 89.74	Bhotia Group	1508	18	88.69	86.26 92.00
VI. ISLANDS					Himalayan Group	216	4	89.49	87.75 92.29
Caste	-	0	-	-	North East Frontier Group	964	12	86.97	83.95 90.25
Scheduled Caste	-	0	-	-	Bodo Group	1788	18	85.45	80.27 91.15
Scheduled Tribe	596	8	86.10	82.08 88.91	Naga Group	872	16	83.09	75.00 97.37
Community	-	0	-	-	Kachin Group	-	0	-	-
INDIA					Kuki Chin Group	181	4	88.56	83.54 90.50
Caste	10562	154	86.45	79.97 92.41	III. DRAVIDIAN FAMILY				
Scheduled Caste	3891	46	85.79	80.38 92.83	South Dravidian Group	4510	71	86.60	80.24 110.99
Scheduled Tribe	15100	186	86.26	75.00 122.80	Central Dravidian Group	1871	29	86.08	80.45 90.58
Community	9909	119	86.58	80.18 120.60	North Dravidian Group	752	7	87.37	81.87 89.45
B. REGIONS OF HIMALAYA					IV. INDO-EUROPEAN FAMILY				
A. Western Himalaya					Dard Group	502	5	90.29	89.76 91.88
Caste	575	5	90.12	89.75 90.60	North Western Group	30	1	81.40	81.40 81.40
Scheduled Caste	-	0	-	-	Southern Group	5110	91	84.23	79.97 120.60
Scheduled Tribe	726	9	89.13	88.01 92.00	Eastern Group	1149	19	85.21	80.75 89.61
Community	1273	19	90.29	88.25 99.00	Bihari	6170	63	87.55	83.90 91.68
B. Central Himalaya					Central Group	7561	84	86.63	80.80 122.80
Caste	60	2	88.88	88.50 89.15	Pahari Group	561	8	90.38	89.50 94.15
Scheduled Caste	-	0	-	-	Unspecified	-	0	-	-
Scheduled Tribe	280	4	87.08	86.26 87.75	Other Languages	222	4	85.41	83.38 86.37
Community	319	4	88.05	87.25 88.75	No Information	-	0	-	-
C. Eastern Himalaya					6A. LANGUAGE FAMILY				
Caste	101	2	87.54	84.06 91.23	I. Austro Asiatic Family	5410	50	85.68	80.59 95.30
Scheduled Caste	1098	11	85.30	83.90 87.72	II. Tibeto Chinese Family	5614	73	86.47	75.00 97.37
Scheduled Tribe	4552	59	85.32	75.00 97.37	III. Dravidian Family	7133	107	86.54	80.24 110.99
Community	887	10	86.18	83.95 90.25	IV. Indo European Family	21083	271	86.42	79.97 122.80
5. TRADITIONAL OCCUPATION					Other Languages	222	4	85.41	83.38 86.37
Priesthood	3483	51	86.03	80.80 91.68	No Information	-	0	-	-

the value of facial index is above 88 (leptoprosopic) who are inhabiting Western, Central Himalayas as also among Lepcha, Bhutias of Sikkim state, whereas from Eastern Himalayan region the value of index varies between 84.0 and 87.9 (mesoprosopic) as observed among the speakers of Mon Khmer group (85.33) of Austro-Asiatic and North East Frontier group (86.97), Bodo group (85.45) and Naga group (83.09) of Tibeto-Chinese languages except Kuki Chin group speakers (88.56). In the Southern groups of Indo-European language speakers the value of index is low as observed from the West as compared to the Eastern group (85.21) and Bihari group from Eastern India (87.55). Among the speakers of Munda group the value of index is either low (85.79) as compared to North Dravidian group (87.37) or similar to Central Dravidian group, who are mostly tribals. However, in general, the differences among the speakers of different languages are not significant (Bhasin et al. 1994; Bhasin and Walter, 2001).

The mesoprosopic facial index is observed among various population groups of India and from all the zones of India except North India (Leptoprosopic). In the Himalayan region mesoprosopic type is predominant except in Western Himalayan region where leptoprosopic type is found. The value of index is high in the Himalayan region and it gradually starts declining towards South. Leptoprosopic facial index is observed among caste groups from West Bengal and Tamil Nadu, Kerala and Pondicherry of South India. Euryprosopic type of facial index is observed among caste groups of Gujarat and scheduled caste groups of West and South India zones. However, generally a high frequency is observed among Community or Caste > Caste or Community > Scheduled Tribe > Scheduled Caste except from West and South India where high value is observed among scheduled tribes. No correlations are observed with various climatic factors and altitude by different ethnic groups. The differences observed in the various language families for facial index are not significant (Dravidian > Tibeto-Chinese > Indo-European > Austro-Asiatic).

Considering India as a whole, the biogenetical traits distribution is seen to follow a pattern that might have been expected from the complex structure of its people. Indian populations are living in different natural and climatic regions. The composition of Indian populations is

multiethnic consisting of elements of autochthonous (Pre-Dravidian/Australoids), Caucasoids (Dravidians) who are inhabiting South of India and to some extent Central India, Caucasoids (mostly Aryans) in the North and Northwest India, Mongoloids in the Himalayan region, mostly in North East and East India and Negrito/Negroids in the Andaman Islands. Except for the Caucasoids, the other three groups mainly involve tribal populations and are numerically relatively smaller. The Caucasoids, Mongoloids and autochthonous elements have interbred in varying degrees to give mixtures which are difficult to unravel. The peoples are categorized into caste (on the basis of their traditional occupation, who are considered to have come in successive waves of immigration of known and sometimes unknown antiquity) and tribal (most of them are considered to be original inhabitants of India). The existence of caste and tribal systems made the Indian population genetically isolated. Indians speak different languages of various families—Austro-Asiatic, Tibeto-Chinese, Dravidian and Indo-European, which are represented within specific boundaries. In addition other factors that have influenced the distribution of biological traits in India are genetic drift, mutation and selection.

It is rather difficult to evaluate the effect of environmental factors on the somatometric traits as observed from the foregoing text. Generally in the Himalayan region, the populations are of lower medium stature, head and nose indices are mesocephalic and mesorhinae with leptoprosopic facial index, but the differences are observed between the peoples from Western and Central Himalayas and that of Eastern Himalayas, which are due to genetic composition *i.e.* different racial elements present among these populations (Table 5). Again from the Indus-Ganga-Brahmaputra plains region the differences are well marked between the population groups from the states of Punjab, Haryana, Uttar Pradesh (North India) and that of Bihar and West Bengal (East India) which are due to different racial elements present in these two regions. Further in latter two states, along with Orissa and Madhya Pradesh (peninsular region), the scheduled tribes are in majority and having Australoid (Pre-Dravidian) racial element show significant differences with the other ethnic groups who are predominantly Caucasoid (Aryan). Towards West of India also scheduled tribes who speak Indo-European languages show significant

Table 5: Classification of stature and indices

<i>Particulars</i>	<i>Stature</i>	<i>Cephalic Index</i>	<i>Nasal Index</i>	<i>Facial Index</i>
INDIA	L. Med. (163.06)	Meso (76.06)	Meso(78.53)	Meso (88.34)
<i>Natural Region</i>				
Himalayan	L. Med. (162.13)	Meso (76.38)	Meso (70.68)	Lepto (88.33)
Mountain Complex				
Indus-Ganga-	Med. (164.15)	Dolicho (75.87)	Meso (72.13)	Meso (87.00)
Brahmaputra Plains				
Peninsular Plateau	L. Med. (162.77)	Meso (76.07)	Meso (78.01)	Meso (85.64)
Islands	Short (155.50)	Meso (78.64)	Meso (79.98)	Meso (86.10)
<i>Zones of India</i>				
North India	Med. (164.70)	Dolicho (73.65)	Meso (70.13)	Lepto (88.33)
West India	L. Med. (163.09)	Meso (78.24)	Meso (75.86)	Meso (84.52)
East India	L. Med. (163.05)	Meso (76.82)	Meso (77.06)	Meso (86.27)
Central India	L. Med. (160.32)	Dolicho (74.96)	Meso (82.09)	Meso (87.45)
South India	L. Med. (161.23)	Meso (76.07)	Meso (77.99)	Meso (86.61)
Islands	Short (155.50)	Meso (78.64)	Meso (79.98)	Meso (86.60)
<i>Regions of Himalaya</i>				
Western Himalaya	L. Med. (163.38)	Dolicho (74.00)	Lepto (65.51)	Meso (89.92)
Central Himalaya	L. Med. (163.13)	Meso (76.71)	Meso (72.85)	Meso (87.71)
Eastern Himalaya	L. Med. (161.20)	Meso (78.09)	Meso (75.56)	Meso (85.47)
<i>Ethnic Groups</i>				
Caste	Med. (165.25)	Meso (76.20)	Meso (74.78)	Meso (86.45)
Scheduled Caste	L. Med. (162.11)	Dolicho (75.96)	Meso (75.61)	Meso (85.79)
Scheduled Tribe	Short (157.75)	Dolicho (75.81)	Meso (74.49)	Meso (86.26)
Community	Med. (164.18)	Meso (76.17)	Meso (71.57)	Meso (86.58)
<i>Language Family</i>				
Austro-Asiatic	Short (158.89)	Dolicho (75.63)	Meso (81.72)	Meso (85.68)
Family				
Tibeto-Chinese	L. Med. (161.37)	Meso (77.67)	Meso (73.87)	Meso (86.47)
Family				
Dravidian Family	L. Med. (161.34)	Dolicho (75.52)	Meso (77.55)	Meso (86.54)
Indo-European	Med. (164.52)	Dolicho (75.99)	Meso (74.23)	Meso (86.42)
Family				

L. = Lower, Med. = Medium

Table 6: Correlations with climatic factors and ethnic groups

<i>Ethnic groups</i>	<i>Somatometry</i>			
	<i>Stature</i>	<i>Cephalic Index</i>	<i>Nasa Index</i>	<i>Facial Index</i>
Mean Annual Temperature				
Caste	0.025	0.098	0.126 ¹	0.027
Scheduled Caste	-0.110	0.081	0.033	-0.007
Scheduled Tribe	-0.151 ²	-0.209 ³	0.267 ³	-0.107 ³
Community	0.002	0.084	0.205 ²	-0.470 ³
Total	0.068 ¹	0.027	0.082 ¹	-0.179 ³
Mean Annual Rainfall				
Caste	0.023	-0.141 ¹	0.011	0.314 ³
Scheduled Caste	-0.249 ²	0.247 ²	-0.031	0.400 ²
Scheduled Tribe	-0.272 ³	0.317 ³	-0.157 ²	0.052 ¹
Community	-0.187 ²	0.078	0.016	0.189 ¹
Total	-0.331 ³	0.055	0.126 ³	0.176 ³
Mean Annual Humidity				
Caste	-0.052	-0.028 ³	0.088	0.190 ¹
Scheduled Caste	-0.253 ²	0.321 ³	-0.025	0.215
Scheduled Tribe	-0.308 ³	0.252 ³	-0.170 ²	-0.040 ²
Community	-0.206 ²	0.156 ¹	0.129	0.261 ²
Total	-0.296 ³	0.125 ³	0.093 ²	0.103 ¹
Mean Altitude				
Caste	-0.089	0.065 ²	-0.010	-0.169 ¹
Scheduled Caste	0.177 ¹	-0.236 ²	0.174	-0.418 ²
Scheduled Tribe	0.201 ³	0.078	-0.173 ²	0.045 ¹
Community	-0.068	-0.086	-0.021	0.192 ¹
Total	-0.047	-0.022	0.015	-0.004

1. Significant at P < 0.05 2. Significant at P < 0.01 3. Significant at P < 0.001

differences with other ethnic groups. As one moves towards South India, there are a number of scheduled tribes which are numerically smaller as compared to the tribes from the adjoining states of West India, Madhya Pradesh and Orissa. Though they speak Dravidian languages, they are showing significant differences from other ethnic groups of South India who are Caucasoids (Dravidian). It appears that they are basically Australoid (Pre-Dravidian) having some admixture with Caucasoid (Dravidians) and adopted their language. From Islands, Andaman is inhabited by Andamanese and Onges who are considered to be Negritos and Nicobarese and Shompens, who are having Mongoloid affinities. The differences which are observed among the various populations of India are due to different racial elements present among them and the division of the peoples into caste and tribal systems *i.e.* different ethnic groups. The differences are well marked between tribals and other populations whereas among caste, scheduled caste and community the differences are invariably in relation to their proximate category *e.g.* between castes and scheduled castes between communities and castes etc. This may be explained as caste system is based on traditional occupational groups and it has been observed that there is gene flow among them as the differences observed between various occupational groups are not high, whereas communities are those which are not identifiable on the basis of caste system or are not denoted as tribes and which are much more heterogeneous than either a caste group or a tribe. Further the differences observed between scheduled tribes and other ethnic groups can also be explained by the fact that they are the oldest inhabitants in their ecological niches as compared to other ethnic groups who have migrated to this area only a few thousand years back. The effect of different climatic factors (temperature, rainfall, humidity) and altitude on stature and different indices in relation to ethnic

groups shows significant differences but correlations observed are not high.

The differences observed among Indian populations are mainly due to the presence of various racial elements and prevailing of ethnic groups among them. Differences due to climatic factors are observed but no association is established between biological traits and climatic factors since the values of correlations observed are not high.

3. GENETIC DISTANCES

For anthropometric measurement (stature) and indices the number of investigations is maximum from East India (about 45 per cent for Eastern Himalayan region) followed by North and South India.

The distribution of the available studies shows that about 30 per cent of the total reports are on scheduled tribes followed by castes, communities and scheduled castes. Again in all the zones of India, in general, more studies are available among scheduled tribes (around 35 per cent), followed by castes (about 26 per cent), communities and scheduled castes (about 13 per cent) except from North India where the distribution is high in castes, communities, scheduled castes followed by scheduled tribes. Further about 50 per cent of the caste groups studied from India belong to North India zone and scheduled tribes to East India zone.

In the language families—the maximum studies are from Indo-European family followed by Dravidian, Tibeto-Chinese and Austro-Asiatic families.

The dendrograms generated on the basis of anthropometric for various Indian populations subdivided by zones (North, West, East, Central and South India) are represented in figure 6. The trend observed in the dendrogram is that West,




Fig. 6. Dendrograms of anthropometric traits according to different zones of India (after Bhasin and Walter 2001)

East, South and Central India together form one major cluster and separating out from them is North India. It is noted that from North India, maximum studies are available among castes and communities and it has been reported by Balakrishnan (1978) that considerable homogeneity has been demonstrated among populations of North-West India and it has been further pointed out by Balakrishnan (1981) that homogeneity is increased in certain regions by excluding the tribal populations.

Figure 7 shows dendrograms for various Indian populations based on social ranking. It is observed that caste group is invariably clustered with the scheduled caste and the community, while the scheduled tribe is distinct from all. An explanation of this phenomenon can be found in the fact that the scheduled castes may have in them a substantial contribution of gene flow from the higher castes in past generations (Majumdar, 1965). Since the communities are grouped on the basis of religion, occupation etc. they represent a fairly heterogeneous group. The distribution of major ethnic components may be present in the proportions similar to those observed in caste groups as the groups forming the communities in one way or the other are the offshoot of the major Indian ethnic stock (mainly the groups based on *Varna* system of Manu) at different periods of time in Indian history. The scheduled tribes, who represent mainly the Australoid (Pre-Dravidian), or Mongoloid strain, understandably are distinct from all the other groups subdivided by social ranking. Similar findings have also been reported by many other investigators (for example Balakrishnan 1978; Bhasin et al. 1986; Char et al. 1989; Chakraborty et al. 1986; Danker-Hopfe et al. 1988; Mukherjee et al. 1979, 1989; Roychoudhury 1984; Saha et al. 1976; Walter et al. 1993).

The dendrogram for different Indian populations subdivided by language families are presented in figure 8. As observed, from the dendrograms, the speakers of Tibeto-Chinese linguistic group show small differences with the Indo-European linguistic group and they are closer to Austro-Asiatic and Dravidian linguistic groups. From this dendrogram it is reasonable to believe that the current population of Indian Region is the product of differential admixture of Austro-Asiatic with Dravidian, Dravidian with Indo-Aryan and Tibeto-Chinese with Indo-European and also with Dravidian and Austro-Asiatic.

Clustering of the population groups of India

subdivided by geographical proximity and social ranking on the basis of anthropometric distances is clearly seen in the dendrogram presented in figure 9.

It has been observed that except for North India, where all the four ethnic groups fall nearer to each other in social rank— caste, community or/and scheduled caste and scheduled tribe while for the rest of the zones it is generally observed that caste, scheduled caste and community groups are closer to each other, whereas scheduled tribe groups are nearer to one another. The ethnic groups of West India are closer to South India as observed before and scheduled tribes of East India and South India. Thus, overall the various ethnic groups of North India are showing less difference, different ethnic groups of West India and South India are falling nearer to each other and scheduled tribes of East India and South India are showing small distances and, in general, scheduled tribes are falling apart from the other ethnic groups.

In the dendrogram generated on the basis of anthropometric traits (Fig. 10) for Indian populations classified by language families and social-ranking it has been observed that Tibeto-Chinese join the main cluster which consists of Indo-European and Dravidian speaking people and Austro-Asiatic (Scheduled Tribe) make a cluster with Dravidian (Scheduled Tribe) and Indo-European (Scheduled Tribe). This is understandable, since as reported earlier the Tibeto-Chinese speakers are dominating Eastern Himalayan region in Eastern India and this zone is the meeting ground of three ethnic groups: people with Mongoloid elements from the Northeastern region, non-tribals of Caucasoid origin from Northwestern region and tribals from the Southern and Southwestern regions. Whereas Austro-Asiatic (Scheduled Tribe) make a cluster with Dravidian (Scheduled Tribe) and Indo-European (Scheduled Tribe) and these are showing small differences with the Tibeto-Chinese (Scheduled Tribe) speakers. It has already been established that Austro-Asiatic speaking tribals are predominating in Bihar, Orissa, as well as also in Central India, Assam and Nicobar Islands, and admixture has taken place among them and other linguistic groups—Dravidian, Indo-European and Tibeto-Chinese, as well as between these groups in varying degrees. The Indo-European and Dravidian speakers are showing small distances and it has been suggested that the Dravidians who entered




Fig. 7. Dendrograms of anthropometric traits according to different ethnic groups of India (after Bhasin and Walter 2001)




Fig. 8. Dendrograms of anthropometric traits according to different language families of India (after Bhasin and Walter 2001)




Fig. 9. Dendrogram of anthropometric traits according to different zones and ethnic groups of India (after Bhasin and Walter 2001)




Fig. 10. Dendrogram of anthropometric traits according to different language families and ethnic groups of India (after Bhasin and Walter 2001)

Indian region early and moved southward with the arrival of Aryans have had admixture with them.

The above discussion was based on the dendrograms generated from different Indian populations subdivided by geographical regions, social status and linguistic division that have been considered separately as well as in combination viz., geographical regions/zones with social ranks and social ranks with linguistic groups. From the dendrograms derived from various geographical regions/zones (North, West, East, Central and South India) it has been observed that there is a general tendency of correlation between morphological distance with geographic distance. The dendrograms based on social ranking and morphological traits show that the scheduled tribes are distant from all the other groups and caste group is invariably clustered with the scheduled caste and the community, which shows that scheduled caste population may have in them a substantial trace of gene flow from upper castes or/and from communities, who are grouped on the basis of religion, occupation, region, etc. Considering now India as a whole, the biological

trait distribution is seen to follow a simpler pattern than that might have been expected from the complex history of its people. The scheduled tribe groups stand out as a uniform and unique population, showing less similarity with the neighbouring peoples and certainly much less when compared to the caste groups of India. From the dendrograms based on different linguistic groups, it has been observed that Austro-Asiatic, Dravidian and Indo-European linguistic categories form one cluster and Tibeto-Chinese category stand out separately. It may be concluded that the current population of India is the product of differential admixture between three distinct population entities viz., Austro-Asiatic (Australoid or Pre-Dravidian), Dravidian and Indo-European (Caucasoid) linguistic groups, whereas the Tibeto-Chinese linguistic group along the northeastern and northern peripheries of the subcontinent is the product of admixture between Mongoloids and Australoid and Caucasoid populations in varying degrees.

The migration of early *Homo sapiens sapiens* from Africa to Asia and Pacific might be through Indian subcontinent. Modern humans are believ-

ed to have arisen in Africa and subsequently migrated to the rest of the world beginning around 100,000 years ago. India may be in the path of this and subsequent migration events out of Africa and thus these migrants were the ancestors of the original inhabitants of Indian subcontinent (Mourant 1983; Cavalli-Sforza et al. 1994).

Descendants of Paleolithic expansion into India believed to have contributed substantially to tribal populations, which constitute 8.1 percent of the total population of India (Census of India, 1991).

Later migration events spread Austriacs – Austro-Asiatic (may be either from Indo-China and South China or a very off-shoot of the Mediterranean people who came to India from the West even before the Dravidian), Dravidian, Tibeto-Chinese (the area of the speeches stretches from Baltistan in the West to the north eastern frontiers of the country and further reaching up to southern most portions of Assam) followed by Indo-European speakers in to the Indian subcontinent. The Indo-European speakers introduced the caste system, a social hierarchy that strictly governs marriage and mating practices. The later migrants i.e., Indo-European speakers of North India had less influence on Dravidian speakers of South India, therefore both the groups are genetically and linguistically distinct from each other (Bhasin et al., 1994; Bhasin and Walter 2001).

The genetic diversity of caste and tribal population groups is well established (Bhasin et al. 1994, Bhasin and Walter 2001; Cordaux et al. 2004).

NOTES

1. Three more states have come into existence recently. Now Bihar has been divided into Bihar and *Jharkhand*; from Madhya Pradesh have resulted the states of Madhya Pradesh and *Chattisgarh*; the earlier state of Uttar Pradesh is now divided into Uttar Pradesh and *Uttaranchal*. The states in italics are newer ones. India at present comprises 28 states.
2. The frequency distributions for the State of Goa and Union Territory of Daman and Diu have been listed under Goa, Daman and Diu.

REFERENCES

- Balakrishnan V 1978. A preliminary study of genetic distances among some populations of the Indian sub-continent. *J Hum Evol*, 7: 67-75.

- Balakrishnan V 1981. Genetic distance studies in India. *Acta Anthropogenet*, 5: 57-65.
- Basu A, Majumdar PP, Ghosh AK and Biswas SK 1980. Human biological variations in Asia with special reference to India. pp. 197-224. In: J. Hiernaux (Ed.). *Human Biological Diversity*. Paris: Masson.
- Beals KL, Smith CL, Dodd SM 1984. Brain size, cranial morphology, climate and time machines. *Curr Anthropol*, 25: 301-330.
- Bhasin MK 1988. *Biology of the Peoples of Indian Region (Bangladesh, Bhutan, India, Maldives, Nepal, Pakistan, Sri Lanka). A Classified and Comprehensive Bibliography*. Delhi: Kamla-Raj Enterprises.
- Bhasin MK 2006a. Genetics of Castes and Tribes of India: Indian Population Milieu. *Int J Hum Genet*, 6(3): 233-274.
- Bhasin MK 2006b. Genetics of Castes and Tribes of India: Glucose-6-Phosphate Dehydrogenase Deficiency and Abnormal Haemoglobins (HBS and HBE). *Int J Hum Genet*, 6(1): 49-72.
- Bhasin MK 2006c. Genetics of Castes and Tribes of India: A Review of Population Differences in Red and Green Colour Vision Deficiency in India. *Int J Hum Genet*, 6(1): 81-88.
- Bhasin MK 2006d. Genetics of Castes and Tribes of India: Taste Sensitivity. *Int J Hum Genet*, 6(2): 145-151
- Bhasin MK, Walter H 2001. *Genetics of Castes and Tribes of India*. Delhi: Kamla-Raj Enterprises.
- Bhasin MK, Walter H 2002. Genetics of Castes and Tribes of India: Serum Protein Polymorphisms KM and GM Systems. *Int J Hum Genet*, 2(2): 119-138.
- Bhasin MK, Walter H, Danker-Hopfe H 1994. *People of India. An Investigation of Biological Variability in Ecological, Ethno-Economic and Linguistic Groups*. Delhi: Kamla-Raj Enterprises.
- Bhasin MK, Walter H and Danker-Hopfe H 1992. *The Distribution of Genetical, Morphological and Behavioural Traits among the People of Indian Region (Bangladesh, Bhutan, India, Maldives, Nepal, Pakistan, Sri Lanka)*. Kamla-Raj Enterprises, Delhi.
- Bhasin MK, Walter H, Chahal SMS, Bhardwaj V, Sudhakar K, Danker-Hopfe H, Dannewitz A, Singh IP, Bhasin V, Shil AP, Sharma MB, Wadhavan D 1986. Biology of the people of Sikkim, India: 1. Studies in the variability of genetic markers. *Z Morph Anthropol*, 77: 49-86.
- Brace CL 1964. A nonracial approach towards the understanding of human diversity. In: *The Concept of Race*. A. Montagu (Ed.). New York: The Free Press of Glencoe, pp. 103-152.
- Brace CL, Hunt KD 1990. A non-racial craniofacial perspective on human variation: A (australia) to Z (uni). *Am J Phys Anthropol*, 83: 341-460.
- Brace CL, Smith SL, Hunt KD 1991. What big teeth you had Grandma! Human tooth size, past and present. In: *Advances in Dental Anthropology*. M.A. Kelley and C.S. Larsen (Eds.). New York: Wiley Liss, pp. 33-57.
- Büchi, EC 1968. Rassengeschichte des indopakistanischen Subkontinents. In: K Saller (Hrg.), *Rassengeschichte der Menschheit*, 1. Lieferung. München-Wien: Oldenbourg, pp. 109-184.
- Cavalli-Sforza LL, Piazza A, Menozzi P 1994. *The History and Geography of Human Genes*. Princeton,

- New Jersey: Princeton University Press.
- Census of India 1991. Series-1, Paper 1 of 1991. *Provisional Population Totals*. New Delhi: Office of the Registrar General & Census Commissioner, India.
- Chakraborty R, Walter H, Mukherjee BN, Malhotra KC, Sauber P, Banerjee S, Roy M 1986. Gene differentiation among ten endogamous groups of West Bengal, India. *Am J Phys Anthropol*, **71**: 295-309.
- Char KSN, Lakshmi P, Gopalam KB, Sastry JG, Rao PR 1989. Genetic differentiation among some endogamous populations of Andhra Pradesh, India. *Am J Phys Anthropol*, **78**: 421-429.
- Cordaux R, Aunger G, Bentley G, Nazidze I, Sirajuddin SM, Stoneking M 2004. Independent origins of Indian caste and tribal paternal lineages. *Curr Biol*, **14**: 231-235.
- Corgnier F 1981. Climate and anthropometric variations in Europe and the Mediterranean area. *Ann Hum Biol*, **8**: 99-107.
- Danker-Hopfe H, Das BM, Walter H, Das PB, Das R 1988. Anthropological studies in Assam, India. 6. Differentiation processes among Assamese populations. *Anthrop Anz*, **46**: 159-184.
- Davies A 1932. A re-survey of the morphology of the nose in relation to climate. *J Roy Anthropol Inst*, **62**: 337-359.
- Devdas RP, Easwaran PP, Ponnammal, K 1977. Diet and nutrition in the first year of life. Part II. Cross-sectional and semi-longitudinal study of physical measurements. *Ind J Nutr Diet*, **14**: 361-365.
- Dutta PC 1978. *The Great Andamanese Past and Present*. Calcutta: Anthropological Survey of India.
- Easwaran PP, Saktivelmani A, Devdas RP 1974. Heights and weight of 7, 8 and 9 year old children in selected schools of Coimbatore city. *Ind J Nutr Dietet*, **11**: 63-71.
- Eickstedt E Frh v 1934. *Rassenkunde und Rassen-geschichte der Menschheit*. Stuttgart: Ferdinand Enke Verlag.
- Eickstedt E Frh v 1952. Rassentypen und Typendynamik von Asien. In: 1. F. Valjavec (Ed.). *Historia Mundi*. Volume. München, pp. 147-166.
- Ganguly P 1976. *Physical Anthropology of the Nicobarese*. Memoir No. 34, Calcutta: Anthropological Survey of India.
- Ganguly P 1979. Progressive decline in stature in India. A study of sixty population groups. In: WA Stini (Ed.). *Physiological and Morphological Adaptation and Evolution*. The Hague: Mutton, pp. 315-337.
- Gopalan C, Swaminathan MC, Krishnakumari VK Rao, HD, Vijayraghavan K 1973. Effect of calorie supplementation on growth of undernourished children. *Am J Clin Nutr*, **26**: 563-566.
- Guha BS 1935. Racial affinities of the people of India. *Census of India, 1931. Report, Vol. 1, Pt. III Ethnographical. Sec. A*. Government of India, Simla.
- Harrison GA, Kuchemann CF, Moore CMAS, Boyce, AJ, Baju T, Mourant AE, Godber MJ, Glasgow BG, Kopec AC, Tills D, Clegg, EJ 1969. The effects of altitudinal variation in Ethiopian populations. *Phil Trans R Soc*, **B 256**: 147-182.
- Hiernaux J, Froment, A 1976. The correlations between anthropological and climatic variables in sub-Saharan Africa. Revised estimates. *Hum Biol*, **48**: 757-767.
- Hornwell J 1923. The origins and ethnological significance of Indian boat designs. *Mem Asiat Soc Beng*, **7**: 139-256.
- Hoyme LE 1965. The nasal index and climate: A spurious case of natural selection in man. Abstract. *Am J Phys Anthropol*, **23**: 336-337.
- Kaplan BA 1954. Environment and Human Plasticity. *Am Anthropol*, **56**: 780-800.
- Karve I, Dandekar, VM 1951. *Anthropometric Measurements of Maharashtra*. Deccan College Monograph S. No. 8. Poona: Deccan College.
- Karve I, Malhotra, KC 1968. A biological comparison of eight endogamous groups of the same rank. *Curr Anthropol*, **9**: 109-124.
- Keith A 1936. Review of B.S. Guha's The Racial Affinities of the Peoples of India. *Man*, **36**: 28-30.
- Kelso AJ 1970. *Physical Anthropology: An Introduction*. Philadelphia: J.B. Lippincott Co.
- Knussmann R (Ed.) 1988. *Anthropologie, Handbuch der vergleichenden Biologie des Menschen Band I: Wesen und Methoden der Anthropologie 1. Teil: Wissenschaftstheorie, Geschichte, morphologische Methoden*. Stuttgart, New York: Gustav Fischer
- Lohman, TG, Roche, AF and Martorell, R 1988. *Anthropometric Standardization Reference Manual*. Illinois: Human Kinetics Books Champaign.
- Mahalanobis PC 1927. Analysis of race mixture in Bengal. *J. Asiat Soc Beng*, **23**: 301-333.
- Mahalanobis PC, Majumdar DN, Rao CR 1949. Anthropometric survey of the United Provinces, 1941. A statistical study. *Sankhya*, **9**: 89-324.
- Majumdar DN 1965. *Races and Cultures of India*. New Delhi: Asia Publishing House, 4th Edition.
- Majumdar DN and Sen AR 1949. Racial elements in Gujarat. *J. Gujarat Res. Soc.*, **11**: 118.
- Majumdar DN, Rao CR 1960. *Race Elements in Bengal. A Quantitative Study*. Calcutta: Asia Publishing House.
- Malhotra KC, Balakrishnan V, Karve I 1981. Anthropometric variation in Tamil Nadu. In: L.D. Sanghvi, Balakrishnan and I. Karve (Eds.). *Biology of the People Tamil Nadu*. Calcutta: Indian Society of Human Genetics and Indian Anthropological Society, pp. 50-74.
- Malhotra MS 1966a. People of India, including primitive tribes. A survey on physiological adaptation, physical fitness and nutrition. In: P.T. Baker and J.S. Weiner (Eds.). *The Biology of Human Adaptability*. New Delhi: Oxford, pp. 329-355.
- Martin R, Saller K 1957. *Lehrbuch der Anthropologie*. Volumes I and II. Gustav Fischer Verlag, Stuttgart.
- Mills CA 1937. Geographic and time variations in body growth and age at menarche. *Hum Biol*, **9**: 43-56.
- Mills CA 1942. Climatic effect on growth and development with particular reference to the effects of tropical residence. *Am Anthropol*, **44**: 1-14.
- Mourant AE 1983. *Blood Relations, Blood Groups and Anthropology*. Oxford: Oxford University Press.
- Mukherjee BN, Majumdar PP, Malhotra KC, Das SK, Kate SL, Chakraborty, R 1979. Genetic distance analysis among nine endogamous population groups of Maharashtra, India. *J Hum Evol*, **8**: 567-570.
- Mukherjee BN, Malhotra KC, Roy M, Banerjee S, Walter H, Chakraborty R 1989. Genetic heterogeneity and population structure in eastern India: Red cell enzyme variability in ten Assamese populations. *Z Morph Anthropol*, **77**: 287-296.

- Newman MT 1953. The application of ecological rules to the racial anthropology of the New World. *Am. Anthropol.*, **55**: 311-327.
- Olivier, G 1963. *Anthropological Repercussions of the Caste System*. Anthropology on the March, Social Sciences Association, Madras.
- Pachauri S, Marwah SM, Rao NSN 1971. A multifactorial approach to the study of the factors influencing birth weight in the urban community of New Delhi. *Ind J Med Res*, **59**: 1318-1341.
- Rakshit HK 1966. The Brahmans of India: An anthropometric study. *Man in India*, **46**: 33-58.
- Rao DH, Satyanarayana K 1974. Nutritional status of tribal pre-school children of Andhra Pradesh. *Ind J Nutr Diet*, **11**: 328.
- Rao DH, Satyanarayana K, Vijayaraghavan K, Sastry JG, Naidu AN, Swaminathan MC 1975. Evaluation of the special nutrition programme in the tribal areas of Andhra Pradesh. *Ind J Med Res*, **63**: 652-660.
- Risley H Sir 1969. *The People of India*. (1915). Delhi: Orient Book Reprint Corporation, Reprinted.
- Roberts DF 1953. Body weight, race and climate. *Am J Phys Anthropol*, **11**: 533-558.
- Roberts DF 1973. *Climate and Human Variability*. Addison Wesley Module in Anthropology, 34.
- Roychoudhury AK 1984. Genetic relationships of Indian populations. In: *Human Genetics and Adaptation*. Volume 1. KC Malhotra, A. Basu (Eds.). Calcutta: Indian Statistical Institute, pp. 146-174.
- St. Hoyme LE, Iscan MY 1989. Determination of sex and race: Accuracy and assumptions. In: MY Iscan, KAR Kennedy (Eds.). *Reconstruction of Life from the Skeleton*. New York: Alan R. Liss, pp. 59-93.
- Saha N, Kirk RL, Shanbhag S, Joshi SR, Bhatia HM 1976. Population genetic studies in Kerala and the Nilgiris (South-West India). *Hum. Hered*, **26**: 175-197.
- Sarkar SS, Deka U, Agarwal KK 1955. Brachycephaly in India. *Man in India*, **35**: 257.
- Shapiro HL 1939. *Migration and Environment*. New York: Oxford University Press.
- Singh IP, Bhasin MK 1989. *Anthropometry*. Kamla-Raj Enterprises, Delhi, first published in 1968.
- Singh IP, Bhasin MK 2004 *A Manual on Biological Anthropology*. Delhi: Kamla-Raj Enterprises.
- Spielman RS, Smouse PE 1976. Multivariate classification of human populations. 1. Allocation of Yanomama Indians to villages. *Am J Hum Genet*, **28**: 317-331.
- Takahashi E 1971. Geographic distribution of human stature and environmental factors- an ecologic study. *Anthrop Soc Nippon*, **79**: 259-286.
- Thomson A, Buxton LHD 1923. Man's nasal index in relation to certain climatic conditions. *Roy Anthropol Inst*, **53**: 92-122.
- Tobias, Ph V 1985.: The negative secular trend. *J Hum Evol*, **14**: 347-356.
- Vijayalakshmi R, Devaki, SR 1976. Nutritional status of expectant mothers belonging to two different income levels. *Ind J Nutr Dietet*, **13**: 63-66.
- Walter H 1982. Human physique and climate. *Anthrop Közl*, **26**: 99-111.
- Walter H, Bhasin MK, Danker-Hopfe H, Khanna A, Hahm U, Witt I 1993. Genetic studies on eleven populations of Jammu and Kashmir, India. *J Hum Ecol*, **4**: 1-14.
- Weiner JS 1954. Nose shape and climate. *Am J Phys Anthropol*, **12**: 1-4.
- Weiner JS, Lourie JA 1969. *Human Biology. A Guide to Field Methods*. Oxford and Edinburgh: Blackwell Scientific Publications.
- Weiner JS, Lourie JA 1981. *Practical Human Biology*. London: Academic Press.
- Wolanski N 1966. The secular trend; microevolution, physiological adaptation and migration, and their causative factors. Problems of World Nutrition. In: *Proceedings of the 7th International Congress of Nutrition, Hamburg*, pp. 96-108.
- Wolanski N 1967. Basic problems in physical development in man in relation to the evaluation of development of children and youth. *Curr Anthropol*, **8**: 35-60.
- Wolpoff MH 1968. Climatic influence on the nasal aperture. *Am J Phys Anthropol*, **29**: 405-423.
- Wolpoff MH 1980. *Palaeoanthropology*. New York: Knopf.