

Cytogenetic Studies of Idiopathic Mental Retardation: A Report

G. K. Chetan, K. R. Manjunatha, H. N. Venkatesh, S. Balu, E. Venkataswamy and S. Roy

*Department of Human Genetics, National Institute of Mental Health and Neuroscience,
Bangalore 560 029, Karnataka, India*

KEYWORDS Idiopathic Mental Retardation. Multiple Congenital Anomalies. Cytogenetics. Chromosomal Aberrations. FISH. Genes

ABSTRACT A majority of subjects with mental retardation and congenital anomalies cannot be classified under any known malformation syndromes. Major problems in understanding the mechanisms involved in the production of this abnormal phenotype and consequently in providing accurate and informed genetic counseling are therefore challenging. The application of recently developed chromosome banding techniques including the high resolution banding and FISH studies to population surveys has enabled accurate identification of individual chromosomes, specific existence and understanding of various chromosomal aberrations and their association with mental retardation. The present study was carried out to delineate any such chromosomal aberrations and their association in subjects with idiopathic mental retardation, multiple congenital malformations and various syndromes.

INTRODUCTION

The mental retardation (MR) of unknown cause is grouped under “idiopathic mental retardation” category. Idiopathic MR subjects show learning disabilities and borderline intelligence whose IQ range is 71-85.

Cytogenetic abnormalities among the subjects with Idiopathic Mental retardation with chromosomal abnormalities have been recorded; like 22q-13.3 deletion in language disability and speech in Autism (Manning et al. 2004), deletions in the 2qter and 6qter regions is seen in an idiopathic mental retarded subject by Li Rong and Zheng (2004).

In a study by Luciani et al. (2003), telomeric 22q13 deletion occurred as a result of a ring chromosome, simple deletion and translocations. The deletion was shown to be highly variable, ranging from 160 kb to 9 Mb. It was also shown that the parental origin of these deletions was more paternal (74%) than maternal (26%). A study by Sismani et al. (2001) detected an 8q subtelomeric deletion in an idiopathic mentally retarded subject by using FISH and MAPH Telomere Assay. A mild developmental delay and dysmorphism and very blue iris in a patient who

showed 15q24, q26 1.1 interstitial deletion was reported by Spruijt et al. (2003).

Idiopathic subjects also show chromosomal anomalies in the form of translocations. A *de novo* balanced translocation between 17p 13.3 and 20q 13.33 was identified by Walter et al. (2004). A study by Anderlid et al. (2002) identified one *de novo* unbalanced translocation and three unbalanced translocation inherited in a patient suffering with idiopathic mental retardation. Granzow et al. (2000), identified an unbalanced cryptic translocation der (5) t (3; 5) (q 27- p15.3) in a family with three cases of unexplained mental retardation and dysmorphic features using the Multiplex FISH Assay. Another unbalanced cryptic translocation between chromosomes 8 and 13: der (13) t (8; 13) (q 24.3- q34) in two sisters was identified by Kleefstra et al. in 2000.

FISH studies in 84 families with idiopathic mental retardation (Ewa Bocian et al. 2004) had revealed a large number of aberrations such as: 46 XY t(7,10) (q36 q26), 46 XY der(13) t(4,13) (p16 q34), 46 XY der(2) t(2,7) (q37 q36), 46 XX der(4) t(4,21) (p16 q22), 46 XY der(6) t(4,6) (q35 q27), 46 XX der(13) t(X, 13) (q28 q34), 46 XY der(10) t(10,19) (q26, p13.3), 46 XY del (4) (p16.1 p16.3).

Cryptic unbalanced chromosomal rearrangements in telomeric bands of human chromosome constitute a significant cause of idiopathic mental retardation. This was supported by numerous investigations, namely Ghaffari et al. (1998), Coco and Penchaszadeh (1982), Flint et al. (1995) Arunkumar (1998), Baker et al. (2002).

Corresponding author:

Dr. G. K. Chetan

Associate Professor

Department of Human Genetics, NIMHANS

Bangalore 560 029, Karnataka, India

Telephone: 091-080-26995121/5782

E-mail: drchetang@gmail.com

Keeping in view the literature on genetic studies in idiopathic mental retardation, the present investigation comprises the study of 100 mental retardation subjects. The report shows that 12 of the subjects had subtle chromosomal defects and details of the genetic implications are discussed in brief (Venkatesh 2006).

MATERIALS AND METHODS

100 patients consisting of 76 males and 24 females in the age group between 1-30 years belonging to different parts of India were selected and included in the study on the basis of mental retardation, multiple congenital malformations, various syndromes and anomalies of the central nervous system (CNS). All patients with syndromes of known gross chromosomal abnormalities like, Down syndrome, Klinefelter syndrome, Turner syndrome and patients with biochemical disorders were excluded from the present study.

Among the one hundred patients with unknown cause of mental retardation, multiple congenital malformation and syndromes were selected for the present study, 60 (50 males and 10 females) patients had mild degree of mental retardation, 25 (16 males and 9 females) patients had moderate degree of mental retardation, 15 (10 males and 5 females) have multiple congenital malformations (Table 1). These subjects were selected from in-patient and out-patient services of Mental Retardation clinic and from the Child Guidance Clinic (CGC) unit at NIMHANS, Bangalore, India. Since the evaluation of mental retardation in young infants was difficult, patients less than one year of age were not included in the present study. Among those included between one year and thirty years of age, a definite abnormality of the CNS was readily identified apart from some degree of mental retardation, multiple congenital malformations and syndromes.

Fifty intellectually and phenotypically normal, healthy individuals from the same age group were selected to serve as controls.

The chromosome preparations were made from peripheral blood lymphocytes using the standard method of Arakaki and Sparkes (1963) modified by Manjunatha (1988). The slides were GTG banded using Seabright's (1971) method of chromosome banding. Karyotypes were made according to the classification adopted by the Paris conference (1971).

For the Fluorescent In Situ Hybridization

(FISH) analysis the cultures were setup similar to those for GTG banding. Both centromeric and Locus Specific Indicator (LSI) probes (Vysis) were used. The denatured probe was applied to the marked region on the slides and a coverslip was placed over the slide. The slides were placed in the Hybridization chamber (HYBRITE, Vysis) overnight at 37°C. The slides were counter-stained with DAPI or PI and observed under Leica DM LB2 Microscope, using the appropriate filters. For the Interphase nuclei a minimum of 100 cells for good signals and at least 5 images were captured. For metaphase spread, a minimum of 15 spreads were analyzed and 3 images were captured.

RESULTS

The cytogenetic investigations were done in 100 subjects with idiopathic mental retardation, multiple congenital malformations and 50 controls. Among the subject group, males and females between the age group 1-30 yrs were selected and studied. A systematic survey on the presence of any subtle chromosomal aberrations revealed that 12 subjects have chromosomal structural anomalies (Table 2), such as, translocations, deletions, inversions, and ring chromosome formations. These findings were compared with the data on the control population.

Among the 12 subjects positive for various chromosomal anomalies, 8 were males aged between 2 1/2 to 17 years and 4 were females aged 3 to 23 years.

Out of the 12 positive families studied, 1 family showed first degree consanguinity, the rest showing no consanguineous union. In these 12 families, one family had two sibs affected with the same chromosome anomaly in their karyotype.

DISCUSSION

The findings varied from inversions of the chromosomal regions, deletions, ring formations,

Table 1: Degree of M.R. / MCA syndromes among the patient Groups (n = 100)

Abnormality	Age group	Males	Females	Total
Mild M.R	1-30 yrs	50	10	60
Moderate M.R	1-30 yrs	16	9	25
MCA	1-30 yrs	10	5	15
Total	1-30 yrs	76	24	100

Table 2: Different chromosomal anomalies found among the positive subjects in the present study (n=12) (see Fig 1 and 2)

Case No	Chromosomal anomaly	Sex	Age(in years)	Clin. Status / Diagnosis
1	del(15)(q11q13)	Male	8	Moderate MR/PWS
2	del(15)(q11q13)	Male	10	Mild MR/PWS
3	del(15)(q11q13)	Male	13	Severe MR
4	inv(9)(p13q21)	Male	4	Mild MR
5	inv(9)(p11q13)	Male	2 1/2	MCA
6	inv(9)(p11q13)	Male	9	Severe MR
7	inv(X)(q13q27)	Female	17	Mild MR
8	del(X)(pter→q21)	Female	16	Moderate MR
9	r(15)	Male	6	Mild MR / MCA
10	r(22)	Male	12	Mild MR
11	t(1;7)(p21;q11.2)	Female	3 1/2	Severe MR
12	t(3;10)(q28.2;q21.1)	Female	16	Moderate MR / MCA

Table 3: The percentage of chromosomal abnormalities reported previously by various authors in subjects with mental retardation.

Number	Investigation	Year	Subjects studied (n)	Subjects with chromosomal anomalies (n)	%
1.	Lubs and Lubs	1973	54	4	7.4
2.	Jacobs PA	1978	475	57	12.0
3.	Faed et al	1979	756	103	13.6
4.	Gripenberg et al	1980	1062	349	32.8
5.	Kondo et al	1980	449	37	8.1
6.	Narahara K	1981	74	11	14.9
7.	Nelson and Smart	1982	720	148	20.5
8.	Rasmussen et al	1982	1905	354	18.5
9.	Ambani et al	1984	709	248	35.0
10.	Fryns et al	1984	1991	424	21.3
11.	Reddy and Thomas	1985	200	39	19.5
12.	Kanata S	1986	121	14	11.6
13.	Kanata S	1986	56	9	16.1
14.	Hedge et al.	1989	1314	312	23.7
15.	Latha P	1996	100	18	18.0
16.	Bhaskar Rao GV	1999	200	40	20.0
17.	Hong et al	1999	604	69	11.3
18.	Riegel et al	2001	254	15	5.9
19.	Anderlid et al	2002	111	10	9.0
20.	Koolen et al	2004	210	14	6.7
21.	Soggard et al	2005	132	9	7.0
22.	Present Study	2007	100	12	12.0

translocations to various chromosomal regions (Table 2, Fig 1a-e and 2). Previous studies by various authors (Table 3) have also found out the association between the chromosomal anomalies and mental retardation. Multiple congenital anomalies, as a separate group do not specifically associate with any chromosomal anomaly. However, when mental retardation is also present there is an association with chromosomal anomalies. Although various syndromes have so far been associated with a "specific chromosomal anomaly" as in the case of Prader-Willi syndrome, del 15 (p11-13), there are still a lot more syndromes for which a definite chromo-

somal anomaly is not attributed. This study has tried to find out any such association and was successful to some extent.

Hence, it can be thought that, the association of these various chromosomal abnormalities with idiopathic mental retardation, congenital malformations and syndromes might have a definite etiology, but their implications in phenotype-genotype correlations are not well understood. However, they are useful for management, prevention, treatment amongst the high risk family members.

The application of the technique of high resolution banding, coupled with FISH analysis




Fig. 1a. Cut-out showing invX (Case No. 7)



Fig. 1b. Cut-out showing r15 (Case No. 9)



Fig. 1c. Cut-out showing r22 (Case No. 10)




Fig. 1d. Cut-out of 1 and 7 chromosome showing breakpoints of translocation $t(1;7)(p21;q11.2)$ (Case No. 11)




Fig. 1e. Cut-out of 3 and 10 chromosome with translocation breakpoints $t(3;10) (q28.2; q21/1)$ Case No. 12




Fig. 2. FISH image showing 15(q11-13) microdeletion in PWS (Case No. 1)

if applied to the routine cytogenetic screening programmes, would definitely help in identifying the subtle chromosomal aberrations, which otherwise could be missed in the normal conventional banding methods. Specialized techniques like PCR probe analysis and chromosome painting, CGH, CGH array and Multicolor FISH analysis would definitely provide us with the full molecular details of the disease under consideration and would be of great help in establishing

the proper phenotype-genotype correlations. This helps us in providing accurate genetic counseling to the affected and also to the members of the families at high-risk, also for a better management and therapeutic regimen.

ACKNOWLEDGMENTS

We thank the Director and CGC of NIMHANS, Bangalore. Financial support from

UGC and ICMR, New Delhi is greatly acknowledged. We are thankful for the patients and families for their co-operation.

REFERENCES

- Ambani LM, Patel ZM, Dhareshwar SS, Krishnamurthy DS, Moghe MS, Mulye VR, Gokarn VV, Peter JJ, Shah LP Merchant, SM 1984. Clinical, biochemical and cytogenetic studies in mental retardation. *Indian J Med Res*, 79: 384-391.
- Anderlid BM, Schoumans J, Anneren G, Sahlen S, Kyllerman M, Vujic M, Hagberg B, Blennow E Nordenskjold M 2002. Subtelomeric rearrangements detected in patients with idiopathic mental retardation. *Am J Med Genet*, 107(4): 275-284.
- Arakaki DT, Sparkes, RS 1963. Microtechnique for Culturing Leukocytes from Whole Blood. *Cytogenetics*, 85: 57-60.
- Arunkumar 1998. Molecular Characterization and Delineation of Subtle Deletion in de novo balanced Chromosomal rearrangements. *Human Genetics*, 103(2): 173.
- Baker E, Hinton L, Callen DF, Aitree M, Dobbie A, Eyre HJ, Sutherland GR, Thompson E, Thompson P, Woollatt E, Haan E 2002. Study of 250 children with idiopathic mental retardation reveals nine cryptic and diverse subtelomeric chromosome anomalies. *Am J Med Genet*, 107(4): 285-93.
- Bhaskar Rao GV 1999. *A Systematic and Comparative Study on the Association and Significance of Chromosomal Aberrations in Subjects with in Idiopathic Mental Retardation, Congenital Malformations and Syndromes*. PhD Thesis. Submitted to NIMHANS (Deemed University), Bangalore.
- Coco R, Penchaszadeh VB 1982. Cytogenetic findings of 200 children with mental retardation and multiple congenital anomalies of unknown cause. *Am J Med Genet*, 12(2): 155.
- Ewa Bocian, Zofia Helias-Rodzewicz, Kamila Suchenek, Ewa Obersztyn, Anna Kutkowska-Kazmierczak, Powel Stankiewicz, Ewa Kostyk, Tadeusz Mazurczak 2004. Subtelomeric rearrangements: results from FISH studies in 84 families with idiopathic mental retardation. *Medical Science Monitor*, 10(4): CR 143-151.
- Faed MJ, Robertson J, Field MA, Mellon JP 1979. A chromosome survey of a hospital for the mentally subnormal. *Clin Genet*, 16: 191.
- Flint J, Wilkie AOM, Buckle VJ, Winter RM, Holland AJ, McDermid HE 1995. The detection of subtelomeric chromosomal rearrangements in idiopathic mental retardation. *Nat Genet*, 9: 132-140.
- Fryns JP, Timmermans J, D'Hondt F, Francois B, Emmery L, Van Den Berghe 1984. Cytogenetic findings in moderate and severe mental retardation, a study of an institutionalised population of 1991 patients. *Acta Ped Scand* (Supplement), 313: 3-23.
- Ghaffari SR, Boyd E, Tolmie JL, Crow YJ, Trainer AH, Connor JM 1998. A new strategy for cryptic telomeric translocation screening in patients with idiopathic mental retardation. *J Med Genet*, 35(3): 225-233.
- Granzow M, Popp S, Keller M, Holtgreve-grez H, Brough M, Schoell B, Raterberg-Ruland I, Hager HD, Tariverdian G, Jauch A 2000. Multiplex FISH telomere integrity assay identifies an unbalanced cryptic translocation der (5)t (3;5) (q27;p15.3) in a family with three mentally retarded individuals. *Hum Genet*, 107(1): 51-57.
- Gripenberg U, Hongell K, Knuutila S, Kahkonen M, Leisti J 1980. A chromosome survey of 1062 mentally retarded patients. Evaluation of a long-term study at the Rinnekoti institution, Finland. *Hereditas*, 92: 223-8.
- Hegde S, Kalyankar GD, Thomas IM, Ambani LM, Dareshwar SS, Patil SM, Menon PSS, Thomas S, Verma IC, Agarwal SS, Das SK, Pandey VK, Singh UR, Sithole P, Ahluwalia TP, Arora R, Saxena NC, Saxena BN 1989. *Multicentric Task Force Study on Identification of Genetic Cause of Mental Retardation*. New Delhi: ICMR.
- Hong KE, Kim JH, Moon SY, Oh SK 1999. Chromosomal abnormalities in child psychiatric patients. *J Korean Med Sci*, 14(4): 377-85.
- Jacobs PA 1978. A Cytogenetic survey of an institution for the mentally retarded I. Chromosome abnormalities. *Clin Genet*, 13(1): 37.
- Kanata S 1986. A chromosomal study on 56 cases with heavy mental' retardation and behavior disorders. *Proc Japan Acad*, 62 B: 165-168.
- Kanata S 1986. A chromosomal study on 121 cases with heavy mental retardation. *Proc Japan Acad*, 62B: 314-316.
- Kleefstra T, Van de Zande G, Merkx G, Mieloo H, Hoovers JM, Smeets D 2000. Identification of an unbalanced cryptic translocation between the chromosomes 8 and 13 in two sisters with mild mental retardation accompanied by mild dysmorphic features. *Eur J Hum Genet*, 8(8): 637-640.
- Kondo I, Hamaguchi H, Nakajima S, Haneda T 1980. A cytogenetic survey of 449 patients in a Japanese institution for the mentally retarded. *Clin Genet*, 17: 177.
- Koolen DA, Nillesen WM, Versteeg MH, Merkx GF, Knoers NV, Kets M, Vermeer S, van Ravenswaaij CM, de Kovel CG, Brunner HG, Smeets D, de Vries BB, Sistermans EA 2004. Screening for subtelomeric rearrangements in 210 patients with unexplained mental retardation using multiplex ligation dependent probe amplification (MLPA). *J Med Genet*, 41(12): 892-899.
- Latha P 1996. *Cytogenetics of Autosomal Fragile Sites: Association and Their Implications in Subjects with Mental Retardation*. Ph. D Thesis, submitted to NIMHANS (Deemed University). Bangalore.
- Li-Rong Aieu, Zhad Zheng-Yanoyno 2004. Two subtelomeric chromosomal deletions in forty six children with idiopathic mental retardation. *Chinese Medical Journal*, 11(9): 1414-1417.
- Lubs HA, Lubs ML 1973. New cytogenetic techniques applied to a series of children with mental retardation. *Ann Genet, Nobel Symposium*, 23.
- Luciani JJ, de Mas P, Depetris D, Mignon-Ravix C, Bottani A, Prieur M, Jonveaux P, Philippe A, Bourrouillou G, de Martinville B, Delobel B, Vallee L, Croquette MF, Mattei MG 2003. Telomeric 22q13 deletions resulting from rings, simple

- deletions, and translocations: cytogenetic, molecular, and clinical analyses of 32 new observations. *J Med Genet*, 40(9): 690-696.
- Manjunatha KR, Rao BSS, Narayanan HS, Girimaji SR, Shobha S, Gandhi DH, Sangvi R 1988. Fragile X syndrome: The first case report from India. *Genome*, 30(1): 205.
- Manning MA, Cassidy SB, Clericuzio C, Cherry AM, Schwartz S, Hudgins L, Enns GM, Hoyne HE 2004. Terminal 22q deletions syndrome: A newly recognized cause of speech and language disability in Autism spectrum. *Pediatrics*, 114(2): 451.
- Narahara K 1981. A cytogenetic study of children with clinically unclassifiable multiple congenital malformations and mental retardation. *Acta Med Okayama*, 35(5): 343-355.
- Nelson MM, Smart RD 1982. The results of chromosome examinations in an institution for mental retardates in the Cape Province. *S Afr Med J*, 62: 25.
- Paris Conference 1971 and S1975. *Standardisation in Human Cytogenetics*. Birth defects; Original Article Series, XI: 9. New York: The National Foundation.
- Rasmussen K, Nielsen J, Dahl G 1982. The prevalence of chromosome abnormalities among mentally retarded persons in a geographically delimited area of Denmark. *Clin Genet*, 22: 244.
- Reddy KS, Thomas IM 1985. Cytogenetic investigations in mentally retarded patients. *Indian J Med Res*, 82: 164-170.
- Riegel M, Baumer A, Jamar M, Delbecque K, Herens C, Verloes A, Schinzel A 2001. Submicroscopic terminal deletions and duplications in retarded patients with unclassified malformation syndromes. *Hum Genet*, 109(3): 286-94.
- Seabright M 1971. A rapid banding technique for Human chromosomes. *Lancet*, II: 971-972.
- Sismani C, Armour JA, Flint J, Girgalli C, Regan R, Patsalis PC 2001. Screening for subtelomeric chromosome abnormalities in children with idiopathic mental retardation using multiprobe telomeric FISH and the new MAPH telomeric assay. *Human Genetics*, 9(7): 527-32.
- Spruijt L, Engelen JJM, Bruinen-Smeijsters IP, Albrechts JCM, Schrander J, Schrander-Stumpel CTRM 2004. A Patient with a de novo 15q24q26.1 interstitial deletion, developmental delay, mild dysmorphism and very blue irises. *Am J Med Genet*, 129(3): 312-315.
- Sogaard M, Tümer Z, Hjalgrim H, Hahnemann J, Friis B, Ledaal P, Pedersen VF, Baekgaard P, Tommerup N, Cingöz S, Duno M, Brondum-Nielsen K 2005. Subtelomeric study of 132 patients with mental retardation reveals 9 chromosomal anomalies and contributes to the delineation of submicroscopic deletions of 1pter, 2qter, 4pter, 5qter and 9qter. *BMC Med Genet*, 17: 6-21.
- Venkatesh HN 2006. *Genetics of Mental Retardation: Molecular Cytogenetic Studies among Idiopathic Mental Retardation*. Thesis Submitted to NIMHANS (Deemed University), Bangalore.
- Walter S, Sandig K, Hinkel GK, Mitulla B, Ounap K, Sims G, Sitska M, Utermann B, Viertel P, Kalscheuer V, Bartsch O 2004. Subtelomere FISH in 50 children with mental retardation and minor anomalies, identified by a checklist, detects 10 rearrangements including a de-novo balanced translocation of chromosomes 17p13.3 and 20q13.33. *Am J Med Genet*, 128A: 364-73.