

## Beta-Globin Gene Haplotypes in Manipur, North-east India

\*Maishnam Rustam Singh<sup>1</sup>, B. Choudhury<sup>1</sup>, T. Shyamacharan Singh<sup>2</sup> and V.R. Rao<sup>3</sup>

<sup>1</sup>*Department of Anthropology, Gauhati University, Guwahati 781 014, Assam, India*

<sup>2</sup>*Department of Anthropology, Manipur University, Imphal 795 003, Manipur, India*

<sup>3</sup>*Anthropological Survey of India, 27, JN Road, Kolkata 700 016, West Bengal, India*

**KEYWORDS** Naga and Kuki Tribes. Hb $\beta$ EE. Genetic Affinity. Polymorphic Restriction Sites. Framework (F.W.)

**ABSTRACT** A sample of 204 individuals belonging to five different population units of Manipur, viz. the Meitei, Gangte, Thadou, other Kuki, and Kabui has been analyzed for beta globin gene cluster, considering seven loci to bring out a first hand information on haplotype frequencies of beta globin in the state. A common trend of allelic frequencies observed in the present data are the high frequencies of HindIII $\epsilon$ , AvaII $\beta$  and HinfI $\beta$  and relatively low frequencies of HindIII $\gamma$ , HindIII $\alpha$ , HindIII $\psi$   $\beta$  and HindIII $\psi$   $\beta$ . The frequency of Framework 1 (presence of both AvaII $\beta$  and HinfI $\beta$  sites) ranges from nil to 17.65% in the present data of Manipur. The frequency of Framework 2 ranges from 33% to 66% and that of Framework 3 from 33% to 55%. One noteworthy observation is the association of Hb $\beta$ EE exclusively with Framework 2, which suggests a similar mutational origin of HbE in South-east Asia and North-east India. Only five beta globin haplotypes are found in the present data. Haplotypes +—+—+ and +—+—+—, the most common in Japanese, Cambodians, Koreans, South Chinese, Samoans and Thais, are the most frequent haplotypes in Manipur too.

### INTRODUCTION

The beta-globin gene cluster associated with haemoglobin S, E, and beta-thalassemia genes has been studied as a means of searching affinities and common origin among human populations (Antonarakis et al. 1982; Kazazian et al. 1984; Chan et al. 1986; Wainscoat et al. 1986; Hundrieser et al. 1988; Currat et al. 2002). Distribution of haplotypes associated with the  $\beta$ A gene is now known for a number of populations. The data indicate that Africans are the most diverged group, and that the first split separated an African group from an Eurasian one (Antonarakis et al. 1985; Oehme et al. 1985; Maggio et al. 1986; Wainscoat et al. 1986; Ramsay and Jenkins 1987; Shimizu 1987; Hundrieser et al. 1988 a and b; Shimizu et al. 1989; Yongvanit et al. 1989; Chen et al. 1990; Long et al. 1990; Trent et al. 1990; Varawalla et al. 1992; Hewitt et al. 1996; Fucharoen et al. 1997; Kaufman et al. 1998; Villalobos-Arámbula et al. 2000). In north-eastern part of India, Hundrieser et al. (1988) studied the beta globin gene cluster in Assam. The present paper brings out a firsthand information on the frequencies of beta globin gene cluster haplotypes in Manipur.

### AREA AND PEOPLE

The state of Manipur lies between the 23.83° and 25.68° N latitude and 93.03° and 94.78° E longitude and has an area of 22,327 sq. km. of which only 8% is the valley. The most populous community of the state is the Meiteis who mainly occupy the valley while the hills are mainly inhabited by two tribal groups, viz. the Kuki and Naga. There are 33 scheduled tribes, who have their own distinct culture and dialects. The present study comprises the Meitei (the major community of Manipur), Thadou and Gangte (as representatives of Kuki tribe), other Kuki tribes and the Kabui (as a representative of Naga tribe).

### MATERIAL AND METHODS

About 4ml of intravenous blood samples from 204 healthy and unrelated individuals from the Meitei (75), Gangte (35), Thadou (45), Kabui (12) tribe and other tribes belonging to Kuki group (37) were collected in vacutainer, containing EDTA after obtaining their written consent. The blood samples were collected during July 2006 and April 2007. The Kabui samples were collected from the Tamenglong district, the Gangte, Thadou and other groups belonging to Kuki were collected from Churachandpur district, and the Meitei from Imphal east and west districts of Manipur. Analyses work was done in the DNA Laboratory of Anthropological Survey of India.

---

\*Address Correspondence to:  
Dr. Maishnam Rustam Singh,  
Khagempalli Panthak,  
Imphal West, 795 001, Manipur, India  
Mobile: +919859423382,  
E-mail: mrluwang@yahoo.co.in

DNA was extracted by phenol-chloroform methods (Sambrook et al. 1989) with minor modifications and the seven restriction sites for beta globin gene cluster (HindII $\epsilon$ , HindIIIG $\gamma$ , HindIIIA $\gamma$ , HindII5' $\psi\beta$ , HindII3' $\psi\beta$ , AvaII $\beta$  and HinfI $\beta$ ) were amplified using the oligonucleotide primers (Weatherall and Clegg 2001) and the products were digested by respective endonucleases. The digested products were separated in agarose gels containing ethidium bromide. The presence or absence of specific bands was then verified under UV- transilluminator, using appropriate molecular weight marker. The frameworks were identified by the presence (+) or absence (-) of the AvaII restriction site in intron 2 of the beta globin gene and of the BhamHI site located 3' to the beta globin gene. The BhamHI RFLP is located within an L1 repetitive element creating amplification problems and a HinfI RFLP located just 3' to the beta globin gene is used instead, because these two RFLPs have been found to exist in linkage disequilibrium (Semenza et al. 1989). Presence of both restriction sites (++) indicates framework 1, presence of AvaII site together with the absence of the HinfI site indicates framework 2 and the absence of AvaII site and presence of HinfI site indicate framework 3 (Antonarakis et al. 1982). Calculation of allelic frequencies of the seven restriction sites and estimation of haplotype

frequencies were estimated following Nei (1987) and Long (1995).

## RESULTS AND DISCUSSION

Gene frequencies of the seven polymorphic restriction sites in the beta globin gene cluster of the present sample of five populations of Manipur are shown in Table 1. A common trend of allelic frequencies observed in these populations is the high frequencies of HindII $\epsilon$ , AvaII $\beta$  and HinfI $\beta$  and relatively low frequencies of HindIIIG $\gamma$ , HindIIIA $\gamma$ , HindII5' $\psi\beta$ , and HindII3' $\psi\beta$ . The frequency of the presence of the HindII $\epsilon$  site in the populations increases from 0.60 in the Thadou to 0.75 in the Kabui, through other Kukis (0.608), Meitei (0.66) and the Gangte (0.729). Frequency of the presence of AvaII $\beta$  site is greatest among the Kabui (0.583) followed successively by the Gangte (0.571), Meitei (0.567), Thadou (0.544) and other Kukis (0.514). The frequency of the presence of HinfI $\beta$  is also above 57% in all the other populations except the Kabui who shows only 45.8%. Sample size of the Kabui, of course, is the smallest, only 12 individuals. On the other hand, the greatest frequency of presence of HindIIIG $\gamma$  site is 0.40, shown by the Thadou, which is followed by other Kukis (0.365), Meitei (0.34), Gangte (0.271) and the Kabui (0.208). The frequency of HindIIIA $\gamma$  ranges from 0.25

**Table 1: Allelic frequencies of seven beta globin loci in five populations of Manipur**

|                      | Meitei<br>(N=75) | Gangte<br>(N=35) | Thadou<br>(N=45) | Other Kuki<br>(N=37) | Kabui<br>(N=12) | Total<br>(N=204) |
|----------------------|------------------|------------------|------------------|----------------------|-----------------|------------------|
| HindII $\epsilon$    |                  |                  |                  |                      |                 |                  |
| —                    | 0.34             | 0.271            | 0.4              | 0.392                | 0.25            | 0.352            |
| ++                   | 0.66             | 0.729            | 0.6              | 0.608                | 0.75            | 0.648            |
| HindIIIG $\gamma$    |                  |                  |                  |                      |                 |                  |
| —                    | 0.66             | 0.729            | 0.6              | 0.635                | 0.792           | 0.654            |
| ++                   | 0.34             | 0.271            | 0.4              | 0.365                | 0.208           | 0.346            |
| HindIIIA $\gamma$    |                  |                  |                  |                      |                 |                  |
| —                    | 0.74             | 0.757            | 0.667            | 0.689                | 0.75            | 0.716            |
| ++                   | 0.26             | 0.243            | 0.333            | 0.311                | 0.25            | 0.284            |
| HindII5' $\psi\beta$ |                  |                  |                  |                      |                 |                  |
| —                    | 0.713            | 0.814            | 0.722            | 0.811                | 0.75            | 0.753            |
| ++                   | 0.287            | 0.186            | 0.278            | 0.189                | 0.25            | 0.247            |
| HindII3' $\psi\beta$ |                  |                  |                  |                      |                 |                  |
| —                    | 0.647            | 0.729            | 0.6              | 0.635                | 0.75            | 0.648            |
| ++                   | 0.353            | 0.271            | 0.4              | 0.365                | 0.25            | 0.352            |
| AvaII $\beta$        |                  |                  |                  |                      |                 |                  |
| —                    | 0.433            | 0.429            | 0.456            | 0.486                | 0.417           | 0.448            |
| ++                   | 0.567            | 0.571            | 0.544            | 0.514                | 0.583           | 0.552            |
| HinfI $\beta$        |                  |                  |                  |                      |                 |                  |
| —                    | 0.353            | 0.429            | 0.378            | 0.338                | 0.542           | 0.370            |
| ++                   | 0.647            | 0.571            | 0.622            | 0.662                | 0.458           | 0.630            |

in the Kabui to 0.333 in the Thadou. Presence of HindII5' $\psi$  $\beta$  ranges from 0.187 in the Gangte to 0.287 in the Meitei and that of HindII3' $\psi$  $\beta$  ranges from 0.25 in the Kabui to 0.365 of Other Kukis.

Frequencies of Beta-globin gene frameworks (F.W.) are displayed in Table 2. Presence and absence in homozygous condition of the beta polymorphic restriction sites *Ava*II $\beta$  and *Hinf*I $\beta$  determines the frameworks. It appears that F.W. 3, the absence of *Ava*II $\beta$  and presence of *Hinf*I $\beta$  sites, shows greatest in other Kuki tribes (0.5556) followed by the Meitei (0.50), Thadou (0.4706), Gangte (0.4615) and the Kabui (0.3333). The frequency of F.W. 2 (presence of *Ava*II $\beta$  and absence of *Hinf*I $\beta$ ) ranges from 0.3333 in the Meitei and Other Kukis to 0.6667 in the Kabui, through the Thadou (0.352) and Gangte (0.538). It is noteworthy that five heterozygous HbAE cases and all the homozygous HbEE cases were met exclusively in F.W.2. The beta thalassaemia carriers were, however, found to be associated with F.W. 3. Frequency of F.W.1 (presence of both *Ava*II $\beta$  and *Hinf*I $\beta$  sites) ranges from nil to 17.65% in the populations of Manipur, which is notably lower than the earlier reports of Asian Indians with 52% (Antonarakis et al. 1982; Kazazian et al. 1984; Old et al. 1984; Hundrieser et al. 1988; Varawalla et al. 1992). The high frequencies of F.W.2 and F.W.3 of the present data need further investigation. In a pioneering work, Antonarakis et al. (1982) reported the evidence of two different Hb $\beta$ E associated DNA frameworks in subjects from Cambodia and Thailand. F.W. 3 was exclusively found in Cambodians, suggesting an independent mutational origin of HbE in this population. Other framework of HbE, F.W. 2 was found among the Thai, Laotian and also among some Cambodians. Such an investigation, conducted by Hundrieser et al. in 1988 among the Kachari of Assam did not provide evidence for a separate mutational origin of Hb $\beta$ E in Assam, for the Kachari subjects were,

by and large, characterized by F.W. 2. It is pertinent to mention that five heterozygous HbAE cases and all the homozygous HbEE cases were met exclusively in F.W.2. It suggests a common mutational origin of HbE in South-east Asia and north-east India. At the present state of knowledge, the presumption of 'Austroasiatic speaking people as the original owner of Hb $\beta$ E' (Flatz 1967; Das et al. 1971; Flatz et al. 1972) is neither supported by the available evidences, nor the high frequency of Hb $\beta$ E in north-east region can be explained by a possible independent mutational origin of this gene.

**Table 2: Frequencies of the beta-globin gene frameworks in five populations of Manipur**

| Ethnic groups | Chromosomes | Framework 1 | Framework 2 | Framework 3 |
|---------------|-------------|-------------|-------------|-------------|
| Meitei        | 72          | 0.167       | 0.333       | 0.500       |
| Gangte        | 26          | 0.000       | 0.539       | 0.462       |
| Thadou        | 34          | 0.176       | 0.353       | 0.471       |
| Other Kuki    | 36          | 0.111       | 0.333       | 0.556       |
| Kabui         | 6           | 0.000       | 0.667       | 0.333       |
| Total         | 174         | 0.1264      | 0.3793      | 0.4943      |

The beta-globin gene cluster haplotype frequencies in five populations of Manipur are displayed in Table 3. Altogether five beta globin haplotypes are found so far in Manipur. Haplotype +—+—, the most frequently encountered in Japanese, Cambodians, Koreans, South Chinese, and Samoans (Antonarakis et al. 1982; Cheng et al. 1984; Chan et al. 1986; Liu et al. 1987; Shimizu 1987; Hundrieser et al. 1988; Shimizu et al. 1989; Trent et al. 1990; Shimizu et al. 1992;), is also found to be most common type in the populations of Manipur. Its frequency is highest in the other Kuki tribes (0.625) followed by the Gangte (0.50), Thadou (0.50) and the Meitei (0.454). Among the Kabui the haplotype +—+—, the most common in Thais (Hundrieser et al. 1988; Yongvanit et al. 1989), is the only representing type. The frequency of haplotype +—+— in the remaining popula-

**Table 3: Beta globin haplotype frequencies in five populations of Manipur**

| Haplotypes | Polymorphic sites |   |   |   |   |   |   | F.W. | Meitei | Gangte | Thadou | Other Kuki | Kabui | Total |
|------------|-------------------|---|---|---|---|---|---|------|--------|--------|--------|------------|-------|-------|
|            | 1                 | 2 | 3 | 4 | 5 | 6 | 7 |      |        |        |        |            |       |       |
| A          | +                 | - | - | - | - | + | + | 1    | 0.182  | 0.000  | 0.250  | 0.000      | 0.00  | 0.085 |
| B          | +                 | - | - | - | - | + | - | 2    | 0.273  | 0.500  | 0.000  | 0.375      | 1.00  | 0.371 |
| C          | -                 | + | + | + | + | + | - | 2    | 0.000  | 0.000  | 0.250  | 0.000      | 0.00  | 0.029 |
| D          | +                 | - | - | - | - | - | + | 3    | 0.454  | 0.500  | 0.500  | 0.625      | 0.00  | 0.486 |
| E          | +                 | - | - | + | - | - | + | 3    | 0.091  | 0.000  | 0.000  | 0.000      | 0.00  | 0.029 |

tions ranges from nil among the Thadou to 0.50 in the Gangte through 0.273 in the Meitei, 0.375 in Other Kuki. The haplotype -++++- is encountered exclusively among the Thadou tribe with a frequency of 0.25 while the haplotype +—+—+ is found only in the Meitei population (0.091). The Meitei and Thadou exhibit the haplotype +—+—++ frequencies of 0.182 and 0.25 respectively. This haplotype +—+—++ is most frequently encountered in Colombian Amerinds, Mediterraneans, Asian Indians, and Mexican Mestizos (Shimizu 2004).

To conclude, it may be recapitulated that the beta-globin gene cluster haplotypes in Manipur show closer similarity with the people of east and south-east Asian populations. A common mutational origin of Hb $\beta$ E in north-east Indian and the South-east Asian populations is also expected.

## ACKNOWLEDGEMENT

Authors acknowledge the financial supports from the University Grant Commission, New Delhi under the UGC Dr. D.S. Kothari Post Doctoral Fellowship Scheme and Anthropological Survey of India, Ministry of Culture, Govt. of India.

## REFERENCES

- Antonarakis SE, Boehm CD, Giardina PJV, Kazazian HH Jr 1982. Non-random association of polymorphic restriction sites in the  $\beta$ -globin gene cluster. *Proc Natl Acad Sci USA*, 79: 137-141.
- Antonarakis SE, Kazazian HH Jr, Orkin SH 1985. DNA polymorphism and molecular pathology of the human globin gene clusters. *Hum Genet*, 69: 1-14.
- Chan V, Chan TK, Cheng MY, Leung NK, Kan YW, Todd D 1986. Characteristics and distribution of  $\beta$ -thalassaemia haplotypes in South China. *Hum Genet*, 73: 23-26.
- Chen LZ, Eastaugh S, Board PG, Kirk-f RL 1990. Evolution of beta-globin haplotypes in human populations. *Mol Biol Evol*, 7: 423-437.
- Cheng TC, Orkin SH, Antonarakis S, Potter MJ, Sexton JP, Markham AF, Giardina OJV, Li A, Kazazian HH Jr 1984. Beta-thalassaemia in Chinese: Use of in vivo RNA analysis and oligonucleotide hybridization in systematic characterization of molecular defects. *Proc Natl Acad Sci USA*, 81: 2821-2825.
- Curat M, Trabuchet G, Rees D, Perrin P, Harding RM, Clegg JB, Langaney A, Excoffier L 2002. Molecular analysis of the beta-globin gene cluster in the Niokholo Mandenka population reveals a recent origin of the beta (S) Senegal mutation. *Am J Hum Genet*, 70: 207-223.
- Das BM, Chakravarti MR, Delbruck H, Flatz G 1971. High prevalence of haemoglobin E in two populations in Assam. *Humangenetik*, 12: 264-266.
- Flatz G 1967. Hemoglobin E: Distribution and population dynamics. *Humangenetik*, 3: 189-234.
- Flatz G, Chakravarti MR, Das BM, Delbruck H 1972. Genetic survey in the population of Assam I. ABO blood groups, glucose-6-phosphate dehydrogenase and haemoglobin type. *Hum Hered*, 22: 323-330.
- Fucharoen S, Winichagoon P 2000. Clinical and hematologic aspects of hemoglobin E beta-thalassemia. *Curr Opin Hematol*, 7: 106-112.
- Hewitt R, Krause A, Goldman A, Campbell G, Jenkins T 1996. Beta-globin haplotype analysis suggests that a major source of Malagasy ancestry is derived from Bantu-speaking Negroids. *Am J Hum Genet*, 58: 1303-1308.
- Hundrieser J, Deka R, Gogoi BC, Papp T, Flatz G 1988a. DNA haplotypes and frameworks associated with the beta-globin gene in the Kachari population of Assam (India). *Hum Hered*, 38: 240-245.
- Hundrieser J, Sanguanserm Sri T, Papp T, Laig M, Flatz G 1988b. Beta-globin gene linked DNA haplotypes and frameworks in three South-East Asian populations. *Hum Genet*, 80: 90-94.
- Kaufman L, Carnese FR, Goicoechea A, Dejean C, Salzano FM, Hutz MH 1998. Beta-globin gene cluster haplotypes in the Mapuche Indians of Argentina. *Genet Mol Biol*, 21: 435-437.
- Kazazian HH Jr, Orkin SH, Antonarakis SE, Sexton JP, Boehm CD, Goff SC, Waber PG 1984. Molecular characterization of seven beta-thalassaemia mutations in Asian Indians. *EMBO J*, 3: 593-596.
- Liu VWS, Woo YK, Lam VMS, Huang CH, Chau AS, Lam STS, Wong HW, Tam JWO 1987. Molecular studies of beta-thalassaemic DNA of Chinese patients. *Birth Defects Orig Artic Ser*, 23: 87-92.
- Long JC, Chakravarti A, Boehm CD, Antonarakis SE, Kazazian HH Jr 1990. Phylogeny of  $\beta$ -globin gene haplotypes and its implications for recent human evolution. *Am J Phy Anthropol*, 81: 113-130.
- Long JC, Williams RC, Urbanek M 1995. An E-M algorithm and testing strategy for multiple locus haplotypes. *Am J Hum Genet*, 56: 799-810.
- Maggio A, Acuto S, Lo Gioco P, Di Marzo R, Giambona A, Sammarco P, Caronia F 1986.  $\beta^A$  and  $\beta^{thal}$  DNA haplotypes in Sicily. *Hum Genet*, 72: 229-230.
- Nei M 1987. *Molecular Evolutionary Genetics*. New York: Columbia University Press.
- Oehme R, Kohne E, Horst J 1985. DNA polymorphic patterns linked to  $\beta$ -globin genes in German families affected with hemoglobinopathies and thalassemias: A comparison to other ethnic groups. *Hum Genet*, 71: 219-222.
- Old JM, Petrou M, Modell B, Weatherall DJ 1984. Feasibility of antenatal diagnosis of  $\beta$ -thalassaemia by DNA polymorphisms in Asian Indian and Cypriot populations. *Br J Haematol*, 57: 255-263.
- Ramsay M, Jenkins T 1987. Globin gene-associated restriction-fragment length polymorphisms in southern African peoples. *Am J Hum Genet*, 41: 1132-1144.
- Sambrook J, Fritsch E, Maniatis T 1989. *Molecular Cloning: A Laboratory Manual*. New York, Cold Spring Harbor Laboratory.
- Semenza GL, Dowling CE, Kazazian HH Jr 1989. HinfI polymorphisms 3' to the human  $\beta$  globin gene detected by the polymerase chain reaction (PCR). *Nucl Acids Res*, 17: 2376.
- Shimizu K 1987. Characteristics of  $\beta^A$  chromosome haplotypes in Japanese. *Biochem Genet*, 25: 197-203.
- Shimizu K, Marubayashi A, Tokimasa K, Harihara S, Omoto K, Imanishi T, Hao L, Jin F 2004. Characteristic  $\beta$ -globin gene cluster haplotypes of Evenkis and Oroqens in North China. *Hum Biol*, 76: 765-778.
- Shimizu K, Park KS, Harihara S, Enoki Y 1992. Relation-

- ships between %HbF or %<sup>C</sup>g and the haplotypes in the  $\beta$ -globin gene cluster in the normal adult Japanese and Korean populations. *Hum Hered*, 42: 216-221.
- Shimizu K, Park KS, Omoto K 1989. The DNA polymorphisms of the  $\beta$ -globin gene cluster and the arrangements of the  $\alpha$ - and the  $\beta$ -globin genes in Koreans. *Hemoglobin* 13: 137-146.
- Trent RJ, Ferguson V, Hertzberg MS, Rutherford J, Mickleson KNP 1990.  $\beta$ -globin gene haplotypes in polynesians are predominantly southern Chinese type. *Hum Hered*, 40: 285-289.
- Varawalla NY, Fitches AC, Old JM 1992. Analysis of  $\hat{\alpha}$ -globin gene haplotypes in Asian Indians: Origin and spread of  $\beta$ -thalassaemia on the Indian subcontinent. *Hum Genet*, 90: 443-449.
- Villalobos-Arambula AR, Bustos R, Casas-Castaneda M, Gutierrez E, Perea FJ, Thein SL, Ibarra B 1997.  $\beta$ -thalassaemia and  $\beta^A$  globin gene haplotypes in Mexican mestizos. *Hum Genet*, 99: 498-500.
- Wainscoat JS, Hill AVS, Boyce AL, Flint J, Hernandez M, Thein SL, Old JM, Lynch JR, Falusi AG, Weatherall DJ, Clegg JB 1986. Evolutionary relationships of human populations from an analysis of nuclear DNA polymorphisms. *Nature*, 319: 491-493.
- Weatherall DJ, Clegg JB 2001. *The Thalassaemia Syndromes*. 4th Edition. USA: Blackwell Science Inc.
- Yongvanit P, Sriboonlue P, Mularlee N, Karnthong T, Areejitranusorn P, Hundrieser J, Limberg R, Schulze B, Laig M, Flatz SD, Flatz G 1989. DNA haplotypes and frameworks linked to the  $\beta$ -globin locus in an Austro-Asiatic population with a high prevalence of hemoglobin E. *Hum Genet*, 83: 171-174.