

Comprehensive Association Analysis of 10 Single Nucleotide Polymorphisms Associated With Osteoporosis among a Taiwanese Population

Chiao-Wen Hwang¹, Ching-Hsiang Lu², Shu-Fen Sun¹, Tzu-Ying Sung¹, Hsin-Yeh Chung¹, Shi-Ying Huang³, Han-Chun Hung³, Chun-Hong Chen^{3,4}, Yu-Min Sun^{3,4}, Yen-You Lin³, Wen-Sheng Liu⁵ and Zhi-Hong Wen^{3*}

¹*Department of Physical Rehabilitation & Medicine, Kaohsiung Veterans General Hospital, Kaohsiung 813, Taiwan*

²*Department of Neurosurgery; Kaohsiung Armed Forces General Hospital, Kaohsiung 802, Taiwan*

³*Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung 804, Taiwan*

⁴*Doctoral Degree Program in Marine Biotechnology, National Sun Yat-sen University, Kaohsiung 804, Taiwan*

⁵*Asia-Pacific Biotech Developing, Inc.*

KEYWORDS Osteoporosis. Bone Mineral Density. Single Nucleotide Polymorphisms. Taiwanese

ABSTRACT In the present study, we tested for an association between single nucleotide polymorphisms (SNPs) and bone mineral density (BMD) of the hip and lumbar spine in a Taiwanese population by analyzing 252 healthy persons (non-osteoporosis) and 193 persons with osteoporosis. we found that age; body mass index; family history; and consumption of coffee or vitamin D were associated with osteoporosis in our Taiwanese population. Our results also indicated that osteoprotegerin (*OPG*) SNP (rs6993813 T→C) and receptor activator of nuclear factor kappa-B ligand (*RANKL*) SNP (rs9594738 C→T) were significantly associated with BMD in our Taiwanese population. Additionally, we propose that the mean threshold value of integration of 7 wild-type SNPs (rs7524102 and rs6696981 of 1p36, rs11898505 of 2p16, rs9479055 of 6q25, rs326340 and rs1289759 of 3q13, and rs9594738 of 13q14) can be used for both a Taiwanese reference for bone density testing and to achieve the effect of grading.