

## **Gradients in Distribution of HLA – DRB1\* Alleles in Castes and Tribes of South India**

**K. Balakrishnan\*, C. Rathika\*, R. Kamaraj\*, R. Subashini#, M. P. Saravanan<sup>§</sup>, K. V. Asha #,  
M. Kananan<sup>‡</sup>, R. Vinoth Kumar#, T. Manikandan#, M. Dhivakar\*, and V. Murali#**

*\*Department of Immunology, Madurai Kamaraj University, Madurai 625 021,  
Tamil Nadu, India*

*<sup>§</sup>K. A.P.V Govt Medical College, Trichy, Tamil Nadu, India*

*<sup>‡</sup>Department of Environmental Biotechnology, Bharathidasan University, Trichy 620 024,  
Tamil Nadu, India*

*#Department of Biotechnology, Bharathidasan University, Trichy 620 024, Tamil Nadu, India*

**KEYWORDS** HLA DRB1 Alleles. South India. Castes and Tribes. Disease Associations. Phylogenetic Analysis

**ABSTRACT** In the present study 520 individuals comprising eleven different populations (castes and tribes) from the states of Tamil Nadu and Kerala, South India were genotyped for HLA – DRB1\* allele profile by PCR-SSP method. HLA DRB1\*15 (subtype of DR2) was the allele consistently showing higher frequency in all populations studied. HLA DRB1\*15 revealed a highest frequency in Kani tribe (45.19%) and the lowest frequency in Narikkuravars (Gypsies) (1.02%). The other predominant alleles based on their order of frequencies observed in each population were DRB1\*10, 07 and 15 among Iyers; DRB1\*07, 04, 15 and 08 among Kallars; DRB1\*03 and 10 among Vanniyars and Vettuva Gounders; DRB1\* 07 and 10 among Sourashtrans; DRB1\*07 and 04 among Pallars; DRB1\*04, 03, 07 and 11 among Narikkuravars; DRB1\*03 among Paliyar and Kani tribes; DRB1\*13, 10, 04, 14 among Nairs; DRB1\*10, 01, 13 and 11 among Namboothiris of Kerala. Alleles such as DRB1\*01, 08, 09, 11, 12, 14 and 16 were either present in low frequencies or completely absent in many of the south Indian populations studied. Predominantly Caucasian allele DRB1\*01 was present in higher frequencies in Namboothiris (12.85%) and Narikkuravars (8.53%) only. Allele DRB1\*01 frequency in all other populations is significantly low. However, alleles DRB1\*07 was present in many populations with higher frequencies (highest in Kallars with 23.58%). This could have been due to the higher prevalence of HIV/TB infectious and the presence of ancestral haplotype 57.1 in Indian populations. Implications of this differential distribution of these HLA-DRB1\*alleles in different castes and tribes of South India are discussed in the context of high prevalence of infectious diseases such as AIDS and TB.