

Major Histocompatibility Complex Class I-related Chain A and B Gene Expression in Sepsis Patient

Sule Menziletoglu Yildiz¹, Gulizar Atli², Ersel Gulec³, Birol Guvenc^{1,4,5} and Dilek Ozcengiz³

¹*Blood Bank, Cukurova University, Faculty of Medicine, Balcali Hospital, 01330, Saricam, Adana, Turkey*

²*Department of Biology, Cukurova University, Faculty of Science and Letters, 01330, Saricam Adana, Turkey*

³*Department of Anesthesiology, Cukurova University, Faculty of Medicine, Balcali Hospital, 01330, Saricam Adana, Turkey*

⁴*Hemapheresis, Stem Cell and Cryopreservation Unit, Cukurova University, Balcali Hospital, 01330, Saricam Adana, Turkey*

⁵*Division of Internal Medicine, Hematology Department, Cukurova University, Faculty of Medicine, Balcali Hospital, 01330, Saricam Adana, Turkey*

KEYWORDS Expression. Gene. Major Histocompatibility Complex Class I-related Chain A and B. Sepsis

ABSTRACT Major histocompatibility complex class I-related chain A and B (MICA/B) function in the regulation of protective responses due to stress induced expression. One of the major causes of the morbidity is sepsis. The objective of this study was to measure the mRNA levels of these genes in the control and sepsis patient groups. Data showed increases in mRNA concentrations of MICA and MICB in the Patient Group in contrast to the Control Group. However this level was found higher than MICA mRNA concentration in the patient group. The current study is the first report according to the researchers' knowledge representing the significant increases in the MICA and MICB expressions in the sepsis. This might provide a useful data for emphasizing these molecules as sensitive biomarkers. Further research is needed to enlighten the detail mechanisms of the MICA/B roles and to develop new treatment approaches.

INTRODUCTION

Major histocompatibility complex (MHC) class I genes encoding polymorphic peptide-binding chains mediate the intracellular antigen recognition by cytotoxic T cells. MICA functions in presentation of antigen or recognition of T cell (Bahram et al. 1994). Wang et al. (2016) indicated that MICA and MICB may broadly regulate protective responses due to stress induced expression and recognition of MICA and MICB. It was reported that MICA and MICB also increased in the patients who had chronic liver disease and hepatocellular malignancy (Chen et al. 2006; Kahraman et al. 2010).

Sepsis and also related syndromes are a major and common cause of morbidity and mortality in intensive care units (Cherian et al. 2007). The inflammatory response involving leukocytes and other inflammatory cells activation leads to oxidative stress in critical illness. Previous studies demonstrated a link between several postulated molecular mechanisms of sepsis generation and reactive oxygen species imbalance through antioxidant systems. However, there is not much research about sepsis and MICA and MICB gene expression in the related field.

This study aimed to investigate the response of MICA and MICB to evaluate the potential of these genes as sensitive biomarkers and provide fundamental data for developing the new treatment approaches in relation to sepsis.

MATERIAL AND METHODS

The current study was carried out in line with the decision of Ethic Committee of Cukurova

Address for correspondence:
Sule Menziletoglu Yildiz (PhD)
Associate Professor
Blood Bank, Cukurova University,
Faculty of Medicine, Balcali Hospital,
01330, Saricam, Adana, Turkey
Telephone: +903223386060/3027
Fax: +903223386938
E-mail: suleyildiz01@yahoo.com

University at Balcali Hospital. Individuals were separated into two groups as control and sepsis patients. All groups were examined physically and controlled for several laboratory parameters.

Blood specimens (3 mL) from patients were collected from the venous blood into EDTA containing tube. For isolation of WBCs, plasma and cells were separated by centrifugation (4000 U min⁻¹, 20°C, 5 min) immediately after blood collection. After centrifugation the supernatant above the pellet was removed and the cell pellet was treated with red blood cell lysis buffer to lyse contaminating erythrocytes. The pellet was then washed twice using PBS-Dulbecco® (Gibco, Paisley, UK) and centrifuged at each step (13000xg, 30sec). The final pellet was dissolved in 2 mL of buffer from the RNA isolation kit. RNA isolated from WBCs of the patients and controls was stored frozen at -80°C, and gene expression analysis was performed. mRNA was isolated from WBCs using the High Pure RNA Isolation Kit (Roche Diagnostics, Mannheim, Germany). Quantitative PCR was performed using the LightCycler® instrument with 4 µL of Light-Cycler Taqman master mix, 0.2 mM of oligonucleotide primers and 5 µL of cDNA solution. Amplification was performed for 10 minutes at 95 °C, followed by 40 cycles of 10 minutes at 95 °C, 30 seconds at 60 °C, and 1 seconds at 72 °C. Melting curve data were collected from 95 °C to 60 °C, at 72 °C steps. Relative gene expressions were calculated from the threshold cycles in relation to housekeeping gene, to untreated controls or healthy donors, respectively. PCR primers were as follows (Mica forward 5'-GTA TTG GGACCG GAACAC AC-3', Mica reverse 5'-ATG CTC TGG AGG GTG TGA GA-3', Micb forward 5'-TGC CAT GAA GAC CAA GAC AC-3', Micb reverse 5'-GGG GCA CTG TTC TCC TGAT-3', SDHA forward 5'-ACACAGACCTGG TGG AGA CC-3', SDHA reverse 5'-CAAAGG GCT TCT TCT GTT GC-3') The Real-Time PCR was performed using the following cycle parameters: initial enzyme activation at 95 °C for 10 min; followed by 45 cycles of 95 °C for 10 s, 60 °C for 20 s, and 65 °C for 30 s. Following the amplification phase, a cooling step was performed at 40 °C for 10 s (ramp rate of 1.5 °C/s).

Statistical Analysis

Statistical analysis of data (Mean ± Standard error) were performed by SPSS Statistics 15.0

(SPSS, Chicago, IL) package programs. Homogeneity of variance was controlled between the groups to evaluate the data distribution before the analysis. Differences between control and patient group were compared statistically by both One way ANOVA and paired Student t (P<0.05).

RESULTS

Individuals were aged <18 years and >60 years in both groups and half of individuals were female. MICA and MICB gene expressions were negative in the all individuals of control group. In the patient group MICA gene expression were only negative in 9th patient. Negative gene expressions were found in MICB in three patients (individual 7, 9 and 13). Nevertheless, positive MICA and MICB gene expressions were recorded in the other patients except these (Table 1). Therefore, three patients were eliminated during the statistical analysis for being out of the normal distribution.

Table 1: Gene expression response of control and sepsis patient groups

Individuals (N=22)	Control group		Sepsis patient group	
	Mica	Micb	Mica	Micb
1	Negative	Negative	Positive	Positive
2	Negative	Negative	Positive	Positive
3	Negative	Negative	Positive	Positive
4	Negative	Negative	Positive	Positive
5	Negative	Negative	Positive	Positive
6	Negative	Negative	Positive	Positive
7	Negative	Negative	Positive	Negative
8	Negative	Negative	Positive	Positive
9	Negative	Negative	Negative	Negative
10	Negative	Negative	Positive	Positive
11	Negative	Negative	Positive	Positive
12	Negative	Negative	Positive	Positive
13	Negative	Negative	Positive	Negative
14	Negative	Negative	Positive	Positive
15	Negative	Negative	Positive	Positive
16	Negative	Negative	Positive	Positive
17	Negative	Negative	Positive	Positive
18	Negative	Negative	Positive	Positive
19	Negative	Negative	Positive	Positive
20	Negative	Negative	Positive	Positive
21	Negative	Negative	Positive	Positive
22	Negative	Negative	Positive	Positive

The mRNA concentration of MICA was measured as $4.91 \times 10^{-7} \pm 0.55 \times 10^{-7}$ in the control group though its concentration was 0.978 ± 0.175 in the

sepsis patient group. There were observed an 2×10^6 fold increase in the patient group ($P < 0.001$) (Table 2).

MICB mRNA concentration was observed as $2.06 \times 10^{-7} \pm 0.60 \times 10^{-7}$ in the control group whereas it was $2.75 \times 10^5 \pm 0.48 \times 10^5$ in the group of sepsis patients. It was found as 1.4×10^{12} fold increase in the MICB mRNA concentrations of sepsis patient group ($P < 0.001$) (Table 2).

DISCUSSION

Septic shock caused by a Gram-negative bacteria infection results in death generally. A specific response corresponding to stress and stimulus type is induced by several factors impairing the homeostasis. Patient can cope with this by releasing various agents to eliminate or restrict the injury development. Thus, it was also emphasized that the significance of evaluation of selected systems which participate in neutralizing the stress inducers should be taken into account (Madej et al. 2007). In addition, it is also important to evaluate the evidence of relation between sepsis and stress biomarkers. On this basis, there were a lot of studies demonstrating the reactive oxygen species as relevant mediators in sepsis and septic shock in both animals and humans (Tsuiji et al. 1996; Kwiecien et al. 2016). Increased level of thiobarbituric acid reactive substances and decreased levels of antioxidants are reported indicating the lipid peroxidation resultant by oxidative stress in the plasma of patients with septic shock (De Winter et al. 2005). In addition, Kapoor et al. (2006) suggested the increased antioxidant enzyme activities (superoxide dismutase, catalase and glutathione peroxidase) in the neonates with septicemia. Authors concluded that patients defense mechanisms deteriorated against free radicals.

Although there are a lot of data showing the close relationship between oxidative stress biomarkers and sepsis, there are no studies investigating the function of MICA and MICB gene

expression in sepsis. The present study is a significant step in the direction to demonstrate the MICA and MICB gene expression increasing as a cellular stress in sepsis.

Polymorphic stress-regulated molecules Human MHC class I chain-related MICA and MICB are known as cell stress sensors. They have been characterized as the human ligands for NKG2D (Nielsen et al. 2015). MIC molecules expressed on a wide range of transformed epithelial cell surface are sensitive to immune responses mediated by NK and T cells. Despite their absence in normal adult tissue, their expressions are found in epithelial, endothelial and activated CD4⁺ and CD8⁺ T cells in the condition of stress veya stress situations (Andresen et al. 2007; Jordanova et al. 2008).

The current study demonstrated the alterations in the levels of MICA and MICB gene expression in comparison between sepsis patient group and controls. MICA and MICB are transcribed in endothelial cells, keratinocytes, fibroblasts, monocytes, gastrointestinal epithelial tissues and epithelial cell lines in humans under normal conditions. Responsive induction of MIC molecules were shown on different cell surfaces after several condition such as heat shock (Groh et al. 1998), oxidative stress (Yamamoto et al. 2001), and infections (Groh et al. 2001). Low levels of MICA expression in normal cells might be a protective mechanism to prevent the NK cell-mediated MIC-dependent cytotoxicity (Molinero et al. 2006). This was accompanied with the researchers' data which the MIC A and MIC B expressions were very low in the control group in contrast to patients with sepsis. Wang et al. (2008) recorded that IFN- γ upregulated the MIC molecule expression on the surface of monocyte in the human PBMC population. Several responses were reported including decreases and increases in MICA and MICB gene expression in the literature. For instance, Jinushi et al. (2005) studied the role of MIC A showing elevated transcripts in patients with hepatocel-

Table 2: mRNA concentrations of MIC A and MIC B in the sepsis patient and the control group. Each data point represents the mean $\pm$ standard error (n=22). Asterisks indicates the significant difference ($P < 0.001$) between control and the patient group

mRNA concentration	Control group	Sepsis patient group	Fold increase
MIC A	$4.91 \times 10^{-7} \pm 0.55 \times 10^{-7}$	$0.978 \pm 0.175^*$	2×10^6 ($P < 0.001$)
MIC B	$2.06 \times 10^{-7} \pm 0.60 \times 10^{-7}$	$2.75 \times 10^5 \pm 0.48 \times 10^5^*$	1.4×10^{12}

($P < 0.001$)

lular carcinoma. In the healthy individuals and patients who had autoimmune hepatitis, primary sclerosing cholangitis and primary biliary cirrhosis, MIC A/B expression was also examined (Holdenrieder et al. 2007). Kahraman et al. (2010) demonstrated that the induction of MIC A/B in stress conditions may function as a mechanism for the damaged, infected, or malignantly transformed cells determination. The exact mechanisms remain to be clarified by further studies. Therefore, assessment of MIC A/B serum levels could be identified as a novel index to assess. MIC A/B contribution to disease severity in patients with nonalcoholic steatohepatitis (NASH) was the first study suggesting the significant function of MIC A/B in liver injury conducted by Kahraman et al. (2010). In addition, their results has a provision for further research on the basis of investigation the mechanisms of MIC A/B expression associated with NASH, as well as their potential importance as antigen-presenting molecules for activating hepatic NK cells. This study also found the higher expression rate in MICB than MICA in the patient group which was also in accordance with the study showing higher expression levels of MICB in the patients with NASH (Kahraman et al. 2010).

It was indicated that stress (such as ischaemia-reperfusion (I/R) injury) related MIC gene expressions are observed in epithelial cells. Zou et al. (2017) also suggested a substantial increased expression in MICA in pulmonary epithelial cells resultant by scald indicating the importance of MIC A that might reflect the level of organ injury related with burns. In addition, Sconocchia et al. (2009) investigated the relationship between MIC A/B expression and innate inflammatory infiltrate in renal cell carcinoma (RCC) and they demonstrated the possible reason of expression of these molecules in most RCC cells might expose them to NK cell recognition. Furthermore, the significance of MICA expression in the regulation of macrophage/NK cell crosstalk during inflammation was also stated due to the correlation between proinflammatory cytokine secretion and the level of MICA upregulation on macrophages (Eissmann et al. 2010). Similarly, an induction of MIC expression was also observed during cytomegalovirus infection (Nielsen et al. 2015). It is concluded that the over expressed MICB gene in the patients with multiple sclerosis is an significant criterion of this depending upon the interaction between

MICB and its receptor on CD8+T or NK cells (Nielsen et al. 2015).

CONCLUSION

The significant increases observed in the expressions of both MICA and MICB in the sepsis patient group is the first report according to the researchers' knowledge. Therefore, the current study might provide useful data for emphasizing these molecules as sensitive biomarkers in sepsis. Nevertheless, there is a need for further research to enlighten the detail mechanisms of the MIC A and MIC B roles in sepsis and also to develop new treatment approaches in this area.

REFERENCES

- Andresen L, Jensen H, Pedersen MT, Hansen KA, Skov S 2007. Molecular regulation of MHC class I chain-related protein A expression after HDAC-inhibitor treatment of Jurkat T cells. *The Journal of Immunology*, 179: 8235-8242.
- Bahram S, Bresnahan M, Geraghty DE, Spies T 1994. A second lineage of mammalian major histocompatibility complex class I genes. *Proceedings of the National Academy of Sciences of the United States of America*, 91: 6259-6263.
- Chen Y, Cheng M, Tian Z 2006. Hepatitis B virus down-regulates expressions of MHC class I molecules on hepatoplastoma cell line. *Cellular & Molecular Immunology*, 3: 373-378.
- Cherian S, Jameson S, Rajarajeswari C, Helena V, Latha L, Anu Rekha MR, Nagamma T, Raju VS, Kini PG, Rao A 2007. Oxidative stress in sepsis in children. *Indian Journal of Medical Research*, 125: 143-148.
- De Winter BY, Van Nassauw L, De Man JG, De Jonge F, Bredenoord AJ, Seerden TC, Herman AG, Timmermans JP, Pelckmans PA 2005. Role of oxidative stress in the pathogenesis of septic ileus in mice. *Neurogastroenterology & Motility*, 17: 251-261.
- Eissmann P, Evans JH, Mehrabi M, Rose EL, Nedvetzki, Davis DM 2010. Multiple mechanisms downstream of TLR-4 stimulation allow expression of NKG2D ligands to facilitate Macrophage/NK Cell Crosstalk. *The Journal of Immunology*, 184: 6901-6909.
- Groh V, Rhinehartn R, Randolph-Habecker J, Topp MS, Riddell SR, Spies T 2001. Costimulation of CD8 α phbeta T cells by NKG2D via engagement by MIC induced on virus-infected cells. *Nature Immunology*, 2: 255-260.
- Holdenrieder S, Eichhorn P, Beuers U, Samtleben W, Stieber P, Nagel D, Stieber P 2007. Soluble NKG2D ligands in hepatic autoimmune diseases and in benign diseases involved in marker metabolism. *Anti-cancer Research*, 27: 2041-2045.
- Jinushi M, Takehara T, Tatsumi T, Hiramatsu N, Sakamori R, Yamaguchi S, Hayashi N 2005. Impairment of natural killer cell and dendritic cell functions by the soluble form of MHC class I-related chain A in

- advanced human hepato-cellular carcinomas. *Journal of Hepatology*, 43: 1013-1020.
- Jordanova ES, Gorter A, Ayachi O, Prins F, Durrant LG, Kenter GG, Burg SH, Fleuren GJ 2008. Human leukocyte antigen class I, MHC class I chain-related molecule A and CD8+/ regulatory T-cell ratio: Which variable determines survival of cervical cancer patients? *Clinical Cancer Research*, 14: 2028-2035.
- Kahraman A, Schlattjan M, Kocabayoglu P, Yildiz-Meziletoglu S, Schlensak M, Fingas CD, Wedemeyer I, Marquitan G, Gieseler RK, Baba HA, Gerken G, Canbay A 2010. Major histocompatibility complex class I-related chains A and B (MIC A/B): A novel role in non-alcoholic steatohepatitis. *Hepatology*, 51: 92-102.
- Kapoor K, Basu S, Das BK, Bhatia BD 2006. Lipid peroxidation and antioxidants in neonatal septicemia. *Journal of Tropical Pediatrics*, 52: 372-375.
- Kwiecien S, Magierowska K, Magierowski M, Surmiak M, Hubalewska-Mazgaj M, Pajdo R, Sliwowski Z, Chmura A, Wojcik D, Brzozowski T 2016. Role of sensory afferent nerves, lipid peroxidation and antioxidative enzymes in the carbon monoxide-induced gastroprotection against stress ulcerogenesis. *Journal of Physiology & Pharmacology*, 67(5): 717-729.
- Madej P, Plewka A, Madej J A, Nowak M, Plewka D, Franik G, Golka D 2007. Ozonotherapy in an induced septic shock. I. Effect of ozonotherapy on rat organs in evaluation of free radical reactions and selected enzymatic systems. *Inflammation*, 30: 1-2.
- Molinero LL, Domaica CI, Fuertes MB, Girart MV, Rossi LE, Zwirner NW 2006. Intracellular expression of MIC in activated CD4 + T lymphocytes and protection from NK cell-mediated MIC-dependent cytotoxicity. *Human Immunology*, 67: 170-182.
- Nielsen MM, Dyring-Andersen B, Schmidt JD, Witherden D, Lovato P, Woetmann A, Ødum N, Poulsen SS, Havran WL, Geisler C, Bonefeld CM 2015. NKG2D-dependent activation of dendritic epidermal T cells in contact hypersensitivity. *Journal of Investigative Dermatology*, 135(5): 1311-1319.
- Sconocchia G, Spagnoli GC, Del Principe D, Ferrone S, Anselmi M, Wongsena W, Cervelli V, Schultz-Thatter E, Wyler S, Carafa V, Moch H, Terracciano L, Tornillo L 2009. Defective infiltration of natural killer cells in MICA/B-positive renal cell carcinoma involves β^2 -Integrin-mediated interaction. *Neoplasia*, 11: 662-671.
- Tsuji R, Isobe N, Kawasaki H. Mechanism of prolongation of pentobarbital-induced sleeping time by empenethrin in mice 1996. *Toxicology*, 108(3): 185-190.
- Wang H, Ruan Z, Wang Y, Han J, Fu X, Zhao T, Yang D, Xu W, Yang Z, Wang L, Chen Y, Wu Y 2008. MHC class I chain-related molecules induced on monocytes by IFN-g promote NK cell activation. *Molecular Immunology*, 45: 1548-1556.
- Wang WY, Tian W, Zhu FM, Liu XX, Li LX, Wang F 2016. MICA, MICB polymorphisms and linkage disequilibrium with HLA-B in a Chinese Mongolian population. *Scandinavian Journal of Immunology*, 83(6): 456-462.
- Yamamoto K, Fujiyama Y, Andoh A, Bamba T, Okabe H 2001. Oxidative stress increases MIC and MICB gene transcription in the human colon adenocarcinoma cell line CaCo-2. *Biochimica et Biophysica Acta*, 1526: 10-12.
- Zou H, Li L, Han Y, Ma R, Liao Q, Tian J, Zhang X, Ren X, Song G, Guo Q, Li X, Ding H, Jiang G 2017. Upregulation of CD54 and downregulation of HLA ABC contribute to the novel enhancement of the susceptibility of HL-60 cells to NK cell-mediated cytotoxicity induced by ATRA plus VPA. *Oncology Reports*, 37(1): 105-114.

Paper received for publicaion on March 2016

Paper accepted for publicaion on February 2017