

Full text open access online (Since 2001)

©  Kamla-Raj IJHG 2024

PRINT: ISSN 0972-3757 ONLINE: ISSN 2456-6330

Int J Hum Genet, 24(0): 000-000 (2024)

DOI: 10.31901/24566322.2024/

MiR-155 Promotes Ischemic Stroke by Inhibiting mTOR Protein Expression

**Wenli Huang, Quanlong Hong, Huimin Wang, Shujie Gong, Zhihua Zhu
and Zhuquan Hong***

***Department of Neurology, Quanzhou First Hospital Affiliated to Fujian Medical University,
Quanzhou, Fujian, China, 362000***

KEYWORDS BV2 Cell. Ischemic Stroke. miR-155. mTOR Protein. Oxygen-Glucose Deprivation

ABSTRACT The researchers aimed to determine the serum miR-155 expression in ischemic stroke (IS) patients. Serum samples were obtained to detect miR-155 expression. National Institute of Health Stroke Scale (NIHSS) was applied to assess neurological deficits. An oxygen-glucose deprivation (OGD) model of mouse microglial BV2 cells was established. Transfection of BV2 cells with miR-155-mimic/mimic-negative control (NC) or miR-155-inhibitor/inhibitor-NC was conducted. Markedly elevated miR-155 expression in serum of IS patients was detected in comparison with that in healthy individuals ($P<0.001$). NIHSS score declined progressively on 1st, 7th and 14th days after onset ($P<0.005$). The survival rate of BV2 cells after OGD treatment decreased evidently ($P<0.001$). Moreover, miR-155 expression was significantly reduced, but the protein expression of mTOR increased in OGD group ($P<0.05$). The downregulation of miR-155 facilitated mTOR mRNA and protein expressions ($P<0.005$). Elevated miR-155 is associated with the pathogenesis of IS and negatively correlated with prognosis. MiR-155 suppresses mTOR protein expression in mouse microglial BV2 cells and facilitates OGD-induced stress response.