

Full text open access online (Since 2001)

©  Kamla-Raj IJHG 2024

PRINT: ISSN 0972-3757 ONLINE: ISSN 2456-6330

Int J Hum Genet, 24(0): 000-000 (2024)

DOI: 10.31901/24566322.2024/

Expression Analysis and Bioinformatics-Based Identification of miR-4534, miR-551a and miR-1294 in Breast Cancer

**Bodour Alassil*, Filiz Özbas Gerçeker*, Alper Aytekin#, Nazli Bozman*,
M. Avni Gökcalp#, Latif Yilmaz# and Havva Yesil^a**

**Department of Biology, Faculty of Arts and Science, University of Gaziantep,
Gaziantep, Türkiye*

Telephone: +905363443656, E-mail: b85_assil@hotmail.com

*# Department of General Surgery, School of Medicine, Gaziantep University,
Gaziantep, Türkiye*

*^a Department of Internal Medicine, School of Medicine, Gaziantep University,
Gaziantep, Türkiye*

KEYWORDS Wnt Pathway. Hippo Pathway. Gene Regulation. Target Genes. MicroRNAs. P53 Pathway

ABSTRACT This study aimed to ascertain the potential role of specific microRNAs (miR-551a, miR-1238, miR-4534, and miR-1294) in breast cancer by analysing their expression levels and identifying the target genes and pathways they regulate. Expression levels in 82 samples from 41 patients were measured using quantitative Real-Time PCR. Bioinformatics tools were utilised to identify the potential target genes and pathways. Findings revealed that miR-1294 and miR-551a had significantly lower expression in tumour tissues compared to normal tissues, while miR-4534 was overexpressed in tumour tissues. No significant difference was observed in miR-1238 expression between the two. miR-1294, miR-551a, and miR-4534 were found to regulate genes associated with Mapk, ErbB, Hippo Wnt, EGFR, and P53 signalling pathways. The study concludes that miR-551a, miR-1294, and miR-4534 may play a role in the development of breast cancer.