

## Plant Diversity in the Homegardens and their Significance in the Livelihoods of War Khasi Community of Meghalaya, North-east India

H. Tynsong\* and B. K. Tiwari\*\*

\*Regional Centre, National Afforestation and Eco-Development Board, \*\*Department of Environmental Studies, North-Eastern Hill University, Shillong 793 022, Meghalaya, India

**KEYWORDS** Cash Income. Medicinal Plants. Conservation. Management Zones. Marketing

**ABSTRACT** The study was conducted with an objective of investigating the plant species diversity in homegardens and their contribution towards livelihoods of War Khasi community of Meghalaya. In total, 197 plant species were recorded with an average of 89 plant species per homegarden. The average size of homegarden was 750 m<sup>2</sup> with the annual gross production of Rs.3514 per homegarden; this contributes about 7 % towards average annual gross income per household. Computing in terms of per unit area, the average annual gross income was found to be Rs.73,748/ha. The study revealed that 35% of the produce was used for self consumption and 65% was sold in the local markets. The important plant species which contributed to the household income were: *Piper betle* (24%), *Mangifera indica* (19%), *Litchi chinensis* (15%) and *Areca catechu* (11%). We conclude that although homegardens contribute only a small part of total income, they are particularly important because of low labour input for management and locational advantage.

### INTRODUCTION

The cultivation of fruits, vegetables and ornamentals in homegardens has a long tradition in north-east India, especially among the hill tribes residing in the states of Assam, Manipur, Nagaland and Meghalaya. In general, homegardens are characterized by different vegetation strata composed of trees, shrubs and herbs in association with annual and perennial agricultural crops and small livestock within the house compounds. Homegarden may be defined as, 'a bounded piece of land cultivated with a diverse mixture of annual and perennial crops on which a house is built' (Karyono 1990). The major functions of homegardens particularly in rural areas are subsistence production and income generation (Kumar and Nair 2004). Because of high plant diversity in the homegardens, a wide spectrum of multiple-use plant products can be generated with relatively low labour, cash or other inputs. In seasons of scarcity homegardens with their diverse products available year round, contribute towards food security. They also fulfil many social, cultural and ecological needs. The multi-layered, forest-like vegetation structure of homegardens contributes substantially to the ecological sustainability of the village ecosystems (Kehlenbeck and Maass 2004). This vegetation structure prevents soil erosion, provides habitat

to wild plants and soil microorganisms and promotes a favourable micro-climate for the household. Because of their richness in plant species, homegardens are regarded as an ideal production system for *in situ* conservation of plant species. The species composition of homegardens varies according to climatic and edaphic factors of the place and socio-economic condition of the people.

Homegarden systems provide an additional food supply and cash income for the people (Das and Das 2005). It is reported that in Indonesia and Nicaragua, homegardens contribute 21.1% and 35% of the total income respectively. Studies from south-west Bangladesh (Motiur et al. 2006) and north-eastern Bangladesh (Motiur et al. 2005) reported that on an average 15.9% and 11.8% of household income is derived from homegardens respectively. Many homegarden systems around the world have been studied for plant diversity and for their economic value (Motiur et al. 2005). However, researches on homegardens of India are very few (Kingston et al. 2006). The review of literatures clearly showed that there is a lack of research on plant diversity and economic value of homegardens in the north-eastern India where tradition of homegardens is very old (Das and Das 2005). This study aims at analyzing plant species richness in these homegardens and investigates the

contribution of the fruits, vegetables and spices grown in the homegardens towards cash income of the people.

### STUDY AREA

The study was conducted in south Meghalaya, India (25° 7'–25° 18' N latitude and 91°–92° E longitude). Five surveyed villages were: Nongkwai (25° 20' N, 91° 54' E); Mawriang (25° 12' N 91° 85' E); Umkrem (25° 14' N, 91° 55' E'); Nolikata (25° 15' N, 91° 13' E) and Ranikor (25° 15' N, 91° 11' E) (Fig. 1). The altitude of the study area ranged from 450 m to 700 m above mean sea level. The mean annual maximum and minimum temperature was 23°C and 13°C respectively. The population is sparse and the people are predominantly employed in plantations and forestry related activities. Agriculture is limited to some small valleys where mainly tuber crops are grown. Arecanut, orange, betel leaf, jack fruit, bayleaf, honey and broom grass are the principal produce of the region. The region is inhabited by *War Khasi* people, a tribal community having long tradition of forest conservation (Tiwari 2005). The size of homegardens varies between 200 m<sup>2</sup> to 3500 m<sup>2</sup>. The average size was about 750 m<sup>2</sup>. The total number of household in the study area varied from 59 at Nolikata to 200 at Nongkwai. The average household con-

sists of 8 members. The natural vegetation of the study area ranges from tropical evergreen to subtropical evergreen forests (Balakrishnan 1983).

### METHODS

#### Field Sampling

Four complementary approaches were adopted, namely; (a) formal interview with the village headmen and secretary, (b) direct observation, (c) interaction with the head of the selected households through questionnaires and (d) phytosociological study in sample homegardens. The interview schedule comprised of a general introduction concerning management of homegardens, farming practices, domesticated and wild plants and their products found in the homegardens and the final section of discussion related to various socio-economic characteristics of the household. The discussion and interviews were conducted following the methods described by High and Shakleton (2000). The survey was administered to a random sample of 30 households in each village. Homegardens size was measured, excluding the area occupied by the house. In case of tree, shrub, climber and epiphytic plant species whole homegarden was used as sample plot while in the case of herb species a 1 x 1 m quadrat was used.




Fig. 1. Location of the study area

### Plant Survey and Analysis

For tree, shrub, climber and epiphytes complete inventories were carried in 150 homegardens; where as in the case of herbs, 1 m<sup>2</sup> plots (1x1 m) for each homegardens was studied to assess plant diversity. The tree species diameter at breast height (dbh  $\geq$  5 cm) were individually measured and numbered and their density and frequency per plot were computed. The information on the uses and importance of plant species were collected by using questionnaires, focused group discussion, key informant interview, and Participatory Rural Appraisal (PRA) methods as described by Mukherjee (1993). The data and information about management of homegardens, plants, income and population were also collected using questionnaires and through discussions and interviews with elders of the sample household. Shannon and Wiener Index of diversity (Shannon and Wiener 1963) was calculated to analyze the diversity of homegardens for each village studied using the formula  $H' = -\sum p_i \ln p_i$ , where,  $p_i$  = proportion number of  $i$ th species. The frequency, density and Importance Value Index (IVI) of the species were determined following the methods of Misra (1968) and Mueller-Dombois and Ellengberg (1974). The basal cover (m<sup>2</sup>ha<sup>-1</sup>) was calculated by using the formula: Basal Cover (BC) = density x average tree basal area (Khan et al. 1997). The species richness is the total number of all the species. Plant specimens were collected and identified with the help of Flora of Assam (Kanjilal et al. 1934-1940) and Flora of Jowai (Balakrishnan 1981-1983). The identifications were confirmed by consulting the herbaria of Botanical Survey of India, North-eastern Circle, Shillong. The nomenclatures of the species are as per the regional flora.

### Income

Household income from homegarden was calculated by asking the respondents the amount of homegarden products they sold and consumed how much income was earned from the previous year's sale of homegarden products and summing the sales and consumption values, allowed us to calculate the proportion of total income earned from the homegardens. Gross annual income of sampled households from agriculture, off-farm and other sources of income was also collected

in order to compute the contribution of the homegardens towards peoples' gross income. Gross income was calculated by adding the amount of money earned from all the products collected from homegardens including those used for self consumption and sale. We converted other currencies at 1 US\$ = 42.7 Indian Rupees, 1US\$ = 7.47 South African Rands and 1US\$ = 107.60 Sri Lanka Rupees as on July, 2008.

### Computation of Expenditures in Gardening

The expenditures in maintaining of the homegardens and marketing of plant products included management cost, sorting cost, harvesting cost, marketing cost, plantation cost, planting material and transport cost. These costs were calculated by multiplying wage rate by number of days spent. The plant products collected from homegardens which require sorting and processing before selling included betel leaf, betel nut and bayleaf. The money spent in arranging of betel leaf, processing of betel nut and packing of bayleaf was included in sorting cost. Sorting and arranging of the products was done by women and packing of bayleaf was done by men. Harvesting of plant products from homegardens was mostly done by men except for few species in which young boys and women were also involved. Harvesting of areca nut, bayleaf and betel leaf was done by skilled men. In the study area, marketing of forest product was exclusively the duty of women. Marketing cost was calculated by multiplying the period of time spent in selling the products by the daily wage of women. Majority of plants in the homegardens are cultivated by men. The plantation cost was calculated by multiplying the time spent in plantation of each species by daily wage of the cultivator. This was added to the amount of money spent in buying the seed or the propagules to be planted in the homegardens. The plant species in which the seed/propagules are to be purchased included black pepper @Rs.5/sapling, betel nut @Rs.5/sapling, litchi @Rs.10/sapling and maize @Rs.5/kg. In case of other plant species, people used their own seed saved or stored by them. Transport cost was the total amount of money spent in transporting the plant products from village to the market (by head load/vehicle/boat). In the study area, ordinarily the daily wage rate of workers during the study period was Rs.150 for skilled men, Rs.100 for unskilled men, Rs.45 for women

and Rs.30 for boys aged between 12-15 yr. The cost of materials was calculated from market rate.

## RESULTS

### Management of Homegardens

The management of homegardens includes fencing, weeding and tree lopping. In the fencing of gardens, *Bambusa vulgaris* is used as pole and *B. tulda* as spokes. Villagers use seeds, seedlings and vegetative propagules to regenerate homegarden plants. Soil fertility of homegardens is maintained naturally from leaf litter, faeces of reared animals and kitchen waste manure. In the day to day maintenance of homegardens, both men and women are involved.

### Plant Diversity

We recorded a total of 197 plant species (70 trees, 41 shrubs, 50 herbs, 23 climbers and 13 epiphytes) belonging to 77 families comprising of 73 angiosperm, 1 gymnosperm and 3 pteridophytes. Moraceae was the dominant family with 20 species followed by Orchidaceae 12 species, Asteraceae 10 and Euphorbiaceae 9 species. The homegardens showed four distinct strata. Strata A or canopy layer was composed of big tree (>15 m height), sub-canopy or B strata was composed of middle sized tree (8 m to 15 m height), under-canopy or C strata was composed of small trees (<8 m height) and ground vegetation ( $\leq$  2 m height) included shrub and herbs. The common tree species in the A strata were: *Artocarpus heterophyllus*, *Mangifera indica*, *Bombax ceiba* and *Duabanga grandiflora*. B strata was composed of *Areca catechu*, *Artocarpus lakoocha*, *Cinnamomum tamala* and *Baccaurea sapida* and C strata composed of *Averrhoa carambola*, *Psidium guajava* and *Zizyphus mauritiana*. The shrub layers were dominated by *Clerodendron colebrookianum*, *Hibiscus rosa-sinensis*, *Hibiscus sabdariffa*, *Manihot esculenta* and *Citrus limon*. The dominant plants of herb layers were: *Ageratum conyzoides*, *Houttuynia cordata*, *Drymaria cordata*, *Oplismenus compositus* and *Borreria pilosa*. The species name, frequency, density and IVI of ten most frequent trees, shrubs, herbs, climbers and epiphytes encountered during the study are given in table 1.

The homegardens appear to be assorted mixture of tree, shrub, herb, climber and epiphytes

mainly ornamental orchids. However, the locations of most plants were found to be deliberate which could be distinguished into several management zones. The present study recorded six major management zones based on their purpose, location and species composition viz., (a) arecanut (*Areca catechu*) grove, (b) banana grove, (c) vegetable garden, (d) fruit grove, (e) ornamental and (f) mixture of different plant species (Fig. 2). These zones fulfil specific objectives for household ranging from food, medicinal, cash, timber, fuelwood etc. Arecanut and fruit zones were the most important zones because of economic importance of the plants grown in these zones.

### Expenditure and Cash income

The total amount of money spent by the people living in the five villages of study area in management of homegardens and other expenditures related to collection and processing of plant products from their homegardens was found to be Rs.8,66,438 (Table 3).

The gross production of homegardens of the five villages of south Meghalaya was worth Rs.19,69,671 per annum out of which Rs.6,88,442 (34.95%) was used for self-consumption, and Rs.12,81,229 (65.04%) was sold in local markets. The most important plant species found in the homegardens which contributed to the household income were: *Piper betle* (24.15%), *Mangifera indica* (19.30%), *Litchi chinensis* (15.16%) and *Areca catechu* (10.8%) (Table 4). The average annual gross income from homegardens was Rs.3,514.58 per household which accounted to Rs.73,748.39 per hectare (Table 5). Homegardens contributed 7% of the total household income.

## DISCUSSION

### Plant Diversity

Majority of the plant species recorded in the homegardens of south Meghalaya were typical plants found in homegardens throughout the tropics e.g., banana (sweet and plantains), coconut, jack fruit, guava, mango, papaya, citrus, the tuber, yam, cassava, taro, chilli, sugarcane etc. (Mendez et al. 2001; Nautiyal et al. 2008). These plants provide a broad basis for self-sufficiency in food for households. The total number of plant


**Table 1: The frequency (%), density (plant ha<sup>-1</sup>) and IVI of 10 most frequent tree, shrub, herb, climber and epiphyte species encountered in the homegardens of south Meghalaya**

| Plant species                                     | Nongkwai |      |       | Mawriang |      |       | Umikrem |      |       | Nolikata |      |       | Ranikor |      |       |
|---------------------------------------------------|----------|------|-------|----------|------|-------|---------|------|-------|----------|------|-------|---------|------|-------|
|                                                   | F        | D    | IVI   | F        | D    | IVI   | F       | D    | IVI   | F        | D    | IVI   | F       | D    | IVI   |
| Tree                                              |          |      |       |          |      |       |         |      |       |          |      |       |         |      |       |
| <i>Areca catechu</i> L.                           | 33.33    | 41   | 13.89 | 27.00    | 155  | 26.64 | 90.00   | 189  | 60.60 | 63.33    | 287  | 53.28 | 90.00   | 280  | 51.46 |
| <i>Artocarpus heterophyllus</i> Ham.              | 50.00    | 32   | 40.00 | 24.00    | 87   | 19.71 | 76.67   | 96   | 29.13 | 53.33    | 69   | 70.18 | 46.67   | 53   | 30.87 |
| <i>Baccaurea sapida</i> (Roxb.) Müll. Arg.        | 13.33    | 9    | 3.98  | 4.00     | 9    | 3.28  | 6.00    | 12   | 11.53 | 6.67     | 5    | 2.06  | 3.33    | 2    | 0.81  |
| <i>Carica papaya</i> L.                           | 20.00    | 15   | 6.06  | 3.00     | 11   | 4.87  | 10.00   | 13   | 3.41  | 20.00    | 31   | 7.81  | 23.33   | 30   | 7.75  |
| <i>Citrus maxima</i> S. Dutta & S.C.Bhattach.     | 56.67    | 45   | 19.88 | 16.00    | 55   | 10.24 | 53.33   | 65   | 18.69 | 10.00    | 13   | 4.06  | 10.00   | 7    | 2.58  |
| <i>Litsea chinensis</i> Sonn.                     | 20.00    | 19   | 10.87 | 22.00    | 94   | 16.10 | 73.33   | 111  | 36.56 | 10.00    | 5    | 2.82  | 3.33    | 1    | 0.69  |
| <i>Mangifera indica</i> L.                        | 63.33    | 60   | 39.39 | 25.00    | 96   | 17.50 | 83.33   | 113  | 60.10 | 40.00    | 34   | 19.64 | 63.33   | 59   | 23.00 |
| <i>Musa</i> spp.                                  | 33.33    | 51   | 16.69 | 10.00    | 57   | 12.57 | 23.33   | 30   | 10.30 | 26.67    | 118  | 23.01 | 20.00   | 54   | 10.42 |
| <i>Prunus persica</i> (L.) Batsch.                | 23.33    | 24   | 7.93  | 3.33     | 4    | 2.66  | 3.33    | 4    | 1.08  | 16.00    | 3    | 0.62  | 3.33    | 2    | 0.80  |
| <i>Psidium guajava</i> L.                         | 56.67    | 63   | 20.52 | 40.00    | 34   | 19.64 | 16.67   | 26   | 7.01  | 36.67    | 27   | 10.96 | 16.67   | 17   | 4.77  |
| Shrub                                             |          |      |       |          |      |       |         |      |       |          |      |       |         |      |       |
| <i>Allamanda cathartica</i> L.                    | 10.00    | 3    | 4.61  | 26.67    | 73   | 13.47 | 30.00   | 86   | 23.09 | 3.33     | 4    | 5.04  | 3.33    | 8    | 1.54  |
| <i>Capsicum frutescens</i> L.                     | 13.33    | 28   | 11.78 | 13.33    | 61   | 9.33  | 13.33   | 72   | 23.09 | 13.33    | 10   | 17.88 | 10.00   | 4    | 12.56 |
| <i>Citrus assamensis</i> S. Dutta & S.C.Bhattach. | 13.33    | 22   | 10.24 | 20.00    | 31   | 7.56  | 13.33   | 26   | 8.31  | 3.33     | 3    | 1.06  | 3.33    | 4    | 1.66  |
| <i>Citrus karna</i> Raf.                          | 20.00    | 22   | 12.81 | 3.33     | 2    | 0.95  | 3.33    | 3    | 1.45  | 3.33     | 4    | 5.04  | 3.33    | 4    | 5.67  |
| <i>Citrus limon</i> Burm.                         | 13.33    | 22   | 10.24 | 26.67    | 28   | 8.63  | 26.67   | 33   | 13.48 | 10.00    | 9    | 13.98 | 16.67   | 10   | 23.16 |
| <i>Clerodendron colebrookianum</i> Walp.          | 36.67    | 45   | 24.84 | 33.33    | 34   | 10.75 | 33.33   | 40   | 16.80 | 3.33     | 4    | 5.67  | 20.00   | 31   | 7.56  |
| <i>Hibiscus rosa-sinensis</i> L.                  | 6.67     | 5    | 2.90  | 26.67    | 31   | 8.98  | 26.67   | 36   | 14.11 | 6.67     | 20   | 15.18 | 10.00   | 28   | 26.64 |
| <i>Hibiscus sabdariffa</i> L.                     | 3.33     | 3    | 2.05  | 3.33     | 3    | 1.06  | 6.67    | 5    | 2.90  | 3.33     | 14   | 9.58  | 13.33   | 5    | 4.96  |
| <i>Manihot esculenta</i> Crantz.                  | 6.67     | 2    | 3.08  | 43.33    | 108  | 20.79 | 40.00   | 127  | 32.88 | 16.67    | 36   | 32.57 | 16.67   | 46   | 44.64 |
| <i>Plogocanthus parviflorus</i> T. Anders.        | 6.67     | 8    | 4.35  | 3.33     | 4    | 2.31  | 3.33    | 4    | 1.18  | 3.33     | 3    | 1.45  | 16.67   | 98   | 14.06 |
| Herb                                              |          |      |       |          |      |       |         |      |       |          |      |       |         |      |       |
| <i>Ageratum conyzoides</i> L.                     | 30.00    | 1621 | 16.31 | 43.33    | 1898 | 21.60 | 43.33   | 1798 | 26.80 | 63.33    | 3053 | 42.04 | 60.00   | 3164 | 45.46 |
| <i>Bidens pilosa</i> (Blume.) Sherff.             | 10.00    | 522  | 5.34  | 10.00    | 522  | 5.48  | 16.67   | 733  | 10.63 | 10.00    | 133  | 3.74  | 10.00   | 244  | 5.10  |
| <i>Canna</i> sp.                                  | 6.67     | 167  | 2.57  | 23.33    | 455  | 8.26  | 23.33   | 455  | 10.38 | 20.00    | 511  | 9.51  | 6.67    | 178  | 3.53  |
| <i>Centella asiatica</i> (L.) Urban.              | 13.33    | 977  | 8.64  | 20.00    | 855  | 9.84  | 20.00   | 855  | 12.56 | 3.33     | 22   | 0.93  | 6.67    | 266  | 4.31  |
| <i>Drymaria cordata</i> (L.) Roem. & Schult.      | 6.67     | 100  | 2.21  | 3.33     | 44   | 1.06  | 3.33    | 44   | 1.32  | 6.67     | 189  | 3.32  | 3.33    | 111  | 1.96  |

Table 1: Contd..

| Plant species                                  | Nongkwai |      |       | Mawriang |      |       | Umikrem |      |       | Nolikata |     |        | Ranikor |     |       |
|------------------------------------------------|----------|------|-------|----------|------|-------|---------|------|-------|----------|-----|--------|---------|-----|-------|
|                                                | F        | D    | IVI   | F        | D    | IVI   | F       | D    | IVI   | F        | D   | IVI    | F       | D   | IVI   |
| <i>Houttuynia cordata</i> Thunb.               | 56.67    | 3541 | 33.42 | 56.67    | 3541 | 34.54 | 20.00   | 1099 | 14.49 | 13.33    | 278 | 5.82   | 6.67    | 178 | 3.53  |
| <i>Oplismenus compositus</i> P. Beauv.         | 26.67    | 1077 | 12.52 | 26.67    | 877  | 11.56 | 33.33   | 1832 | 24.15 | 3.33     | 100 | 1.38   | 16.67   | 722 | 11.25 |
| <i>Oxalis corniculata</i> L.                   | 20.00    | 733  | 8.98  | 3.33     | 233  | 2.18  | 3.33    | 233  | 2.81  | 6.67     | 144 | 2.44   | 13.33   | 488 | 8.23  |
| <i>Sansevieria trifasciata</i> Hort. ex Prain. | 6.67     | 300  | 3.30  | 6.67     | 366  | 3.76  | 6.67    | 366  | 4.83  | 20.00    | 311 | 6.61   | 3.33    | 122 | 2.06  |
| <i>Tagetes erecta</i> L.                       | 3.33     | 44   | 1.07  | 3.33     | 44   | 1.06  | 3.33    | 44   | 1.06  | 13.33    | 311 | 6.09   | 23.33   | 733 | 13.33 |
| Climber                                        |          |      |       |          |      |       |         |      |       |          |     |        |         |     |       |
| <i>Dioscorea hispida</i> Dennst.               | 13.33    | 4    | 6.22  | 13.33    | 13   | 7.13  | 16.67   | 16   | 9.26  | 36.67    | 54  | 24.76  | 3.33    | 4   | 7.58  |
| <i>Dolichos lablab</i> L.                      | 20.00    | 37   | 14.84 | 36.67    | 55   | 23.04 | 3.33    | 4    | 5.59  | 6.67     | 4   | 12.13  | 3.33    | 4   | 5.59  |
| <i>Elaeagnus latifolia</i> L.                  | 20.00    | 12   | 10.86 | 16.67    | 11   | 7.90  | 16.67   | 13   | 8.87  | 6.67     | 6   | 14.15  | 10.00   | 4   | 13.59 |
| <i>Hemidesmus indicus</i> (L.) R.Br.           | 20.00    | 33   | 13.98 | 20.00    | 34   | 13.32 | 3.33    | 4    | 1.97  | 10.00    | 7   | 4.74   | 6.67    | 11  | 12.77 |
| <i>Hodgsonia heteroclita</i> Kh.f.&T.          | 10.00    | 13   | 6.64  | 3.33     | 2    | 1.58  | 3.33    | 3    | 1.77  | 16.67    | 51  | 15.18  | 16.67   | 27  | 10.99 |
| <i>Merremia hederacea</i> (Burn.f.) Hallierf.  | 6.67     | 9    | 4.73  | 10.00    | 7    | 4.74  | 10.00   | 16   | 26.77 | 3.33     | 4   | 5.59   | 6.67    | 6   | 3.85  |
| <i>Mikania micrantha</i> Kunth.                | 16.67    | 49   | 16.4  | 16.67    | 51   | 15.18 | 16.67   | 27   | 10.99 | 10.00    | 28  | 35.86  | 30.00   | 133 | 92.10 |
| <i>Passiflora edulis</i> Sim.                  | 6.67     | 10   | 4.81  | 6.67     | 10   | 4.17  | 16.67   | 24   | 10.33 | 3.33     | 3   | 6.57   | 43.33   | 29  | 42.14 |
| <i>Piper betle</i> L.                          | 40.00    | 97   | 33.77 | 70.00    | 200  | 61.48 | 70.00   | 237  | 64.23 | 16.67    | 30  | 46.98  | 10.00   | 19  | 19.94 |
| <i>Piper nigrum</i> L.                         | 10.00    | 28   | 76    | 30.00    | 88   | 26.75 | 46.67   | 166  | 44.11 | 13.33    | 33  | 62.15  | 10.00   | 46  | 31.58 |
| Epiphyte                                       |          |      |       |          |      |       |         |      |       |          |     |        |         |     |       |
| <i>Aerides multiflorum</i> Roxb.               | 10.00    | 13   | 46.73 | -        | -    | -     | 6.67    | 16   | 31.79 | -        | -   | -      | -       | -   | -     |
| <i>Aerides odoratum</i> Lour.                  | -        | -    | -     | -        | -    | -     | 10.00   | 12   | 35.65 | -        | -   | -      | -       | -   | -     |
| <i>Coelogyne corymbosa</i> Lindl.              | -        | -    | -     | -        | -    | -     | -       | -    | -     | -        | -   | -      | 10.00   | 24  | 78.54 |
| <i>Cymbidium elegans</i> Lindl.                | -        | -    | -     | -        | -    | -     | -       | -    | -     | -        | -   | -      | 10.00   | 14  | 59.42 |
| <i>Cymbidium mastersii</i> Griff. Ex Lindl.    | -        | -    | -     | 10.00    | 28   | 76    | 13.33   | 33   | 62.15 | -        | -   | -      | -       | -   | -     |
| <i>Dendrobium chrysanthum</i> Wall. Ex Lindl.  | -        | -    | -     | -        | -    | -     | -       | -    | -     | -        | -   | -      | 3.33    | 4   | 17.15 |
| <i>Dendrobium longicornu</i> Lindl.            | -        | -    | -     | -        | -    | -     | 3.33    | 21   | 28.32 | 3.33     | -   | -      | -       | -   | -     |
| <i>Dendrobium ochreatum</i> Wall. Ex Lindl.    | 6.67     | 8    | 29.92 | 3.33     | 4    | 44.76 | -       | -    | -     | -        | 5   | 94.7   | -       | -   | -     |
| <i>Eria paniculata</i> Lindl.                  | 10.00    | 12   | 44.91 | -        | -    | -     | -       | -    | -     | -        | -   | -      | -       | -   | -     |
| <i>Vanda cristata</i> Wall. Ex Lindl.          | 10.00    | 14   | 48.52 | -        | -    | -     | -       | -    | -     | -        | 6   | 105.31 | -       | -   | -     |

F-frequency, D-density, IVI-important value index



**Fig. 2. Different management zones: a-arecanut grove, b-banana grove, c-vegetable garden, d-fruit grove, e-ornamental and f-mixture of different plant species**  
(Illustration: H.Tynsong, drawn at Umkrem village on 18<sup>th</sup> Dec. 2009)

**Table 2: Diversity and community characteristics of homegardens at Nongkwai, Mawriang, Umkrem, Nolikata and Ranikor villages of south Meghalaya**

| Plant component | Village  | F  | G  | S  | D     | BC    | PE   | SD   | SI   |
|-----------------|----------|----|----|----|-------|-------|------|------|------|
| Tree            | Nongkwai | 27 | 31 | 41 | 588   | 12.96 | 0.56 | 3.14 | 0.06 |
|                 | Mawriang | 26 | 36 | 46 | 867   | 8.15  | 0.88 | 4.01 | 0.06 |
|                 | Umkrem   | 17 | 24 | 28 | 823   | 31.02 | 0.74 | 2.51 | 0.12 |
|                 | Nolikata | 20 | 28 | 31 | 852   | 45.01 | 0.78 | 2.70 | 0.11 |
|                 | Ranikor  | 25 | 34 | 41 | 861   | 39.80 | 0.75 | 2.81 | 0.10 |
| Shrub           | Nongkwai | 20 | 27 | 32 | 420   | -     | 0.89 | 3.08 | 0.06 |
|                 | Mawriang | 22 | 31 | 36 | 932   | -     | 0.86 | 3.11 | 0.06 |
|                 | Umkrem   | 11 | 16 | 22 | 621   | -     | 0.85 | 2.89 | 0.08 |
|                 | Nolikata | 10 | 13 | 15 | 229   | -     | 0.95 | 2.52 | 0.09 |
|                 | Ranikor  | 8  | 10 | 12 | 169   | -     | 0.66 | 1.66 | 0.13 |
| Herb            | Nongkwai | 20 | 25 | 32 | 18393 | -     | 0.87 | 3.03 | 0.07 |
|                 | Mawriang | 23 | 33 | 34 | 16828 | -     | 0.85 | 3.00 | 0.07 |
|                 | Umkrem   | 22 | 28 | 29 | 12687 | -     | 0.86 | 2.89 | 0.08 |
|                 | Nolikata | 14 | 19 | 23 | 12032 | -     | 0.95 | 2.52 | 0.09 |
|                 | Ranikor  | 22 | 27 | 28 | 11444 | -     | 0.86 | 2.84 | 0.09 |
| Climber         | Nongkwai | 8  | 8  | 20 | 578   | -     | 0.85 | 2.49 | 0.11 |
|                 | Mawriang | 10 | 16 | 18 | 545   | -     | 0.80 | 2.34 | 0.15 |
|                 | Umkrem   | 1  | 7  | 10 | 512   | -     | 0.78 | 1.49 | 0.16 |
|                 | Nolikata | 8  | 10 | 11 | 124   | -     | 0.91 | 2.19 | 0.13 |
|                 | Ranikor  | 8  | 8  | 10 | 236   | -     | 0.76 | 1.76 | 0.26 |
| Epiphyte        | Nongkwai | 1  | 5  | 5  | 55    | -     | 0.83 | 1.59 | 0.21 |
|                 | Mawriang | 2  | 3  | 4  | 56    | -     | 1.15 | 1.27 | 0.30 |
|                 | Umkrem   | 2  | 3  | 5  | 100   | -     | 0.96 | 1.56 | 0.22 |
|                 | Nolikata | 2  | 2  | 2  | 11    | -     | 0.99 | 0.69 | 0.50 |
|                 | Ranikor  | 2  | 4  | 4  | 54    | -     | 0.92 | 1.28 | 0.30 |

F-families, G-genera, S-species, D-density ( $\text{ha}^{-1}$ ), BC-basal cover ( $\text{m}^2\text{ha}^{-1}$ ), PE- Pielou's evenness index, SD- Shannon's diversity index, SI- Simpson's dominance index, '-' indicates absent

species as well as the average number of plant species per homegarden in south Meghalaya was rather high and comparable to plant diversity of homegardens in Assam, India (Das and Das 2005) but lower than the number of plant species reported in Nicaragua (Mendez et al. 2001) and South Africa (High and Shakleton 2000). The average number of plant species per homegarden was very high (89 plant species) and much higher than the number of plant species found in Cuba (22 plant species). The mean Shannon indices of plant species (2.37) was also considerably higher as compared to Cuba (1.79) (Wezel and Bender 2003) and central Sulawesi, Indonesia (2.32) (Kehlenbeck and Maass 2004) but comparatively much lesser than the mean Shannon indices of arecanut agroforests (3.04) of south Meghalaya (Tynsong 2009). Mean Shannon indices vary widely in tropical homegardens and are reported to range from 0.93 to 3.00 (Karyono 1990). The high plant species richness in the homegardens of south Meghalaya may be due to its location in the lower altitude as also observed elsewhere (Quiroz et al. 2002).

The homegardens of all the five villages possessed a multi-layered vegetation structure which could offer advantages in controlling of soil erosion and efficient use of resources viz., light and space. Hohegger (1998) also observed similar multi-layered homegardens in Sri Lanka, which are very old and larger in size. Presence of large number of medicinal plants in the homegardens clearly showed that medicinal plants still play an important role in healing of people's ailments. Medicinal plants were found in larger numbers in the homegardens of herbal practitioners. Most medicinal plants were collected from the wild and domesticated in the homegardens. These included: *Calamus floribundus*, *Centella asiatica*, *Fagopyrum cymosum*, *Hemidesmus indicus*,

*Kaempferia rotunda*, *Mussaenda roxburghii*, *Merremia hedereaeca*, *Piper peepuloides*, *Smilax* sp. and *Trichosanthes palmata*. This corroborates with the finding of Kingston et al. (2006) and Das and Das (2005) who also reported that homegardens preserve large number of plant species having medicinal properties. During interviews it was noted that herbal practitioners have a sound knowledge of medicinal plant uses and techniques to grow them in their homegardens. For herbal practitioners, majority of plant species found in the homegardens such as fruit trees, tubers, vegetables, spices etc., have potential medicinal use. Large numbers of ornamental plants are also grown in the homegardens. Some of them were collected from the wild and domesticated in the homegardens viz., *Cassia tora*, *Dracaena fragrans*, *Luculia pinceana*, *Morinda angustifolia*, *Raphidophora calophyllum* and a variety of orchids. The study also revealed that War Khasi community were concerned in the cleaning, beautifying and maintaining the surroundings of their settlements.

The homegardens contribute a great deal to food supply especially for the people living in the rural areas because of high production and diversity of cultivated edible species. Although the extent of household dependency on homegardens varies considerably, its contribution is quite significant towards livelihood of the people because of low investment and easy accessibility. Homegardens function as *ex situ* as well as *in situ* conservation plots for plant genetic resources of the region. In the villages studied, 103 wild plant species were recorded from the homegardens which showed that homegardens are also a home to many wild plant species, thus they serve as a repository of wild plants.

**Table 3: Annual expenditure (Rs.) incurred in management of homegardens and marketing of the products collected from homegardens of five villages of south Meghalaya**

| Activities                                | Nongkwai | Mawriang | Umkrem | Nolikata | Ranikor | Total    |
|-------------------------------------------|----------|----------|--------|----------|---------|----------|
| Management cost                           | 184333   | 78342    | 120738 | 54378    | 105583  | 543375   |
| Sorting cost                              | 2784     | 1567     | 7132   | 7187     | 1069    | 19739    |
| Harvesting cost                           | 16607    | 14528    | 32461  | 28396    | 35080   | 127072   |
| Marketing cost                            | 2818     | 1800     | 3516   | 1806     | 10029   | 19969    |
| Plantation cost                           | 29180    | 14056    | 15291  | 12245    | 3357    | 74130    |
| Planting material (seed, vegetative etc.) | 1967     | 14558    | 6928   | 3733     | 9927    | 37113    |
| Transport cost                            | 1825     | 5777     | 5974   | 12707    | 18757   | 45040    |
| Grand Total                               | 239514   | 130628   | 192040 | 120453   | 183803  | 8,66,438 |

Standard daily wages of workers during the survey period was Rs.150 for Men (skilled labourers) and Rs.100 (unskilled labourers), Rs. 45 for women and Rs.30 for boys.

**Table 4: Production and percentage contribution of various plant species found in homegardens towards annual income of households of Nongkwai, Mawriang, Umkrem, Nolikata and Ranikor villages of south Meghalaya**

| Plant species                   | Production (kg)/year |       | Rate(Rs. /kg) | Amount (Rs.)     |           | Gross Production (Rs.) | %      |
|---------------------------------|----------------------|-------|---------------|------------------|-----------|------------------------|--------|
|                                 | Self consumption     | Sale  |               | Self consumption | Sale      |                        |        |
| <i>Ananas comosus</i>           | 2315                 | 248   | 10            | 23150            | 2480      | 25630                  | 1.30   |
| <i>Areca catechu</i>            | 772                  | 3197  | 50            | 38600            | 159850    | 198450                 | 10.08  |
| <i>Artocarpus heterophyllus</i> | 30938                | 36397 | 0.83          | 25678.54         | 30209.51  | 55888.05               | 2.84   |
| <i>Averrhoa carambola</i>       | 192                  | 1029  | 15            | 2880             | 15435     | 18315                  | 0.93   |
| <i>Baccaurea sapida</i>         | 614                  | 904   | 17            | 10438            | 15368     | 25806                  | 1.31   |
| <i>Bambusa vulgaris</i>         | 825                  | 1925  | 0.5           | 412.5            | 962.5     | 1375                   | 0.07   |
| <i>Calamus floribundus</i>      | 96                   | 80    | 50            | 4800             | 4000      | 8800                   | 0.45   |
| <i>Carica papaya</i>            | 274                  | 716   | 30            | 8220             | 21480     | 29700                  | 1.51   |
| <i>Cinnamomum tamala</i>        | 12                   | 2016  | 8             | 96               | 16128     | 16224                  | 0.82   |
| <i>Citrus assamensis</i>        | 47                   | 57    | 60            | 2820             | 3420      | 6240                   | 0.32   |
| <i>Citrus karna</i>             | 214                  | 680   | 37.5          | 8025             | 25500     | 33525                  | 1.70   |
| <i>Citrus limon</i>             | 101                  | 21    | 30            | 3030             | 630       | 3660                   | 0.19   |
| <i>Citrus maxima</i>            | 6917                 | 2627  | 10            | 69170            | 26270     | 95440                  | 4.85   |
| <i>Citrus reticulata</i>        | 300                  | 168   | 30            | 9000             | 5040      | 14040                  | 0.71   |
| <i>Cocos nucifera</i>           | 488                  | 758   | 10            | 4880             | 7580      | 12460                  | 0.63   |
| <i>Cucurbita pepo</i>           | 31                   | 57    | 10            | 310              | 570       | 880                    | 0.04   |
| <i>Elaeagnus latifolia</i>      | 130                  | 2560  | 20            | 2600             | 51200     | 53800                  | 2.73   |
| <i>Emblica officinalis</i>      | 38                   | 144   | 15            | 570              | 2160      | 2730                   | 0.14   |
| <i>Garcinia cowa</i>            | 20                   | 200   | 30            | 600              | 6000      | 6600                   | 0.34   |
| <i>Litchi chinensis</i>         | 2001                 | 9946  | 25            | 50025            | 248650    | 298675                 | 15.16  |
| <i>Mangifera indica</i>         | 18069                | 7271  | 15            | 271035           | 109065    | 380100                 | 19.30  |
| <i>Musa sp.</i>                 | 1304                 | 2443  | 15            | 19560            | 36645     | 56205                  | 2.85   |
| <i>Piper betle</i>              | 3199                 | 10393 | 35            | 111965           | 363755    | 475720                 | 24.15  |
| <i>Piper longum</i>             | 3                    | 17    | 60            | 180              | 1020      | 1200                   | 0.06   |
| <i>Piper nigrum</i>             | 102                  | 915   | 90            | 9180             | 82350     | 91530                  | 4.65   |
| <i>Tamarindus indica</i>        | 320                  | 1141  | 15            | 4800             | 17115     | 21915                  | 1.11   |
| <i>Thysanolaena maxima</i>      | 48                   | 328   | 20            | 960              | 6560      | 7520                   | 0.38   |
| <i>Zea mays</i>                 | 59                   | 197   | 3             | 177              | 591       | 768                    | 0.04   |
| <i>Ziziphus mauritiana</i>      | 352                  | 1413  | 15            | 5280             | 21195     | 26475                  | 1.34   |
| Total                           |                      |       | -             | 6,88,442         | 12,81,229 | 19,69,671              | 100.00 |

**Table 5: Gross income, total expenditure and net income (Rs.) generated by homegardens of Nongkwai, Mawriang, Umkrem, Nolikata and Ranikor villages of south Meghalaya**

| Village  | Gross income (Revenue)  |                       | Net income              |                       |
|----------|-------------------------|-----------------------|-------------------------|-----------------------|
|          | Household <sup>-1</sup> | Hectare <sup>-1</sup> | Household <sup>-1</sup> | Hectare <sup>-1</sup> |
| Nongkwai | 2851.83                 | 45997.17              | 855.88                  | 26037.67              |
| Mawriang | 3232.76                 | 53356.27              | 1055.63                 | 34695.13              |
| Umkrem   | 4519.17                 | 128698.00             | 2385.39                 | 96691.33              |
| Nolikata | 4590.90                 | 98495.62              | 2181.84                 | 58344.62              |
| Ranikor  | 2378.26                 | 42194.89              | 1207.54                 | 5434.29               |
| Average  | 3,514.58                | 73,748.39             | 1,537.26                | 44,240.61             |

### Economic Value

Homegardens are a vital source of income for subsistence economy and contribute to the self-sufficiency of many rural households perched in remote places often secluded from markets and modern production centres. This is also true for the homegardens in other tropical countries (Kumar and Nair 2004; Peyre et al. 2006). The people usually work in the homegardens during

leisure time or after coming back from agriculture fields or early in the morning before going to the field. During summer season, farmers of the hill areas of the region prefer to work in their homegardens if the rain is heavy. This is because they have to walk on foot, for a minimum of 1-3 hours to reach to their agricultural fields. The undulated terrain of the land with narrow foot-paths are not convenient to walk during heavy rains.

The average annual gross production of homegardens in south Meghalaya was worth Rs.3514 per household, which was higher than the Kandyan Gardens of Sri Lanka (Rs.237) (Jacob and Alles 1987) but lesser as compared to homegardens in south west Bangladesh (Rs.11435) (Motiur et al. 2006), north-eastern Bangladesh (Rs.97441) (Motiur et al. 2005) and South Africa (Rs.6661) (High and Shakleton 2000). In terms of per hectare area the average annual gross income in south Meghalaya was Rs. 44,241 which was much higher than the Kandyan Gardens in Sri Lanka (Rs.1621) (Jacob and Alles

1987) and also the homegardens of South Africa (Rs.16520) (High and Shakleton 2000). The percentage contribution of homegardens towards peoples' average annual gross income was 7%, which was lesser than the contribution of homegardens in Indonesia (21.1%) (Soemarwoto 1987), Nicaragua (35%) (Mendez et al. 2001), south west and north-east Bangladesh (15.9% and 11.8%, respectively). With 7% contribution of homegardens towards average gross annual household income, it compares well with those from south and south-east Asia, where it varies between 6-54% (Kabir and Webb 2008).

### CONCLUSION

Homegardens of south Meghalaya enhance the livelihood of the people by providing food, construction materials, medicines and by contributing significantly to the gross annual income of the household. The income from homegardens depends on their size, number of marketable crops and management regime. Homegardens also ensure availability of products within the homestead which is particularly important in hill areas where agricultural fields are generally far off and travel to the fields is difficult and time consuming. The study brings to fore the biodiversity value of the homegardens. It is revealed that they are rich repository of wild as well as domesticated plants. Homegardens harbour species found in the wild and also provide habitat for cultivation of economically useful plants. Thus these predominantly man made ecosystems can be placed between forest and agriculture ecosystems in the forest-agriculture continuum, that provide food security to the people and act as safety net in cases of exigency.

### RECOMMENDATIONS

Homegardens are an important element of rural landscape of many tropical countries particularly in south and south east Asia and they play a vital role in the predominantly bioresource based economy of the region. However, they have not received due attention in the policies and programmes of government, the reason being they neither fall in agriculture nor in forestry sectors. Based on the study it is recommended that promotion of homegardens be included in the projects aiming to enhance the livelihoods of rural poor on sustainable basis.

### ACKNOWLEDGEMENTS

We are thankful to the Head, Department of Environmental Studies, North-Eastern Hill University, Shillong for providing necessary laboratories facilities. We are also deeply indebted to Dr. (Mrs.) S. J. Phukan, Scientist (D) Deputy Director, BSI, Eastern Circle, Shillong; for allowing us to consult the herbaria and deputing her staff to help in the identification of plant specimens. A word of gratitude to Mr. Sumendro who helped us in preparing the map.

### REFERENCES

- Balakrishnan NP 1981-1983. *Flora of Jowai, Meghalaya, Vol. I & II*. Howrah: Botanical Survey of India.
- Das T, Das AK 2005. Inventorying plant biodiversity in homegardens: A case study in Barak Valley, North-East Assam. *Current Science*, 89: 155-163.
- High C, Shakleton CM 2000. The comparative value of wild and domestic plants in homegardens of a South African rural village. *Agroforestry Systems*, 48: 141-156.
- Hochegger K 1998. *Farming like the Forests-Traditional Homegarden System in Sri Lanka*. Weikersheim, Germany: Margraf.
- Jacob VJ, Alles WS 1987. Kandyan Gardens of Sri Lanka. *Agroforestry Systems*, 5: 125-137.
- Kabir ME, Webb EL 2008. Can homegardens conserve biodiversity in Bangladesh? *Biotropica*, 40: 95-103.
- Kanjilal UN, Kanjilal PC, Das A, De RN, Bor NL 1934-1940. *Flora of Assam*. 5 Vols. Shillong: Govt. Press.
- Karyono 1990. Homegardens in Java: their structure and functions. In: K Landauer, M Brazil (Eds.): *Tropical Homegardens*. Tokyo, Japan: United Nations University Press, pp. 138-146.
- Kehlenbeck K, Maass BL 2004. Crop diversity and classification of homegardens in Central Sulawesi, Indonesia. *Agroforestry Systems*, 63: 53-62.
- Khan ML, Shaily M, Kamaljit SB 1997. Effectiveness of the protected area network in biodiversity conservation, a case study of Meghalaya state. *Biodiversity and Conservation*, 6: 853-865.
- Kingston C, Mishra BP, Nisha BS, Jeeva S, Livingstone C, Laloo RC 2006. Diversity and distribution of economically important plants in traditional homegardens of Kanyakumari district Tamil Nadu, Southern Peninsular India. *J Natcon*, 18: 4154.
- Kumar BM, Nair PKR 2004. The enigma of tropical homegardens. *Agroforestry Systems*, 61: 135-152.
- Mendez VE, Lok R, Somaribba E 2001. Interdisciplinary analysis of homegardens in Nicaragua: microzonation, plant use and socio-economic importance. *Agroforestry Systems*, 51: 85-96.
- Misra R 1968. *Ecology Work Book*. New Delhi: Oxford and IBH.
- Motiur MR, Furukawa Y, Kawata I 2005. Homestead forest resources and their role in household economy: A case study in the villages of Gazipur Upazila of central Bangladesh. *Small-scale Forest Economics, Management and Policy*, 4: 359-376.
- Motiur RM, Furukawa Y, Kawata I, Rahman M, Alam M 2006. Role of homestead forest in household economy and factors affecting forest production: a case study

- in southwest Bangladesh. *Journal of Forest Research*, 11: 89-97.
- Mueller-Doimbois D, Ellenberg H 1974. *Aims and Methods of Vegetation Ecology*. New York: John Wiley.
- Mukherjee N 1993. *Participatory Methods and Rural Knowledge. Participatory Rural Appraisal Methodology and Applications*. New Delhi: Concept Publishing Company.
- Nautiyal S, Vimla B, Rao KS, Maikhuri RK 2008. The Role of Cultural Values in Agrobiodiversity Conservation: A Case Study from Uttarakhand, Himalaya. *Journal of Human Ecology*, 23: 1-6.
- Peyre A, Guidal A, Wiersum KF, Bongers F 2006. Dynamics of homegarden structure and function in Kerala, India. *Agroforestry Systems*, 66: 101-115.
- Quiroz C, Guitierrez M, Rodriguez D, Perez D, Ynfante J, Gamez J, Perez de Fernandez T, Marques A, Pacheco W 2002. Homegarden and *in situ* conservation in Agrobiodiversity- Venezuelan component. In: JW Watson, PB Eyzaguirre (Eds.): *Homegardens and in situ Conservation of Plant Genetic Resources in Farming Systems. Proceedings of the Second International Homegardens Workshop*, 17-19 July 2002. Witzenhausen, Germany. Rome Italy: International plant genetic resources Institute (IPGRI), pp. 73-82.
- Soemarwoto O 1987. Homegardens: A traditional agro-forestry system with promising future. In: H Steppeler, PKR Nair (Eds.): *A Decade of Development*. Nairobi: ICRAF, pp-157-170.
- Shannon CE, Wiener W 1963. *The Mathematical Theory of Communication*. Urbana: University Illinois Press.
- Tiwari BK 2005. Forest biodiversity management and livelihood enhancing practices of War Khasi of Meghalaya, India. In: Y Thomas, M Karki, K Gurung, D Parajuli (Eds.): *Himalayan Medicinal and Aromatic Plants, Balancing Use and Conservation*. His majesty Government of Nepal Ministry of Forests and Soil Conservation. pp. 240-255.
- Tynsong H, Tiwari BK 2008. Traditional knowledge associated with fish harvesting practices of War Khasi community of Meghalaya. *Indian Journal of Traditional Knowledge*, 7: 618-623.
- Tynsong H 2009. *Plant Diversity and NTFP Management in Community Forests of War Area Meghalaya*. Ph.D. Thesis, Unpublished. Shillong: North-Eastern Hill University.
- Wezel A, Bender S 2003. Plant species diversity of homegardens of Cuba and its significance for household food supply. *Agroforestry Systems*, 57: 39-49.