



Extraction of High Quality Genomic DNA for Genetic Diversity and Molecular Marker Studies from Leaves of Makhana (*Euryale ferox*)

Nitish Kumar^{1*}, Hemant Kumar¹ and Swati Kumari¹

**¹Centre for Biological Sciences (Biotechnology), School of Earth, Biological and
Environmental Sciences, Central University of South Bihar, Patna 800 014, Bihar, India**

**²Centre for Education (Life Science), Central University of South Bihar, Gaya Campus,
Gaya 823 001, Bihar, India**

KEYWORDS DNA Extraction. *Euryale ferox*. Molecular Diversity. Molecular Marker. Phenol

ABSTRACT Genetic diversity study of plants through molecular markers depends on both quality and quantity of DNA. The standard protocols and commercially available kits failed to isolate high quality DNA from leaves tissue of Makhana (*Euryale ferox* Salisb.) due to the occurrence of the high amount of polysaccharides, polyphenols and proteins. A cetyl trimethyl ammonium bromide (CTAB) protocol has been optimized for isolation of high quality genomic DNA from the leaves of *E. ferox*. The optimized protocol yielded up to 4081 ng per gram leaves tissue of good quality DNA, which was free of polysaccharide, polyphenols and protein. The problems of DNA degradation, contamination, and low yield due to irreversible binding of phenolic compounds and co-precipitation of polysaccharides with DNA were avoided by this method. The isolated DNA has excellent spectral qualities, and efficiently digested by restriction endonucleases and it is also suitable for development of molecular markers such as RAPD and ISSR.