

PTC Taste Sensitivity Among the Muslims of Calcutta, West Bengal

J. Hoque¹ and A.R. Bandyopadhyay²

1. Department of Anthropology, University of Calcutta, Calcutta 700 019, West Bengal, India

2. Department of Anthropology, Palli Charcha Kendra, Visva Bharati, Sriniketan 731 236, Birbhum, West Bengal, India

KEY WORDS PTC Tasting Ability. Muslims. Polymorphism. Variation.

ABSTRACT A total of 344 individuals from Bengalee and non-Bengalee Sunni sect Muslim community of Calcutta, West Bengal was tested for PTC tasting ability. Although no sex difference was found among these two groups of Muslims, the higher mean threshold value for tasters among the Bengali Muslims compared to that of the non Bengalee Muslims was evident.

INTRODUCTION

Variable taste sensitivity of phenylthiocarbamide (PTC) has been widely used as a polymorphic genetic marker in human populations. The frequency of tasters in India ranges from 98% in the Khasis of Assam (Miki et al., 1960) to 26% among the Southern Pahira (Basu et al., 1966).

Although the PTC tasting ability was searched in Muslims of different states of India (Srivastava, 1959; Mahapatra and Das, 1968; Kalla, 1971; Das, 1984-85; Devi and Singh, 1990). Only a few studies on Muslims (Bhattacharjee, 1956; Dash Sharma and Haque, 1978) of West Bengal has been done (Bhasin et al., 1992). The present paper deals with the PTC taste sensitivity of Muslims of Calcutta.

SUBJECTS AND METHODS

Subject for the present study comprised of 344 individuals of Bengalee (225) and non-Bengalee (119) Sunni Muslims from Calcutta, West Bengal. Bengalee Muslims are the original inhabitants of West Bengal, while non-Bengalee Muslims are mostly migrated from outside the state e.g. from Uttar Pradesh, Bihar etc. The data were collected from the Muslim

schools as well as different Muslims families of Calcutta, West Bengal. PTC tasting ability was tested by the sorting techniques of Kalmus et al. (1981), using 13 serial dilutions of PTC.

RESULTS AND DISCUSSION

The distribution of taste threshold showed bimodality in both sexes of Bengalee as well as non-Bengalee Muslim samples of Calcutta (Table 1).

The distribution of taster and non-taster proportions in both Bengalee and non-Bengalee Muslim samples, however, showed no sex differences (Table 2). Examination on inter group variation revealed higher taster individuals than the expected (130.8) among the Bengalee Muslims compared to that of the non-Bengalee Muslims. Moreover, allele frequency estimation (Table 2) also revealed the higher frequency of *t* allele ($\chi^2 = 1.78$, d.f.1, $P < 0.20$) among the non-Bengalee Muslims compared to that of the Bengalee Muslim samples of Calcutta.

Finding of higher recessive *t* allele in non-Bengalee Muslims of Calcutta seems to be related to higher incidence of consanguineous marriage among them (Basu, 1985; Hoque and Bandyopadhyay, 1992, 1993).

Allele frequency of *t* in the present study, however, showed good agreement with other Muslim data from West Bengal (Bhattacharjee, 1956; Das-Sharma and Haque, 1978; Hoque, 1994) and as well as from other States (Srivastava, 1956; Kalla, 1971).

Table 1: P.T.C. taste threshold distribution among the Muslims of Calcutta

Popula- tion	Sex	n	<1	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Bengali	Male	118	17	5	8	6	2	5	9	11	10	14	9	8	7	4	3
Muslims	%		14.40	4.23	6.78	5.08	1.69	4.23	7.62	9.32	8.47	11.86	7.62	6.78	5.93	3.39	2.54
	Female	107	22	4	7	6	5	2	6	7	9	11	8	7	6	5	2
	%		20.56	3.74	6.54	5.60	4.67	1.87	6.60	6.54	8.41	10.28	7.48	6.54	5.60	4.67	1.87
	Total	225	39	9	15	12	7	7	15	18	19	25	17	15	13	9	5
	%		17.33	4.00	6.67	5.33	3.11	3.11	6.67	8.00	8.44	11.11	7.55	6.67	5.78	4.00	2.22
Non- Bengali	Male		14	4	5	3	1	5	7	6	8	7	3	2	1	1	1
Muslims	%		20.59	5.88	7.35	4.41	1.47	7.35	10.29	8.82	11.76	10.29	4.41	2.94	1.47	1.47	1.47
	Female	51	12	5	4	3	4	5	3	3	2	4	2	2	1	1	0
	%		23.30	9.80	7.84	5.88	7.84	9.80	5.88	5.88	3.92	7.84	3.92	3.92	1.96	1.96	0.00
	Total	119	26	9	9	6	5	10	10	9	10	11	5	4	2	2	1
	%		21.84	7.56	7.56	5.04	4.20	8.40	8.40	7.56	8.40	9.24	4.20	3.36	1.68	1.68	0.84

Table 2: Allele frequencies of PTC tasting ability among the Muslims of Calcutta

Population	Sex	Position of antimode	Taster		Non-taster		Allele frequency	
			No. Obs.	% Obs.	No. Obs.	% Obs.	T	t
Bengalee	Male	4	80	67.80	38	32.20	0.425	0.575
Muslims	Female	5	61	57.00	46	43.00	0.337	0.663
	Total	5	136	60.44	89	39.56	0.371	0.629
Non-Bengalee	Male	4	41	60.30	27	39.70	0.370	0.630
Muslims	Female	3	27	52.94	24	47.06	0.314	0.686
	Total	4	64	53.79	55	46.21	0.320	0.680

ACKNOWLEDGEMENTS

Authors are grateful to Prof. A.R. Banerjee and Prof. D.P. Mukherjee of Department of Anthropology, University of Calcutta for their useful advice in the preparation of manuscript. Thanks are also due to Muslims of Calcutta for their cooperation in data collection.

REFERENCES

- Basu, A., Sarkar, J., Roy, B., Dasgupta, P. and Sarkar, S.S.: Anthro-po-genetic investigation in Dalma and Ajodhya hills and their neighborhood. *Sci. Cult.*, 32: 273-275 (1966).
- Basu, M.P.: *Anthropological Profile of the Muslims of Calcutta*. Anthropological Survey of India, Govt. of India, Calcutta (1985).
- Bhasin, M.K., Walter, H. and Danker-Hopfe, H.: *Distribution of Genetical, Morphological and Behavioral Traits Among Peoples of Indian Region*. Kamla-Raj Enterprises, Delhi (1992).
- Bhattacharjee, P.N.: A genetic survey in the Rarhi Brahmin and Muslim of West Bengal: ABO, MN, Rh blood groups, ABH secretion, sickle cell, PTC taste, middle phalangeal hair and colour blindness. *Bull. Anthropol. Surv. India*, 5: 18-28 (1956).
- Dash Sharma, P. and Haque, M.: Phenylthiourea tasting ability in a Muslim village (Sabal Singlapur) of Hooghly district (W.B.). *Indian Anthropologist*, 8: 55-58 (1978).
- Devi, N.G.S. and Singh, T.S.: Gene difference at two loci between the Muslims of Manipur and Assam. *J. Ind. Anthropol. Soc.*, 25: 35-38 (1990).
- Hoque, J. and Bandyopadhyay, A.R.: Consanguinity among the Muslim of West Bengal. *J. Ind. Anthropol. Soc.*, 27: 90-94 (1992).

- Hoque, J. and Bandyopadhyay, A.R.: Distribution of red cell H substance among the Muslim of Calcutta. *J. Hum. Ecol.*, 4: 135-138 (1993).
- Hoque, J.: *Anthropo-genetic Study of the Sunni and the Shia Muslims of Bengal*. Ph.D. (Thesis submitted), University of Calcutta, Calcutta (1994).
- Kalla, A.K.: P.T.C. tasting ability and colour blindness among the Muslims. *Anthropologist*, 18: 45-48 (1971).
- Kalmus, H., Mourant, A.E. and Sunderland, E. : PTC taste-tasting. pp. 107-114. In: *Practical Human Biology*. J.S. Weiner and J.S. Lourie (Eds.). Academic Press, London (1981).
- Mahapatra, U.D. and Das, P.: Taste threshold for phenylthiocarbamide in some endogamous groups of Assam. *Anthropologist*, 15: 25-32 (1968).
- Miki, T., Tanaka, T. and Furuhashi, T.: On the distribution of ABO blood groups and the taste ability for phenylthiocarbamide (PTC) of the Lepchas and the Khasis. *Proc. Jap. Acad.*, 36: 78-80 (1960).
- Srivastava, R.P.: Measurement of taste sensitivity of Phenylthiourea (PTC) in Uttar Pradesh. *East. Anthropol.*, 12: 267-272 (1959).