

Seasonal Variations of Diatoms Epiphytic on the Roots of Water Hyacinth [*Eichhornia crassipes* (Mart.) SOLMS]

Parminder Kaur and Naresh K. Mehra

Limnology Unit, Department of Zoology, University of Delhi, Delhi 110 007, India

KEY WORDS Epiphytic diatoms. Water Hyacinth. Seasonal Variations. Water Quality.

ABSTRACT In this investigation, an attempt has been made to study the seasonal variations of epiphytic microflora, especially diatoms, attached to the adventitious roots of water hyacinth growing in lotic waters polluted by urban and agro-industrial wastes. Epiphytic algae were largely represented by Bacillariophyceae, Chlorophyceae, Euglenophyceae, and Cyanobacteria in the decreasing order of abundance. A total of 59 genera belonging to above algal groups were recorded. Although, qualitatively diatoms ranked second to green algae, quantitatively their relative percentage to total algal population was the highest throughout the period of investigation and attained maxima during spring. The dominant forms were *Navicula*, *Nitzschia*, *Synedra*, *Fragilaria*, *Cyclotella* and *Gomphonema* at various sites. Data were subjected to analysis of community parameters viz., Shannon-Weiner index (H') and Morisita's similarity index, as also correlations between the epiphytic diatoms and environmental variables.

INTRODUCTION

The role of periphytic communities in monitoring eutrophication and pollution has attracted the attention of limnologists since past two decades (Lowe, 1974; Lange-Bertalot, 1979; Goldsborough and Robinson, 1985; Sabater et al., 1987; Watanbe et al., 1988; Nather Khan, 1990; Rushforth and Brock, 1991; Nwankwo and Akinsoji, 1992; Müller, 1994). However, most of the earlier studies on water quality evaluation as also seasonal variations include exposure of artificial substrates for periphytic colonization in free-water column (Castenholz, 1960; Saldeckova, 1962; Patrick 1973; Blinn, 1986; Sabater et al., 1987). In comparison, a few investigations have been conducted to study seasonal periodicity of algal assemblages on natural substrates e.g., floating and submerged macrophytes, rocks etc.

The objective of this study was to investigate the seasonal variations in abundance as well as species composition of algal assemblages with special reference to diatoms epiphytic on floating macrophyte *Eichhornia crassipes* growing in lotic waters and relate them with physico-chemical parameters.

MATERIALS AND METHODS

Study Sites

These investigations were conducted in a sub-tropical basin of river Yamuna and an associated drain-Najafgarh drain. River Yamuna flows for 48 kms through the city states of Delhi from north to south, leaving the city near Okhla barrage. The river has an average depth of 1.25 m and breadth varies between 200-400 m along the course (Fig. 1). Najafgarh drain and 16 other drains of various catchment areas discharge untreated sewage and agro-industrial wastes directly into the river Yamuna. As a result, various stages of eutrophication are noticed in the drains as also in the river basin. Four stations were selected for this study. Stations A and B were located on the western bank of river Yamuna. Stations C and D were situated on Najafgarh drain, which alone contributes 51% of untreated BOD load into the river Yamuna.

Methods of Analysis

Water samples for the analysis of physico-chemical parameters and epiphytic diatoms attached to the roots of water hyacinth were collected once every month for a year. Procedures adopted for the analysis of chemical parameters viz., hardness, chlorides, alkalinity, silica,




Fig. 1. Map depicting river Yamuna through its course in the National Capital Territory of Delhi showing location of various drains and sampling stations viz., A, B, C, and D

ortho-phosphates, nitrites, BOD, COD, DO were according to Standard Methods (APHA, 1989) and nitrates according to Fresenius et al. (1987). For epiphytic samples, water hyacinth plants were collected with minimum disturbance. A method was standardised for the quantification of epiphyton on the roots of water hyacinth. Roots were detached from plants and taken in proportionate amount of filtered water sample in a beaker and shaken vigorously. Cells were counted in Sedgewick-Rafter cell. Results have been expressed as number of cells/root unit (for details refer Kaur, 1991; Kaur and Mehra, 1994). Diatoms were identified using frustule size, shape, and ornamentation (Tiffany and Britton, 1952; Desikachary, 1959; Krammer and Lange-Bertalot, 1986, 1988, 1989).

Data Analysis

Data were subjected to Shannon-Weiner index (H'). Similarity in genera among various seasons at different stations was determined using Morisita's similarity index. The distribution of epiphytic diatoms were correlated with physico-chemical variables and other epiphytic algal groups using computer programme by Krebs (1989).

RESULTS

Temperature at all the stations ranged between 15° and 34°C. pH of the water ranged from neutral to slightly alkaline at the study sites. Conductivity was highest at station C and it ranged between 1310 and 1630 μScm^{-1} , whereas at other stations it varied from 510 and 820 μScm^{-1} . In all the stations water was generally hard. During the study period, DO was absent at stations C and D, whereas at stations A and B it remained within a low range. Maximum concentration of chlorides was recorded during winter at stations A, B and C and during monsoon at station D. The concentration of silicates was high at all the sampling sites and recorded maxima during winter. Higher concen-

trations of nitrates were recorded in all the sampling sites and they attained maxima during winter at stations A and D, and during spring at stations B and C. Relatively higher concentrations of ortho-phosphates were observed at stations C and D. BOD was high at all the stations and recorded maxima during monsoon. Range of variations for the selected environmental variables are shown in table 1.

The averages of some of the selected physico-chemical parameters were subjected to the weight score of Crude Pollution Index (DeZwart, 1991) to ascertain water quality. Data analysis revealed that quality of water at stations A, B and C varied from moderately to highly polluted during the study period, where-

Table 1: Range of variations in physico-chemical characteristics at the study sites

Stations/ Parameters	A	B	C	D
Temp°C	16-30	15-31.5	18.5-32	19.5-34
pH	7.3-7.6	6.8-7.8	7.3-8.3	7.2-7.6
DO	1.8-4.8	0.8-1.8	0	0
Conductivity, μScm^{-1}	570-790	575-740	1310-1630	510-820
TOT. HARD.	188-218	160-220	280-390	152-226
Chlorides	42-105	37-98	23.2-192.2	38.4-194
Silicates	7.6-18.5	5.2-30.4	6.4-33.0	8-21.2
Nitrates	0.1-1.06	0.1-0.89	0.21-0.29	0-0.49
Phosphates	0.16-0.87	0.42-0.85	1.09-1.38	0.95-1.36
BOD ₅	31.4-87.2	36.4-76.0	51.3-126	54.3-128.6

Results other than temperature, pH and conductivity, are expressed in mgL^{-1} . [TOT. HARD.-Total hardness]

as station D continued to be highly polluted throughout the period of investigation.

Epiphytic algae on the roots of water hyacinth were largely represented by Bacillariophyceae, Chlorophyceae, Cyanobacteria and Euglenophyceae. A total of 59 genera of algae have been identified, of which 18 belonged to diatoms, 29 to green algae, 6 to blue-green algae and remaining to euglenoids. Although, qualitatively diatoms ranked second to green algae, quantitatively they were most abundant at all the stations during the period of investigation. An analysis of epiphytic diatom popu-

lation from different stations revealed that their abundance varied markedly during different seasons (Figs. 2-5).

Maximum diversity in diatom population was recorded at station A, whereas maximum density was observed at station C. However, among all the stations, peak for diatom population was recorded during spring. Maxima with 1310 diatoms/root unit and 1945 diatoms/root unit were recorded at stations A and B, respectively (Figs. 2,3). The most common forms were *Fragilaria*, *Pinnularia*, *Cymbella*, *Gomphonema*, *Melosira* and *Tabellaria* (Table 2). Among the eight taxa recorded from each of the remaining two stations, *Gomphonema*, *Nitzschia*, *Cyclotella*, and *Stauroneis* with 2135 diatoms/root unit contributed to maxima at station C, whereas *Epithemia*, *Nitzschia*, *Cyclotella*, and *Fragilaria* contributed to the peak with 1985 diatoms/root unit at station D (Table 2, Fig. 5).

Species diversity of epiphytic diatoms ranged between 1.4 and 3.48 at station A, be-

Table 2: List of epiphytic diatoms and their relative abundance during the period of investigation. [- =absent; x =rare; xx =frequent; xxx=very frequent; xxxx =abundant]

	Stations			
	A	B	C	D
<i>Cyclotella</i>	xx	xx	xxxx	xxxx
<i>Cymbella</i>	x	xx	-	xxx
<i>Diatoma</i>	xx	-	xx	-
<i>Epithemia</i>	-	-	-	xxx
<i>Fragilaria</i>	xx	xxx	xxx	xx
<i>Frustulia</i>	x	x	-	x
<i>Gomphonema</i>	xxx	xx	xxx	x
<i>Gyrosigma</i>	xx	xx	xxx	xxx
<i>Gomphonosis</i>	x	x	-	-
<i>Melosira</i>	x	xx	xx	xxx
<i>Meridion</i>	-	-	x	x
<i>Navicula</i>	xx	xxx	xx	xxx
<i>Neidium</i>	x	x	xxx	xxx
<i>Nitzschia</i>	xx	xx	xxxx	xxxx
<i>Pinnularia</i>	xx	xxx	xxx	x
<i>Stauroneis</i>	xxx	xxx	xxx	xxx
<i>Surirella</i>	xx	xx	xxx	xx
<i>Synedra</i>	x	xxx	xxxx	xxx
<i>Tabellaria</i>	x	xx	xx	xxx

tween 2.1 and 3.0 at station B, between 1.8 and 2.7 at station C and between 2.5 and 2.9 at station D (Fig. 6).

Comparison of epiphytic diatoms during different seasons among four stations revealed that maximum similarity was recorded between stations A and B during autumn and monsoon, between stations B and D during spring and between stations A and C during summer (Table 4).

Correlations between epiphytic biota and environmental variables revealed that weak to strong negative correlation existed between

Table 3: Correlation coefficient analysis of total epiphytic diatoms with physico-chemical parameters and between diatoms and epiphytic flora

	Stations			
	A	B	C	D
A. Physico-Chemical parameters				
Temperature	-.501*	-	.622*	-.558**
pH	-.820*	-.722*	-.521**	-.576**
Chlorides	-	.543**	-	-.533**
B.O.D. ₅	-.764*	-.634	-	-.525**
NO ₃ -N	-.681*	.711*	.804*	.522**
PO ₄ -P	.944*	-.524**	.664*	.685*
SiO ₂	.744*	.864*	.741*	.630*
B. Epiphytic Flora				
Chlorophyceae	.735*	-	.595**	-
Cyanobacteria	.860*	.925*	.789*	.745*
Euglenophyceae	.936*	.905*	.597**	.616*
Total Algae	.942*	.910*	.802*	.831*

Level of significance : * = at 1% level

** = at 5% level

- = not significant

diatoms and temperature, DO, pH, Cl, BOD, whereas positive correlations were recorded with conductivity, PO₄-P, total algae, Chlorophyceae and Cyanobacteria (Table 3).

DISCUSSION

The physico-chemical characteristics of River Yamuna have been studied in detail during last 30 years (Rai, 1962, 1974a, b; Mathur, 1965; Kundra et al., 1977; Subramanian and




Fig. 2. Seasonal variations in the total epiphytic algal cells highlighting the relative abundance of diatoms at station A. Abbreviations of predominant species of diatoms are as follows : PIN - *Pinnularia*; CYC - *Cyclotella*; SYN - *Synedra*; FRA - *Fragilaria*; NIT - *Nitzschia*; STA - *Stauroneis*; GOM - *Gomphonema*; NAV - *Navicula*




Fig. 3. Seasonal variations in the total epiphytic algal cells highlighting the relative abundance of diatoms at station B. Abbreviations of predominant species of diatoms are as follows : PIN - *Pinnularia*; SYN - *Synedra*; NIT - *Nitzschia*; NAV - *Navicula*; STA - *Stauroneis*; GOM - *Gomphonema*




Fig. 4. Seasonal variations in the total epiphytic algal cells highlighting the relative abundance of diatoms at station C. Abbreviations of predominant species of diatoms are as follows : SUR - *Surirella*; FRU - *Frustulia*; NIT - *Nitzschia*; GOM - *Gomphonema*; PIN - *Pinnularia*; CYC - *Cyclotella*; SYN - *Synedra*; GYR - *Gyrosigma*; STA - *Stauroneis*




Fig. 5. Seasonal variations in the total epiphytic algal cells highlighting the relative abundance of diatoms at station D. Abbreviations of predominant species of diatoms are as follows : TAB - *Tabellaria*; NEI - *Neidium*; GYR - *Gyrosigma*; NIT - *Nitzschia*; NAV - *Navicula*; SUR - *Surirella*; FRA - *Fragilaria*; MEL - *Melosira*; EPI - *Epithemia*; DIA - *Diatoma*; CYM - *Cymbella*; STA - *Stauroneis*; GOM - *Gomphonema*; PIN - *Pinnularia*




Fig. 6. Shannon-Weiner diversity indices for epiphytic diatoms at four sampling stations during different seasons

Table 4: Similarity Index of the epiphytic diatoms among four stations during different seasons. Bold values indicate maximum similarities between the stations

Seasons	Stations/ Stations	A	B	C	D
Autumn	A	1.01			
	B	.80	1.01		
	C	.35	.20	1.01	
	D	.72	.52	.42	1.01
Winter	A	1.01			
	B	.83	1.02		
	C	.66	.39	1.01	
	D	.52	.55	.33	1.01
Spring	A	1.01			
	B	.76	1.01		
	C	.59	.45	1.00	
	D	.59	.80	.54	1.00
Summer	A	1.01			
	B	.55	1.01		
	C	.57	.38	1.00	
	D	.29	.34	.28	1.00
Monsoon	A	1.01			
	B	.41	1.01		
	C	.23	.21	1.01	
	D	.00	.07	.38	1.01

Sitasawad, 1984; Bhargava, 1985; Trivedi et al., 1985; Khurshid, 1989; Menon, 1990) and that of Najafgarh drain by Bulusu and Sharma (1966) and Dakshini and Soni (1979, 1982). A comparison of the present data with those of earlier studies indicates that water quality has further deteriorated at both the sites. Although, the nutrient load in the segments of river Yamuna as also in Najafgarh drain has increased, they still support a rich and diverse biotic community. Recent studies conducted by DeZwart (1991) revealed that as a result of above physico-chemical changes, the composition of plankton populations have also undergone significant variations during last thirty years.

Earlier studies on species composition and seasonal dynamics of phytoplankton were made on river Yamuna at Delhi (Rai, 1962, 1974 b; Hortobagyi, 1969). These studies revealed that diatoms dominated the phytoplankton followed by green and blue-green algae. In addition, seasonal progressions involving basi-

cally year-round predominance of diatoms with short pulse of green algae have also been reported in Asian rivers (Kundra et al., 1977; Trivedi et al., 1985; Nather Khan, 1990 and Nwankwo and Akinsoji, 1992), European rivers (Swale et al., 1969; Kiss, 1984; Simm, 1985; Descy, 1987), and Australian rivers (Chessman, 1985).

In the present study, it was observed that among the periphytic algal communities, qualitatively green algae dominated the epiphytic assemblages on the roots of water hyacinth, whereas numerically diatoms were more predominant throughout the investigation. This is in conformity with earlier reports of Nwankwo (1984, 1986). Similar observations have also been made with reference to other macrophytes i.e., *Lemna minor* (Goldsborough and Robinson, 1985), on *Pistia* sp. (Nwankwo and Akinsoji, 1988), *Potamogeton* sp. (Cattaneo and Kalff, 1978), *Phragmites australis* (Müller, 1994) and on artificial substrates (Castenholz, 1960; Cattaneo and Kalff, 1978; Nather Khan, 1990).

During this investigation, it was observed that among the epiphytic algal community, diatoms recorded maxima during spring at all the stations. This is in concurrence with observations of Kiss (1984) in river Hungary, Goldsborough and Robinson (1985) in Delta marsh, Canada, Watenbe et al. (1988), Acs and Kiss (1991) in river Danube, Hungary, Nwankwo and Akinsoji (1992) on coastal waters in Nigeria and Buczko and Acs (1994) in Danube river.

Lund (1950), Lange-Bertalot (1979), Sabater et al. (1987) and Nather Khan (1990) have observed that following spring maxima, diatom population declined at the onset of summer, which have been attributed to exhaustion of dissolved silica. In our study also, decline in diatom population during summer was due to lowering of dissolved silica levels.

Seki et al. (1988) reported that forms like *Fragilaria*, *Melosira* and *Synedra* were the most abundant diatoms during the growth period of water hyacinth. In our study, predominance of above forms was observed during sum-

mer, which is the growth period of macrophyte under observation.

Seasonal periodicity in the composition as also abundance of diatoms have been recorded at different stations and this can be correlated to the changing physico-chemical profiles of sampling sites. Our observations are in accordance with those of Lowe (1974), Lange-Bertalot (1979), Kiss (1984), Sabater et al. (1987), Nather Khan (1990), and Müller (1994).

Sabater et al. (1987, 1988) and Biggs (1989) recorded negative correlations of epiphytic diatoms with temperature, DO, SiO₂-Si and positive correlations with PO₄-P and conductivity. Similar correlations have been recorded in this study also.

The use of epiphytic diatoms in monitoring water quality has been well documented in literature. Attempt has been made in this study to analyze the composition of epiphytic diatoms in relation to water quality. Patrick and Hohn (1956) had observed that due to organic pollution, rare and intolerant species do not survive, whereas more tolerant species tend to become exclusively dominant. In our study also certain tolerant forms of diatoms viz., *Nitzschia*, *Cyclotella* and *Synedra* were recorded from all the stations, but they were more predominant at stations on Najafgarh drain which has a regular inflow of untreated sewage.

During the period of investigation, species diversity was low in the highly polluted stations on Najafgarh drain. Similar observations have been reported by Dakshini and Soni (1979, 1982).

Sabater et al. (1987), Nather Khan (1990), and Nwankwo and Akinsoji (1992) have listed some species of diatoms as indicators of water quality. Some of these taxa have also been recorded in the present study.

CONCLUSIONS

1. Seasonal changes of epiphytic diatoms on the roots of water hyacinth were recorded at all the stations during spring season.

2. Differences in species diversity at stations could be related to changes in water quality.
3. Some diatom genera reported to act as bio-indicators were recorded in this study also.

ACKNOWLEDGEMENTS

Thanks are due to Council of Scientific and Industrial Research, CSIR, New Delhi, for the award of a research fellowship to P.Kaur during the present work.

REFERENCES

- Acs, E. and Kiss, K.T. : Investigation of periphytic algae in the Danube at God (1669 river km, Hungary). *Algalogical Studies*, **62** : 42-67 (1991).
- American Public Health Association. *Standard Methods for the Examination of Water and Waste water*, 17th ed. APHA, Washington DC. (1989).
- Bhargava, D.S. : Water quality variations and control technology of Yamuna river. *Environ. Pollut.*, **37A** : 355-376 (1985).
- Biggs, B.J.F. : Biomonitoring of organic pollution using periphyton, South Branch, Canterbury, New Zealand. *New Zealand J. Mar. and Freshwat. Res.*, **23** : 263-274 (1989).
- Blinn, D.W. : Pros and cons of artificial versus natural substrata - In : M. Ricard, (Ed.), *Proc. Eighth Int. Diatom Symp. - Koelitz, Koenigstein* 776-781 pp. (1986).
- Buckzo, K. and Acs, E. : Algalogical studies on the periphyton in the branch - system of the Danube at Cikolasziget (Hungary). *Verein. Int. Ver. Limnol.*, **25** : 1680-1683 (1989).
- Bulusu, K.R. and Sharma, V.P. : Survey of Najafgarh drain downstream of industrial area, *Environ. Hlth.*, **8** : 103-111 (1966).
- Castenholz, R.W. : Seasonal changes in the algae of freshwater and saline lakes in the lower Grand Coulee, Washington. *Limnol. Oceanogr.*, **5** : 1-28, (1960).
- Cattaneo, A. and Kalff, J. : Seasonal changes in the epiphytic community of natural and artificial macrophytes in lake Memphremagog (Que & Vt). *Hydrobiologia*, **60** : 135-144 (1978).
- Chessman, B.C. : Artificial-substratum periphyton and water quality in the lower La Trobe River, Victoria. *Aust. J. Freshwater Res.*, **36** : 855-871 (1985).
- Dakshini, K.M.M. and Soni, J.K. : Water quality of sewage drains entering Yamuna in Delhi, *Indian J. Environ. Health*, **21** : 354-360 (1979).
- Dakshini, K.M.M. and Soni, J.K. : Diatom distribution and status of organic pollution in sewage drains. *Hydrobiologia*, **87** : 205-209, (1982).
- Desay, J.P. : Phytoplankton composition and dynamics in the River Meuse (Belgium). *Arch. Hydrobiol.*, **78** : 225-245 (1987).
- Desikachary, T.V. : *Cyanophyta*. Indian Council of Agricultural Research, New Delhi, (1959).
- DeZwart, D. : *Report on an Expert Mission for the Evaluation of Yamuna River Biomonitoring Data*. Report No 768602008, National Institute of Public Health and Environmental Protection, Bithoven, Netherlands : 65pp and Annex 3 of 192pp., illustr., (1991).
- Fresenius, W., Quentin, K.E. and Schenider, W. : *Water Analysis : A Practical Guide to Physico-Chemical and Microbiological Water Examination and Quality Assurance*, Springer - Verlag, Berlin (1988).
- Goldsborough, L.G. and Robinson, G.G.C. : Seasonal succession of diatom epiphyton on dense mats of *Lemna minor*. *Can. J. Bot.*, **63** : 2332-2339 (1985).
- Kaur, P. : *Evaluations of Water Quality of Lotic and Lentic Water Bodies of Delhi by Analysis of Periphytic Communities on the Roots of Water Hyacinth*. M.Phil. Dissertation, University of Delhi, Delhi, India (1992).
- Kaur, P. and Mehra, N.K. : Evaluation of a new method for qualitative analysis of epiphytic biota on the roots of water hyacinth. *Verh. Int. Ver. Limnol.*, **25** : 2287-2290 (1994).
- Khurshid, S. : Study of physico-chemical and bacteriological characteristics of Yamuna River at Union Territory of Delhi to assess the quality and pollution level. *J. Inst. and Publ. Hlth. Engineers, India* (1989).
- Kiss, K.T. : Changes in trophic conditions in the River Danube at God. *Annal. Univ. Sci. Budapest. Sect. Biol.*, **24-26** : 47-59 (1984).
- Krammer, K. and Lange-Bertalot, H. : *Bacillariophyceae 1. Naviculaceae*. Gustav Fischer-Verlag, Stuttgart (1986).
- Krammer, K. and Lange-Bertalot, H. : *Bacillariophyceae 2. Bacillariaceae, Epithemiaceae, Surirellaceae*. Gustav Fischer, Stuttgart (1988).
- Krammer, K. and Lange-Bertalot, H. : *Bacillariophyceae 3. Centrales, Fragilariaceae, Eunotiaceae*. Gustav Fischer, Stuttgart (1988).
- Kundra, R., Nagpal, J.L., Verma, S.R. and Srivastava, S.K. : Raw water quality of Wazirabad and Okhla reservoir in Delhi. *Ind. J. Environ. Hlth.*, **19** : 329-339 (1977).
- Krebs, E. : *Ecological Methodology*, Harper and Row, New York (1989).
- Lange-Bertalot, H. : Pollution and tolerance of diatoms as a criterion of water quality estimation. *Nova Hedwigia*, **64** : 285-304 (1979).
- Lowe, R.L. : *Environmental Requirements and Pollution Tolerance of Freshwater Diatoms*. National Environmental Research Centre, Office of Research and Development, U.S. Protection Agency, Cincinnati, Ohio (1974).
- Lund, J.W.G. : Studies on *Asterionella formosa* Hass II. Nutrient depletion and the spring maximum. *J. Ecol.*, **38** : 1-35 (1950).
- Mathur, R.P. : Pollution aspects of River Yamuna at Delhi.

- Proc. Symp. on Water Pollution Control, Vol. 3* Central Public Health Engineering Research Institute, Nagpur, India (1965).
- Menon, C.L. : *Ecology of Benthic Fauna in the Floodplain of River Yamuna Affected by Sewage Pollution*. Ph.D. Thesis, Jawahar Lal Nehru University, New Delhi, India (1990).
- Müller, U. : Seasonal development of epiphytic algae on *Phragmites australis* (Cav.) TRIN. ex STEN in a eutrophic lake. *Arch. Hydrobiol.*, **129** : 273-292 (1994).
- Nather Khan, I.S.A. : Assessment of water pollution using diatom community structure and species distribution - A case study in a tropical river basin. *Int. Revue ges. Hydrobiol.*, **75** : 317-338 (1990).
- Nather Khan, I.S.A. : Effect of urban and industrial wastes on species diversity of the diatom community in a tropical river, Malaysia. *Hydrobiologia*, **224** : 175-184 (1991).
- Nwankwo, D.I. : *Seasonal Changes of Phytoplankton of Lagos Lagoon and the Adjacent Sea in Relation to Environmental Factors*. Ph.D. thesis, University of Lagos (1984).
- Nwankwo, D.I. : The phytoplankton of a sewage disposal site in Lagos Lagoon 1. *Nig. J. Biol. Sci.*, **1** : 89-96 (1986).
- Nwankwo, D.I. and Akinsoji, A. : The periphyton algae of a eutrophic creek and their possible use as indicators. *Nig. J. Bot.* **1** : 47-54 (1988).
- Nwankwo, D.I. and Akinsoji, A. : Epiphyte community on water hyacinth *Eichhornia crassipes* (Mart.) SOLMS in coastal waters of south-western Nigeria. *Arch. Hydrobiol.*, **124** : 501-511 (1992).
- Patrick, R. : Use of algae, especially diatoms, in the assessment of water quality. In : *Biological Methods for the Assessment of Water Quality*. J. Cairns and K.I. Dickson (Eds.) ASTM, Philadelphia, 76-95 pp. (1973).
- Patrick, R. and Hohn, M.H. : The diatometer, A method for indicating the conditions of aquatic life. *Division of Refining, Vol 36 (111)* : 332-339 (1956).
- Rai, H. : Hydrobiology of river Yamuna at Okhla, Delhi, India. *Technology of Water*, **6** : 77-98 (1962).
- Rai, H. : Limnological studies on the river Yamuna at Delhi, India. Part I : Relation between chemistry and the state of pollution in the river Yamuna. *Arch. Hydrobiol.*, **73** : 369-393 (1974a).
- Rai, H. : Limnological studies on the river Yamuna at Delhi, India. Part II : The dynamics of potamoplankton populations in the river Yamuna. *Arch. Hydrobiol.*, **73** : 492-517 (1974b).
- Rushforth, S.R. and Brock, J.T. : Attached diatom communities from the lower Turckee River, summer and fall 1986. *Hydrobiologia*, **224** : 49-64 (1991).
- Sabater, S., Sabater, F. and Armengol, J. : Relationships between diatom assemblages and physio-chemical variables in the river Ter (N.E. Spain). *Int. Rev. ges. Hydrobiol.*, **73** : 171-179 (1988).
- Sabater, S., Sabater, F. and Tomas, X. : Water quality and diatom communities in two Catalan rivers (N.E. Spain) *Wat. Res.*, **21** : 901-911 (1987).
- Seki, H., Yonemori, M. and Ito, T. : Effect of rapid oligotrophication by an aquatic treatment pilot plant on the microbial community of a mesotrophic bog. I. Effect on the epiphyton community. *Water, Air and Soil Pollution*, **42** : 387-396 (1988).
- Simm, A.T. : The phytoseston of the Vistula river between Gora Kalwaria and Nowy Dzierżewice in 1982. *Ekol. Pol.*, **33** : 439-453 (1985).
- Sladeckova, A. : Limnological investigation methods for periphyton ("aufwuchs") community. *Bot. Rev.*, **28** : 286-350 (1962).
- Subramanian, V. and Sitasawad, R. : A study on water quality in the river Yamuna around Delhi, India. *Water Quality Bulletin*, **9** : 219-222 (1984).
- Swale, E.M.F. : Phytoplankton in two English rivers. *J. Ecol.*, **57** : 1-23., (1969).
- Tiffany, L.H. and Britton, M.E. : *The Algae of Illinois*. The University of Chicago Press, Chicago (1952).
- Trivedi, R.C., Makhijani, S.D., Ansari, P.M. and Kanyotra, J.S. : Pollution of river Yamuna in Delhi and plan for its improvement. *Proc. Nat. Sem. on Pollution Control and Environment Management, Nagpur, India*. 55-61pp (1985).
- Watanbe, T. : Capblancq, J. and Dauta, A. : Utilisation des bioessays << in situ >> (substrates artificiels) pour caractériser la qualité des eaux de rivière à l'aide du periphyton. *Ann. Limnol.*, **24** : 111-125 (1988).