

## Impact of Nutrition Counselling on Food and Nutrient Intake and Haematological Profile of Rural Pregnant Women

Paramjit Kaur Chawla, Ravnit Kaur and R. Sachdeva

The unfortunate state of nutrition in countries like India is attributed to several factors. Poverty and low purchasing power are no doubt major factors contributing to malnutrition, lack of aware-ness and paucity of information also aggravate the problem. There are three population groups who suffer from the ill effects of malnutrition. These vulnerable segments are pre-school children, expectant and nursing mothers. An expectant and nursing mother needs to be provided adequate nutritional intake for maternal and foetal tissue growth and her own usual maintenance requirements. An acute or chronic deprivation of dietary intake may result in poor pregnancy outcome. In India almost one third of babies born are low birth weight mainly attributable to poor maternal health and nutrition during pregnancy (Sachdev and Choudhury, 1994).

Nutrition counselling is one of the prerequisites for improving the nutritional status of any group. The prenatal patient represents an ideal opportunity for nutrition counselling, since at that time more than any other time, she may be highly motivated to understand and accept advice. Hence the present study was conducted to see the impact of nutrition counselling on food and nutrient intake and haematological profile of pregnant women.

### MATERIALS AND METHODS

The study was conducted on 60 rural pregnant women (18-33 years) belonging to low socio-economic status in three villages of Ludhiana district. The subjects were equally divided into two groups-Experimental (E) and Control (C) group. In the experimental group women were imparted nutrition counselling for a period of five months, starting from fifth month till ninth month (end term). Nutrition counselling was imparted once a month by individual contact during 5<sup>th</sup>, 7<sup>th</sup> and 9<sup>th</sup> month of pregnancy. During 6<sup>th</sup> and 8<sup>th</sup> month of pregnancy nutrition counselling was imparted twice a month by group contacts. The subjects were assembled at *dharamshalla* and were made to sit comfortably. Nutrition

counselling was imparted about the functions of food, balanced diet, nutritional requirements during pregnancy, improving nutritive value of foods. The teaching was carried out in vernacular through lectures, discussion and demonstrations. Visual aids like charts, posters and pamphlets were also used. Nutritional status of pregnant women was assessed during 5<sup>th</sup> and 9<sup>th</sup> month of pregnancy using four methods i.e. dietary survey, anthropometric measurements, clinical examination and biochemical investigation of blood. Information regarding dietary intake of subjects was collected by 24 hour recall method for three consecutive days using standardised containers. The nutrient content of diets consumed by subjects were calculated using 'MSU Nutriguide' software (Song et al., 1992). A series of haematological investigations like haemoglobin (Hb), Packed Cell Volume (PCV) and Mean Corpuscular Haemoglobin (MCH) were carried out during 5<sup>th</sup> and 9<sup>th</sup> month of pregnancy using standard methods. Data was analysed statistically. Comparison of food and nutrient intake of E and C group was made by assessing the significance of the difference between their respective means by applying student 't' test (Gupta and Saini, 1988).

### RESULTS AND DISCUSSION

During 5<sup>th</sup> month of pregnancy, there was no statistically significant difference in the food and nutrient intake of subjects in E and C groups.

#### Food Intake

The mean daily intake of different foods is presented in Table 1. The intake of cereals in E and C group was 270 and 271 g during fifth month and 331 and 296 g during ninth month, respectively. As compared to suggested intakes by ICMR (1981) intake of cereals and pulses was inadequate in both the groups during 5<sup>th</sup> and 9<sup>th</sup> month of pregnancy. Intake of cereals and pulses in E group, during ninth month was significantly ( $P < 0.01$ ) higher as compared to C group. The mean daily intake of pulses in group E and C was 28, 27 g during 5<sup>th</sup> month and 35, 29 g during 9<sup>th</sup>

**Table 1: Average daily food intake of the subjects.**

| Food groups<br>(g)     | Fifth month            |                   | Ninth month            |                   | RDA* | t-value<br>(E & C<br>group) |
|------------------------|------------------------|-------------------|------------------------|-------------------|------|-----------------------------|
|                        | Experimental<br>(n=30) | Control<br>(n=30) | Experimental<br>(n=30) | Control<br>(n=30) |      |                             |
| Cereals                | 270 ± 9.7              | 271 ± 1.15        | 331 ± 15.9             | 296 ± 3.07        | 475  | 6.55**                      |
| Pulses                 | 28 ± 0.31              | 27 ± 0.81         | 35 ± 2.82              | 29 ± 1.18         | 60   | 2.80**                      |
| Green leafy vegetables | 89 ± 0.78              | 87 ± 0.81         | 126 ± 3.2              | 107 ± 1.19        | 100  | 7.68**                      |
| Other vegetables       | 72 ± 0.70              | 71 ± 0.50         | 88 ± 3.2               | 73 ± 1.2          | 40   | 7.18**                      |
| Roots and Tubers       | 49 ± 0.67              | 48 ± 0.57         | 87 ± 0.71              | 65 ± 0.69         | 50   | 3.21**                      |
| Milk & milk products   | 308 ± 1.37             | 312 ± 1.06        | 392 ± 5.6              | 338 ± 3.5         | 250  | 4.80**                      |
| Fats and oils          | 18 ± 0.58              | 16 ± 1.13         | 20 ± 0.50              | 20 ± 0.50         | 30   | 0.70NS                      |
| Sugar and jaggery      | 29 ± 0.63              | 30 ± 0.54         | 32 ± 0.43              | 33 ± 0.81         | 30   | 1.67NS                      |

Values are Mean±SEM

\*\*Significant at (p&lt;0.01) level    NS Non Significant    \* ICMR (1981)

month.

Sachdeva and Mann (1995) and Sachdeva et al. (1996) also reported inadequate intake of cereals and pulses among rural pregnant women in Punjab. Consumption of green leafy vegetables (GLV) in group E and C during 5<sup>th</sup> and 9<sup>th</sup> month was 89, 87g and 126, 107g, respectively. In E group intake of GLV was significantly (P<0.01) more than that of C group. It may be due to nutrition counselling imparted to women during the study. At ninth month the intake of GLV was more as compared to suggested intakes which may be due to more availability of GLV in winter season. Intake of other vegetables and roots and tubers was almost double than the suggested intakes. During ninth month the intake of other vegetables was 88 and 73g and roots and tubers was 87 and 65g in E and C group, respectively which were significantly (P<0.01) higher in E group as compared to C group.

The consumption of milk and milk products was above the RDI in both the groups during 5<sup>th</sup> and 9<sup>th</sup> month. After imparting nutrition counselling the consumption of milk and milk products in E group was significantly (P<0.01) higher than that of C group (392 vs 338g). Sachdeva et al (1996) Chawla and Puri (1996) also have reported similar findings. Mean daily intake of fats and oils was inadequate in both the groups, may be due to poor purchasing power of subjects. However, intake of sugar and jaggery was slightly more than suggested intakes in both the groups.

### Nutrient Intake

Table 2 indicates that the average daily intake of energy in E and C group was 1856, 1760 Kcal

during 5<sup>th</sup> month and 2103, 1882 Kcal during 9<sup>th</sup> month, which was less than the RDA (ICMR, 1987). After nutrition counselling improvement in energy intake was significantly (P<0.01) higher in E group as compared to C group (247 vs 122 Kcal). As compared to RDA (ICMR, 1987) protein intake was 88 percent during 5<sup>th</sup> month in both the groups. However, at the end of gestation, intake was 98 and 91 percent in E and C groups, respectively. Intake of invisible fat in E and C group was 33 and 31g during 5<sup>th</sup> month and 36 and 35 g during 9 month of pregnancy. The daily intake of calcium during 5<sup>th</sup> and 9<sup>th</sup> month in E and C group was 1210, 1192mg and 1472, 1237mg, respectively. Intake was more than the RDA, which was attributed to more intake and easy availability of green leafy vegetables and milk and milk products. Group E had significantly (P<0.01) higher intake as compared to C group at 9<sup>th</sup> month. Similar to present findings, Hira et al (1986) have reported adequate intake of calcium by Punjabi pregnant women. Intake of iron in E group was significantly (P<0.01) more than C group during 9<sup>th</sup> month, but it did not meet the recommendations of 38mg. Similarly Rawtani and Verma (1989) reported adequate intake of iron among the pregnant women of Jodhpur.

Increase in retinol intake of E group was significantly (P<0.01) higher as compared to C group (245 vs 55mg) which could be due to the higher intake of milk and milk products, leafy vegetables and fruits rich in vitamin A. The mean daily intake of thiamine in E and C groups was more than the RDA (154 and 138 percent). During ninth month of pregnancy, the intake of riboflavin and niacin in E group was more than the RDA

**Table 2: Average daily nutrient intake of the subjects**

| Nutrients                | Fifth month            |                   | Ninth month            |                   | RDA*            | t-value<br>(E&C group) |
|--------------------------|------------------------|-------------------|------------------------|-------------------|-----------------|------------------------|
|                          | Experimental<br>(n=30) | Control<br>(n=30) | Experimental<br>(n=30) | Control<br>(n=30) |                 |                        |
| Energy (kcal)            | 1856 ± 44.8            | 1760 ± 45.8       | 2103 ± 33.4            | 1882 ± 23.7       | 2525            | 3.75**                 |
| Protein (g)              | 58 ± 1.90              | 57 ± 1.64         | 64 ± 1.20              | 59 ± 1.85         | 65              | 3.80**                 |
| Total fat (g)            | 51 ± 1.61              | 47 ± 1.54         | 56 ± 1.75              | 55 ± 1.73         | 55 <sup>a</sup> | 1.42NS                 |
| Invisible fat (g)        | 33 ± 0.72              | 31 ± 0.63         | 36 ± 0.78              | 35 ± 0.73         | 30              | 1.33NS                 |
| Calcium (mg)             | 1210 ± 42.1            | 1192 ± 38.5       | 1472 ± 42.7            | 1237 ± 33.0       | 1000            | 6.24**                 |
| Iron (mg)                | 29 ± 0.70              | 28 ± 0.87         | 36 ± 3.08              | 33 ± 1.40         | 38              | 3.50**                 |
| Retinol equivalents (ug) | 819 ± 45.0             | 866 ± 34.8        | 1064 ± 36.9            | 921 ± 24.5        | 600             | 4.65**                 |
| Thiamine (mg)            | 1.7 ± 0.50             | 1.70 ± 0.05       | 2.0 ± 0.13             | 1.8 ± 0.04        | 1.3             | 2.75**                 |
| Riboflavin (mg)          | 1.2 ± 0.70             | 1.3 ± 0.08        | 1.6 ± 0.60             | 1.4 ± 0.09        | 1.5             | 2.86**                 |
| Niacin (mg)              | 13.9 ± 0.50            | 13.8 ± 0.30       | 16.6 ± 0.80            | 14.5 ± 0.69       | 16              | 2.87**                 |
| Folic acid (ug)          | 201 ± 9.50             | 186 ± 14.4        | 292 ± 11.7             | 179 ± 7.40        | 400             | 4.80**                 |
| Ascorbic acid (mg)       | 64 ± 5.20              | 66 ± 3.83         | 99 ± 10.7              | 86 ± 6.2          | 40              | 3.16**                 |

Values are Mean±SEM \*\*Significant at 1% (p<0.01) level  
<sup>a</sup> Calculated (10-30% of calories)

NS Non Significant \*ICMR (1989)

whereas it was less than RDA in C group. The average daily intake of folic acid in E and C groups was 201, 186 and 292, 179mg during 5<sup>th</sup> and 9<sup>th</sup> month, respectively. Although intake of folic acid in E group was significantly (P<0.01) more as compared to C group but it did not meet the recommendations i.e. 400mg. Sachdeva and Mann (1995) and Chawla and Puri (1996) also reported inadequate intake of folic acid among pregnant women in Punjab.

The mean daily intake of ascorbic acid was more than the RDA in both the groups during 5<sup>th</sup> and 9<sup>th</sup> month. The increase observed in E group was significantly (P<0.01) more than in C group (35 vs 20mg) which clearly indicated that nutrition education imparted to pregnant women helped them in better selection of foods rich in vitamin C like lemons, tomatoes, green leafy vegetables and sprouted pulses.

#### Clinical Manifestations of Nutrient Deficiency

The data regarding clinical symptoms of nutrient deficiencies (Table 3) revealed extensive prevalence of anaemia during 5<sup>th</sup> month i.e. 86.7 and 80 percent in E and C group, respectively. The most common clinical symptoms of iron inadequacy i.e. pale conjunctiva and pale skin were more evident during 9<sup>th</sup> month in C group (83.3 and 86.6%) as compared to E group (10 and 26.6 %).

At 5<sup>th</sup> month feeling of fatigue was complained by 43.3 percent subjects in each group, at 9<sup>th</sup> month it reduced to 6.7 percent in E group and increased to 60 percent in C group. Symptoms

of anaemia and other nutritional deficiencies decreased during 9<sup>th</sup> month in E group which could be due to impact of nutrition counselling.

#### Biochemical Assessment

Haematological profile of subjects (Table 4) indicated that Hb level was 9.7 and 9.0g/ dl in E and C group, respectively. During 9<sup>th</sup> month Hb level increased in E group (11.4g/dl) and decreased in C group (9.2g/dl). The mean PCV value was 28.9, 28.1 and 33.3, 29.1 percent during 5<sup>th</sup> and 9<sup>th</sup> month in E and C group, respectively. MCHC values during 5<sup>th</sup> month were 33.56 and 33.2 percent in E and C group whereas corresponding values during 9<sup>th</sup> month were 34.23, 33.94 percent which were within normal range. After nutrition counselling Hb and PCV values were significantly (P<0.01) higher in E group as compared to C group.

Although nutrition counselling resulted in better food intake and regular use of ferrous sulphate tablets by the subjects in E group, yet 50 percent subjects had Hb level <11g/dl. However in control group only 16.6 percent subjects had Hb level >11g/dl. Lower Hb values in the present study coincides with the findings reported by Sachdeva et al. (1996).

The present investigation showed an improvement in food and nutrient intake among the subjects at the end of gestation due to increased requirements of pregnancy. Improvement was found to be significantly higher in E group as compared to C group for most of the nutrients but intake was below the recom-

**Table 3: Nutritional deficiency symptoms during pregnancy**

| Symptoms                 | Fifth month            |      |                   |      | Ninth month            |      |                   |      |
|--------------------------|------------------------|------|-------------------|------|------------------------|------|-------------------|------|
|                          | Experimental<br>(n=30) |      | Control<br>(n=30) |      | Experimental<br>(n=30) |      | Control<br>(n=30) |      |
|                          | N                      | %    | N                 | %    | N                      | %    | N                 | %    |
| a) Anaemia               |                        |      |                   |      |                        |      |                   |      |
| i) Pale conjunctiva      | 12                     | 40.0 | 14                | 46.7 | 3                      | 10.0 | 25                | 83.3 |
| ii) Pale skin            | 26                     | 86.7 | 24                | 80.0 | 8                      | 26.6 | 26                | 86.6 |
| iii) Flat nails          |                        |      |                   |      |                        |      |                   |      |
| Fingers                  | 4                      | 13.3 | 3                 | 10.0 | 3                      | 10.0 | 3                 | 10.0 |
| Toes                     | 4                      | 13.3 | 3                 | 10.0 | 3                      | 10.0 | 0                 | 0    |
| iv) Koilonychia          |                        |      |                   |      |                        |      |                   |      |
| Fingers                  | 3                      | 10.0 | 3                 | 10.0 | 0                      | 0    | 1                 | 3.3  |
| Toes                     | 5                      | 16.6 | 4                 | 13.3 | 1                      | 3.3  | 0                 | 0    |
| b) Angular stomatitis    | 4                      | 13.3 | 5                 | 16.6 | 3                      | 10.0 | 4                 | 13.3 |
| c) Cheilosis             | 4                      | 13.3 | 9                 | 30.0 | 3                      | 10.0 | 6                 | 20.0 |
| d) Bleeding gums         | 3                      | 10.0 | 8                 | 26.7 | 2                      | 6.7  | 7                 | 23.3 |
| e) Incidence of headache | 7                      | 23.3 | 9                 | 30.0 | 5                      | 16.6 | 6                 | 20.0 |
| f) Feeling of fatigue    | 13                     | 43.3 | 13                | 43.3 | 2                      | 6.7  | 18                | 60.0 |
| g) Feeling of lethargy   | 11                     | 36.7 | 14                | 46.6 | 8                      | 26.7 | 10                | 33.3 |

**Table 4: Hematological profile of pregnant women**

| Blood Parameters              | Fifth month            |                   | Ninth month            |                   | Normal value | t-value<br>(E & C gp) |
|-------------------------------|------------------------|-------------------|------------------------|-------------------|--------------|-----------------------|
|                               | Experimental<br>(n=30) | Control<br>(n=30) | Experimental<br>(n=30) | Control<br>(n=30) |              |                       |
| Haemoglobin (g/dl)            | 9.7 ± 0.16             | 9.0 ± 0.07        | 11.4 ± 0.17            | 9.2 ± 0.27        | 11.0         | 2.68**                |
| Haematocrit (PCV)%            | 28.9 ± 0.75            | 28.1 ± 0.28       | 33.3 ± 0.42            | 27.1 ± 0.27       | 32.36        | 3.74**                |
| Mean corpuscular              | 33.56 ± 0.63           | 33.02 ± 0.30      | 34.23 ± 0.58           | 33.94 ± 0.60      | 30.35        | 2.17**                |
| Haemoglobin Concentration (%) |                        |                   |                        |                   |              |                       |

Values are Mean ± SEM

\*\* Significant at 1% (p&lt;0.01) level

\* Significant at 5% (p&lt;0.05) level

mendations for some nutrients despite imparting nutrition education.

Inadequate nutrient intake might be due to poverty, low purchasing power, lack of knowledge, ignorance, illiteracy and less availability of foods. Inadequate nutrient intake showed the wide prevalence of iron deficiency anaemia among the subjects. The study shows that in low socio-economic group, nutrition education is likely to result in better nutrient intake and haematological profile. However, there is need to impart concerted nutrition education to pregnant women for a longer duration starting earlier in pregnancy to improve their nutritional status to the desired level.

**KEYWORDS** Nutrition Counselling. Pregnant Women. Food Consumption. Nutrient Intake

**ABSTRACT** Sixty pregnant women belonging to low socio-economic status during 5<sup>th</sup> month of pregnancy were selected from three villages of Ludhiana district. They were assessed for their nutritional status and divided into two groups. Group

I (30 women) was given nutrition education for five months about nutritional needs during pregnancy, served as Experimental (E) group and other 30 women served as control (C) group. Dietary survey revealed that the intake of cereals, pulses, green leafy vegetables and fats and oils was less than Recommended Dietary Intakes (ICMR 1980). However, the intake of milk and milk products, other vegetables, roots and tubers and sugar and jaggery was more than the recommended intakes. Average energy intake was 1856, 1760 Kcal and 2103, 1882 Kcal during 5<sup>th</sup> and 9<sup>th</sup> month by the subjects belonging to E and C group, which was less than the RDA. However, intake of protein was almost adequate. Nutrient intake during 9<sup>th</sup> month improved in both the groups due to increased requirements of pregnancy. In E group intake of all the nutrients was significantly higher as compared to C group. This may be due to more consumption of pulses, milk and milk products and leafy vegetables and also due to adoption of desirable nutritional practices like use of sprouted grains and other nutritious foods. Thus there is wide scope of imparting nutrition counselling to pregnant women to improve their nutritional status and that of newborns.

## REFERENCES

Chawla, P.K. and Puri, R.: Impact of nutrition education on

- food and nutrient intake of pregnant women. *Ind. J. Maternal Child Health*, **7** (1): 11-15 (1996).
- Gopalan, C., Shastri, B. V. and Balasubramanian, S. C.: *Nutritive Value of Indian Foods*. Indian Council of Medical Research, National Institute of Nutrition, Hyderabad, India, pp. 94 (1987).
- Gupta, J. P. and Saini, S. S.: *Introduction to Statistical Methods*. Second Edition, Kalyani Publishers, New Delhi, pp. 137 -142 (1987).
- Hira, C. K., Pathak, V. and Murgai, V.: Nutritional status of pregnant Punjabi women in relation to education and economic level. *Ecol. Food Nutr.*, **19**: 147-153 (1986).
- Rawtani, L. and Verma, M.: A study of nutritional status and food practices of the pregnant and lactating women residing in selected desert areas of Jodhpur. *Ind. J. Nutr. Dietet.*, **26**: 304-310 (1989).
- Sachdev, H. P. S. and Choudhury: *Nutrition in Children-Developing Country Concerns*. Published by Conveners, Maulana Azad Medical College, New Delhi (1994).
- Sachdeva, R. and Mann, S. K.: Efficacy of nutrition counselling and medical supervision on the nutriture of rural pregnant women. *J. Res.*, **31**: 483-497 (1995).
- Sachdeva, R., Sukhwinder, K. and Sangha, J. K.: Efficacy of antenatal nutrition counselling on the nutriture and haematological profile of rural pregnant Punjabi women. *Applied Nutr.*, **21**: 18-24 (1996).
- Song, W. D., Roerr, S., Bnond, J., Kalarala, M., Mann, S., Sehgal, S., Pande, U., Singh, I., Mehta, U. and Rohinidevi, P.: *Nutriguide: Asian Indian Foods Nutritional Analysis Computer Programme*. Michigan State University, USA (1992).

**Authors' Address:** Paramjit Kaur Chawla , Ravnit Kaur and R. Sachdeva, Department of Food and Nutrition, Punjab Agricultural University, Ludhiana 141 001, Punjab, India