

Phenotypic and Genetic Characterization of Sorbitol-Fermenting *Escherichia coli* O157: H7 Isolated from Retail Beef and Mince Beef

Kaemogetso Makhubalo¹, Madira Manganyi², Ajay Kumar³,
Moses Mbewe^{1,2} and Collins N. Ateba^{3,4*}

¹Center for Animal Health Studies, North West University, Mafikeng Campus,
Private Bag X2046, Mmabatho 2735, South Africa

²Department of Water and Sanitation, University of Limpopo, Turfloop Campus,
Private Bag X1106, Sovenga 0727, Polokwane, South Africa

³Department of Biological Sciences, North West University, Mafikeng Campus,
Private Bag X2046, Mmabatho 2735, South Africa

⁴Food Security and Safety Niche Area, Faculty of Agriculture, Science and Technology,
North-West University, Mmabatho, Mafikeng 2735, South Africa

KEYWORDS *E. coli* O157:H7. Shiga-toxin. 16S rRNA Gene. *rfbE*_{O157} Gene. *fliC*_{H7} Gene

ABSTRACT *Escherichia coli* occur as normal flora in the gastrointestinal tract of humans and animals. However, shiga-toxin producing *E. coli* and most especially serotype O157:H7 are a major group of food-borne pathogens. The consumption of undercooked contaminated food products of animal origin particularly beef has identified as a potential source for the transmission of these pathogens to humans. The aim of this paper was to determine the level of *E. coli* O157:H7 contamination in beef and mince beef samples obtained from some supermarkets and evaluate the virulence gene profiles of the isolates. A total of 128 beef and mince beef samples were collected from selected supermarkets and butcheries in the North West Province of South Africa. Samples were analysed for heterotrophic bacterial counts on PCA and potential *E. coli* O157:H7 counts on SMAC. Preliminary (oxidase test, TSI agar test) and confirmatory biochemical (API 20E, O157 and H7 serological assays, *rfbE*_{O157} and *fliC*_{H7} PCR analysis) tests were used to identify *E. coli* O157:H7 isolates while the virulence profiles of the isolates were determined through amplification of the *stx1*, *stx2*, *eaeA* and *hlyA* virulence genes. Despite the fact that all the beef samples were contaminated with heterotrophic bacteria none of those from Ventersdorp and Rustenburg were contaminated with potential *E. coli* O157:H7. Beef samples from Swaartruggens had highest heterotrophic microbial count (5.3x10⁶ CFU/mL) while those from Garankwa had the lowest level of contamination (1.3x10⁴ CFU/mL). All (100 percent) the mince beef samples collected from the different sampling stations were contaminated with both heterotrophic and potential *E. coli* O157 bacteria. Mince beef samples from Potchefstroom, Rustenburg, Klerksdorp, Ventersdorp, Zeerust, Mafikeng, Swaartruggens, Vryburg, Bloemhof and Taung had very high levels (1.8x10³ - 4.1x10⁶) of potential *E. coli* O157:H7 contamination. All the isolates from beef except for those isolated from Rustenburg and Ventersdorp as well as those from mince beef were Gram-negative rods. A large proportion (71.4 percent to 100 percent) of the isolates from both beef and mince beef were oxidase negative. Large proportions (92.9 percent to 100 percent) of isolates from beef as well as small proportions (14.3 percent to 42.9 percent) of those from mince beef fermented the sugars in the TSI medium. Small proportions (7.1 percent to 42.9 percent) of the isolates from both beef and mince beef produced gas and hydrogen sulphide from the TSI agar test. A total of 39 and 38 *E. coli* isolates were obtained from beef and mince beef samples respectively while a large proportion 27 (70.1 percent) of these isolates were positively identified as *E. coli* O157:H7 using a serological assay. Only 8 *E. coli* O157:H7 isolates were positively identified in the beef samples while large proportions (23) of the isolates from mince beef were positive for the *fliC*_{H7} and *rfbE*_{O157} gene fragments. Among the isolates from beef, the *stx2* gene was present in all the 8 isolates while only 3 isolates possessed the *stx1* gene. Similarly, the *stx2* gene was frequently (13) detected among the isolates from mince beef when compared to the *stx1* gene. Putative virulence genes *hlyA* and *eaeA* were detected in 12 and 6 isolates respectively from mince beef. Six isolates from beef possessed both the *hlyA* and *eaeA* gene fragments. The findings of this study revealed that beef and mince beef samples obtained from some supermarkets in the area may pose severe health risks to consumers if they are not properly cooked before consumption.

Address for correspondence:

Collins Njie Ateba
Department of Biological Sciences,
North West University, Mafikeng Campus,
Private Bag X2046,
Mmabatho 2735,
South Africa
Telephone: (+27) 18 389 2247,
E-mail: 16528026@nwu.ac.za

INTRODUCTION

Escherichia coli occur as normal flora in the gastrointestinal tract of humans and animals (Harakeh et al. 2005; Ojo et al. 2010; Ateba and Mbewe 2011; Zheng et al. 2012; Díaz-Sánchez et al. 2013; Abraham et al. 2015; Tamang et al. 2015; Vijay et al. 2015; Liu et al. 2016). However, shiga-

toxin producing *E. coli* and most especially serotype O157:H7 are a major group of food-borne pathogens (Harakeh et al. 2005; Rangel et al. 2005; Seto et al. 2007; Amani et al. 2015; Kargar and Homayoon 2015). Despite the fact that *E. coli* O157:H7 may be present in the gastrointestinal tract of healthy domestic animals such as cattle, sheep and goats without any clinical symptoms (Müller et al. 2002; Cobbold et al. 2000) the pathogen has been associated with a variety of diseases in humans ranging from non-bloody to severe bloody diarrhoea, haemorrhagic colitis, haemolytic uraemic syndrome and thrombotic thrombocytopenic purpura which may be very lethal in immuno-compromised patients (Banatvala et al. 2001; Momba et al. 2008). These infections are usually the cause of renal failure in infected subjects (Weir 2000). Virulent *E. coli* O157:H7 strains harbour a variety of virulence gene determinants including *Stx*₁, *Stx*₂ and variants of *Stx*₂, *hlyA* and *eaeA* gene (Paton and Paton 1998; Law 2000; Schouten et al. 2005). These toxins invade the cells of the colon, epithelia and the endothelium of the human GIT tract resulting in disease (Paton and Paton 1998). *E. coli* O157:H7 infections usually result from the consumption of undercooked contaminated food products of animals origin and meat particularly beef has identified as a potential source for the transmission of these pathogens to humans (Meichtri et al. 2004; Stampi et al. 2004; Ojo et al. 2010; Ateba and Mbewe 2011; Gordillo et al. 2014; Ahmed and Shimamoto 2015; Dong et al. 2015; Hessain et al. 2015; Xu et al. 2016). In most developing countries, proper hygiene practices are not always strictly implemented during the handling and processing (Harakeh et al. 2005). Against this background meat, the equipment used for cutting meat and the environment in which the meat is stored may become contaminated with pathogenic bacteria strains including *E. coli* O157:H7 during slaughter (Elder et al. 2000; Blanco et al. 2003; Conedera et al. 2004; Magwira et al. 2005). Outbreaks of *Escherichia coli* O157:H7 infections through consumption of contaminated foods including undercooked meat products have brought a great safety concern (Gordillo et al. 2014; Ahmed and Shimamoto 2015; Dong et al. 2015; Hessain et al. 2015; Xu et al. 2016). The objectives of this study was to determine the level of *E. coli* O157:H7 on beef and mince beef and to evaluate the virulence gene profiles of the isolates.

MATERIAL AND METHODS

Sampling and Isolation of Bacteria

A total of 128 beef and mince beef samples were collected from selected supermarkets and butcheries in the North West Province of South Africa (Table 1). Fresh beef and minced beef portions were aseptically collected and properly packed in sterile Whirlpak bags. The samples were properly labeled and transported on ice to the laboratory for analysis. One gram from each sample was washed in 5 ml of 2 percent (w/v) peptone water. Ten-fold serial dilutions were prepared and an aliquot of 100 µL from each dilution was spread-plated on Sorbitol MacConkey agar (SMAC) agar and Plate Count agar (PCA) medium. The plates were incubated aerobically at 37°C for 24 hours. Colonies on PCA and SMAC agar plates were counted and results were re-

Table 1: Areas, sources and types of samples collected

Areas of collection	Source samples	Nature of samples	Number of samples collected
Zeerust	Bovine	Meat	4
Lichtenburg	Bovine	Mince meat	4
	Bovine	Meat	4
Koster	Bovine	Mince meat	4
	Bovine	Meat	4
Mafikeng	Bovine	Mince meat	4
	Bovine	Meat	4
Swaartruggens	Bovine	Mince meat	4
	Bovine	Meat	4
Delareyville	Bovine	Mince meat	4
	Bovine	Meat	4
Swarzrenerke	Bovine	Mince meat	4
	Bovine	Meat	4
Rustenburg	Bovine	Mince meat	4
	Bovine	Meat	4
Vryburg	Bovine	Mince meat	4
	Bovine	Meat	4
Taung	Bovine	Mince meat	4
	Bovine	Meat	4
Garankwa	Bovine	Mince meat	4
	Bovine	Meat	4
Brits	Bovine	Mince meat	4
	Bovine	Meat	4
Potchefstroom	Bovine	Mince meat	4
	Bovine	Meat	4
Ventersdorp	Bovine	Mince meat	4
	Bovine	Meat	4
Bloemhof	Bovine	Mince meat	4
	Bovine	Meat	4
Klerksdorp	Bovine	Mince meat	4
	Bovine	Meat	4

corded. Isolates on SMAC were later sub-cultured on SMAC agar and plates were incubated aerobically at 37°C for 24 hours. The isolates on SMAC were stored at room temperature and used for bacteria identification.

Bacteria Identification

Presumptive *E. coli* isolates were identified based on the following criteria:

Cellular Morphology

Isolates were Gram stained using standard techniques (Cruikshank et al. 1975). *E. coli* species are Gram negative rods, thus all isolates that appeared pink were retained and subjected to primary and secondary biochemical identification tests.

Oxidase Test

The oxidase test was performed on all *E. coli* isolates using the TestOxidase™ reagent (PL.390) in accordance with the manufacturer's protocol (Mast Diagnostics, Neston, Wirral, U.K.).

Triple Sugar Iron (TSI) Agar Test

Triple sugar iron agar (Biolab, Merck – South Africa) was used to assay the *E. coli* content against three sugars (glucose, sucrose and lactose) that are present at concentrations 0.1 percent, 1.0 percent and 1.0 percent respectively (Forbes and Weissfeld 1998). Results were interpreted based on standard protocols (Prescott 2002).

Simmons Citrate Agar Test

Simmons citrate agar was used to determine the ability of the isolates to utilize citrate and ammonium ions as the sole source of carbon and nitrogen respectively, (Simmons 1926).

All isolates that satisfied this preliminary identification criterion were subjected to confirmatory biochemical tests.

Confirmatory Biochemical Identification Tests for *E. coli* isolates

Analytical Profile Index (API 20E) Test

API 20E was used for identification of *E. coli* according to the manufacturer's specifications

(BioMérieux, Marcy l'Etoile, France). Indices were generated and the identities of the isolates were determined using the API web software.

Serotyping

Representative *E. coli* isolates from each sample were further analysed for characteristics of *E. coli* O157:H7 using the slide agglutination test with *E. coli* O157 and H7 specific antisera obtained from Mast Diagnostics, U.K.). Agglutination that was strong and clearly visible within one minute was recorded as positive results.

Molecular characterization of *E. coli* O157:H7 Isolates

Extraction of Genomic DNA

Genomic DNA was extracted from bacteria using a modification of the hot (65°C) CTAB (cetyltrimethylammonium bromide)–PVP (Polyvinylpyrrolidone) DNA extraction procedure (Dolye and Dolye 1990).

The quality and purity of DNA was checked using the spectrophotometric method as described earlier by Sambrook et al. (1989).

Polymerase Chain Reaction (PCR) for the Amplification of *rfbE*_{O157:H7}, 16S rRNA and *fliC*_{H7} Gene Fragments

The PCR protocols were used to amplify bacterial *rfbE*_{O157:H7}, 16S rRNA and *fliC*_{H7} gene fragments as described earlier by Paton and Paton (1998), Muyzer et al. (1995) and Gannon et al. (1997), respectively. Oligonucleotide primers sequences that were used in the study are shown in Table 2. The PCR reactions were performed to amplify *rfbE*_{O157:H7} gene by using a DYAD DNA engine model (Model - PTC- 220 DYAD™ DNA Engine Thermocycler with the following conditions: initial denaturation 95°C for 5 minutes, followed by 34 cycles at 95°C for 30 seconds, 65°C for 30 seconds, 72°C for 30 seconds cycles, and finally elongation at 72°C for 3 minutes. For the amplification of 16S rRNA gene amplification following thermal conditions were used: initial denaturation at 94°C for 5 minutes followed by 31 cycles consisting of denaturation at 94°C for 1 minute, annealing at 67°C for 40 seconds and extension at 72°C for 45 seconds, followed by final extension at 72°C for 2 minutes. The *fliC*_{H7} gene segment was amplified by using the fol-

Table 2: Oligonucleotide primers sequences that were used in the study

Primer	Sequence (5' – 3')	Specificity	Amplicon size BP
O157F ^a	CGGACATCCATGTGATATGG	<i>rfbE</i> _{O157:H7}	259
O157R ^a	TTGCCTATGTACAGCTAATCC		
GM5F ^b	CCGTCAATTCCTTTGAGTTT	16S bacterial ribosomal genes	550
907R ^b	CCGTCAATTCCTTTGAGTTT		
<i>fliCh7</i> F ^c	GCGGCTGTCGAGTTCTATCGAGC	H7 gene	625
<i>fliCh7</i> R ^c	CAACGGTGACTTTATCGCCATTCC		

^aPaton and Paton (1998), ^bMuyzer (et al.) 1995, ^cGannon (et al.) 1997

lowing PCR conditions: initial denaturation at 96°C for 3 minutes, followed by 26 cycles at 96°C for 10 seconds, 51°C for 7 seconds and 62°C for 15 seconds; followed by final extension at 62°C for 4 minutes. The amplified PCR amplicons products were stored at 4°C until used.

Agarose Gel Electrophoresis

PCR products were separated by electrophoresis on a 1 percent (w/v) agarose gel using a horizontal Pharmacia biotech equipment system (model Hoefer HE 99X) for 2 hours at 80V using 1 X TAE buffer. A 100 bp DNA molecular weight marker was included in each run and was used to compare the sizes of the amplified fragments. Gels were stained in ethidium bromide (0.001 µg/ml) and visualized under UV light (Sambrook et al. 1989). A Gene Genius Bio Imaging System (Syngene, Synoptics.UK) was used to capture images using the gene snap (version). The images were analyzed using the gene tools (version) software (Synegene, Synoptics, UK).

RESULTS

Detection of Potential *E. coli* Isolates in the Samples

A total of 128 samples of raw beef and mince were examined for the presence of *E. coli* O157:H7 and results revealed that large proportion (60%) of the samples was contaminated with *E. coli* O157:H7 based on morphological characteristics. This included 87.5 percent of beef samples analysed in the study. Beef samples from Garankwa were the most contaminated with 31.8 percent potential *E. coli* O157:H7 contamination when compared to those from Swaartruggens that had the lowest contamination levels (Table 3). On the contrary, none of the samples from Ventersdorp and Rustenburg were contaminat-

ed with potential *E. coli* O157:H7 strains based on colonial morphologies.

Table 3: Total heterotrophic bacterial counts and potential *E. coli* isolates obtained from beef samples

Areas of collection	Colony forming units on plate count agar	Colony forming units on Sorbitol Macconkey Agar
Brits	9.9x 10 ⁵	8.5 x 10 ²
Mafikeng	1.1x10 ⁵	4.7 x10 ²
Garankwa	1.3x10 ⁴	0.4 x10 ⁴
Koster	1.2x10 ⁵	0.3 x10 ³
Potchefstroom	0.9x10 ⁶	0.1 x10 ⁴
Klerksdorp	1.8x10 ⁵	3.3 x10 ³
Ventersdorp	2.9x10 ⁴	0
Zeerust	1.5x10 ⁴	0.2 x 10 ³
Lichtenburg	0.9x10 ⁶	6.5 x10
Swaartruggens	5.3x10 ⁶	3.7 x10 ²
Vryburg	1.4x10 ⁶	6.9 x10 ³
Delaryville	2.8x10 ⁶	6.3 x10 ³
Swazrenecke	5.3x10 ⁵	1.5 x10 ³
Bloemhof	1.4x10 ⁴	0.6 x10 ³
Taung	1.3x10 ⁵	6.4 x10 ³
Rustenburg	1.7x10 ⁶	0

Determination of the Level of Contamination with Heterotrophic Bacteria and Potential *E. coli* isolates in Samples

The level of microbial contamination with potential *E. coli* O157:H7 isolates with respect to heterotrophic bacteria was determined for both beef and mince beef samples and results are shown in Tables 3 and 4 respectively. Despite the fact that all the beef samples were contaminated with heterotrophic bacteria none of those from Ventersdorp and Rustenburg were contaminated with potential *E. coli* O157:H7 (Table 3). Beef samples from Swaartruggens had highest heterotrophic microbial count (5.3x10⁶ CFU/ml) while those from Garankwa had the lowest level of contamination (1.3x10⁴ CFU/ml). In

addition, samples from Vryburg, Delareyville and Taung had very high levels of contamination with potential *E. coli* O157 contaminants (Table 3). All (100%) the mince beef samples collected from the different sampling stations were contaminated with both heterotrophic and potential *E. coli* O157 bacteria (Table 4). Samples from Potchefstroom, Rustenburg, Klerksdorp, Ventersdorp, Zeerust, Mafikeng, Swaartruggens, Vryburg, Bloemhof and Taung had very high levels (1.8×10^3 - 4.1×10^6) of potential *E. coli* O157:H7 contamination. This was a cause for

concern and it is suggested that these samples may facilitate the transmission of *E. coli* O157:H7 strains to humans especially if the food products are consumed undercooked.

Preliminary Identification of *E. coli* Isolates from Beef and Mince Beef Samples

A total of 448 presumptive *E. coli* O157:H7 isolates that were different based on colonial morphologies were obtained from samples collected from the different locations and these isolates were subjected to preliminary identification tests that are specific for *E. coli* (Gram staining, oxidase test, TSI agar test and Simmons citrate agar test). The makeup included 224 isolates from beef and mince beef samples. Results obtained for isolates obtained from beef and mince beef samples are shown in Tables 5 and 6 respectively. All the isolates from beef were Gram-positive rods except for those isolated from Rustenburg and Ventersdorp. However, all the isolates from mince beef were Gram-negative rods. A large proportion (71.4% to 100%) of the isolates from both beef and mince beef were oxidase negative. Despite the fact that a large proportion (85.7% to 92.9%) of the isolates from beef in Vryburg and Lichtenburg did not ferment the sugars from the TSI medium, large proportions (92.9% to 100%) of those from the other stations fermented these sugars. However, none of the isolates from Rustenburg and Ventersdorp as well as only a small proportion (7.1% to 5 (35.7%) produced gas from the TSI medium. In addition, only 33 of the 224 isolates from beef

Table 4: Total heterotrophic bacterial counts and potential *E. coli* isolates obtained from mince beef samples

Area of collection	No. of colony farming units in plate count Agar	No. of colony farming units in Sarbitol Macconkey Agar
Garankwa	2.03×10^3	2×10^2
Brits	2.52×10^5	3.68×10^2
Koster	8.3×10^5	1.2×10^3
Potchefstroom	9.1×10^6	4.1×10^6
Klerksdorp	3.4×10^4	8.5×10^3
Ventersdorp	1.38×10^5	8.5×10^3
Zeerust	2.2×10^6	2.5×10^4
Lichtenburg	1×10^4	5.5×10^2
Mafikeng	1.9×10^5	1.8×10^3
Swaartruggens	19.5×10^4	5.5×10^4
Vryburg	9×10^4	6.5×10^3
Delareyville	1.22×10^5	2.3×10^3
Swazrenecke	6.37×10^5	8.7×10^2
Bloemhof	6.1×10^4	9.5×10^4
Taung	1.3×10^6	8.5×10^4
Rustenburg	1.4×10^6	1.2×10^4

Table 5: Proportion of isolates from beef samples that were positive for preliminary biochemical tests

Areas of collection	Gram staining (Negative rod)	Oxidase test (Negative)	Simmons Citrate agar test (Negative)	Colour change or acid production	Gas production	Hydrogen sulphide production
Brits	14 (100%)	13 (92.9%)	6 (42.9%)	1 (7.1%)	1 (7.1%)	1 (7.1%)
Garankwa	14 (100%)	14 (100%)	1 (7.1%)	2 (14.3%)	3 (21.4%)	0
Rustenburg	0	0	0	0	0	7 (50%)
Vryburg	14 (100%)	13 (92.9%)	12 (85.7%)	1 (7.1%)	7 (50%)	0
Koster	14 (100%)	13 (92.9%)	3 (21.4%)	3 (21.4%)	4 (28.6%)	4 (28.6%)
Bloemhof	14 (100%)	12 (85.7%)	9 (64.3%)	3 (21.4%)	6 (42.9%)	2 (14.3%)
Taung	14 (100%)	4 (28.6%)	4 (28.6%)	1 (7.1%)	0	1 (7.1%)
Swaartruggens	14 (100%)	11 (78.6%)	0	1 (7.1%)	1 (7.1%)	1 (7.1%)
Klerksdorp	14 (100%)	12 (85.7%)	3 (21.4%)	3 (21.4%)	1 (7.1%)	0
Ventersdorp	0	0	0	0	0	4 (28.6%)
Zeerust	14 (100%)	6 (42.9%)	16.1	3 (21.4%)	2 (14.3%)	2 (14.3%)
Delareyville	14 (100%)	11 (78.6%)	6 (42.9%)	1 (7.1%)	0	8 (57.1%)
Swaartruggens	14 (100%)	10 (71.4%)	0	3 (21.4%)	2 (14.3%)	0
Lichtenburg	14 (100%)	12 (85.7%)	13 (92.9%)	3 (21.4%)	1 (7.1%)	0
Potchefstroom	14 (100%)	13 (92.9%)	3 (21.4%)	5 (35.7%)	4 (28.6%)	0
Mafikeng	14 (100%)	7 (50%)	3 (21.4%)	1 (7.1%)	7 (50%)	3 (21.4%)

produced hydrogen sulphide gas (Table 5). Only a small proportion (14.3% to 42.9%) of the isolates from mince beef as well as none of those from Ventersdorp produced acid from breakdown of carbohydrates in the TSI medium. Similarly, none of the isolates from Taung, Ventersdorp, Delareyville and Rustenburg including a small proportion (7.1% to 42.9%) of those from mince beef in the other sample stations produced gas (Table 6). Interestingly, only 30 of the 224 isolates from mince beef produced hydrogen sulphide gas which is generally not a biochemical characteristic of *E. coli* strains.

Serological and API 20E Tests for Isolates from Beef and Mince Beef Samples

None of the beef samples from Rustenburg, Koster, Taung, Swaartruggens, Ventersdorp and Zeerust were positive for *E. coli* contamination using the API 20E assay. Despite this, 39 *E. coli* isolates were obtained from beef in the study. Similarly, 38 *E. coli* isolates were also obtained from mince beef samples. Among these isolates only small proportion 11 (28.2%) of the isolates from beef were positively identified as *E. coli* O157:H7. On the contrary, a large proportion 27 (70.1%) of the *E. coli* isolates were positive for *E. coli* O157 and H7 serological assays (Table 7). Interestingly *E. coli* O157:H7 was not detected in both beef and mince beef samples from Taung.

Genotypic Characterization and Virulence Genes Detection in *E. coli* O157:H7

In order to avoid bias all the 448 isolates were subjected to a universal 16S rRNA PCR assay as well as *E. coli* O157:H7 specific PCR designed to amplify the *fliC_{H7}* and *rfb_{O157}* gene fragments. A further task was to assess the virulence profiles of the isolates through amplification of the *stx1* and *stx2* as well as putative virulence factors enterohemolysin (*hlyA*) and intimin (*eaeA*). Detailed results are shown in Tables 8 and 9 for beef and mince beef isolates respectively. Despite the fact that only 8 *E. coli* O157:H7 isolates were positively identified in the beef samples a large proportion (23) of the isolates from mince beef was positive for the *fliC_{H7}* and *rfb_{O157}* gene fragments. Among the isolates from beef, the *stx2* gene was present in all the 8 isolates while only 3 isolates possessed the *stx1* gene. Similarly, the *stx2* gene was frequently (13) detected among the isolates from mince beef when compared to the *stx1* gene. Putative virulence genes *hlyA* and *eaeA* were detected in 12 and 6 respectively of the mince beef isolates. Among the isolates from beef 6 isolates possessed both the *hlyA* and *eaeA* gene fragments. It was also identified that beef samples from Mafikeng and mince beef samples from Brits had the highest level of contamination with *E. coli* O157:H7 (Tables 8 and 9). Despite the fact that *E. coli* O157:H7 was not detected in both beef samples from Koster, Bloemhof, Taung, Swaartruggens, Zeerust,

Table 6: Proportion of isolates from mince beef samples that were positive for preliminary biochemical tests

Areas of collection	Gram staining (Negative rod)	Oxidase test (Negative)	Simmons Citrate agar test (Negative)	Colour change or acid production	Gas production	Hydrogen Sulphide production
Brits	14 (100%)	8 (57.1%)	3 (21.4%)	2 (14.3%)	3 (21.4%)	0
Garankwa	14 (100%)	13 (92.9%)	1 (7.1%)	2 (14.3%)	2 (14.3%)	0
Mafikeng	14 (100%)	2 (14.3%)	0	2 (14.3%)	1 (7.1%)	6 (42.9%)
Vryburg	14 (100%)	10 (71.4%)	3 (21.4%)	5 (35.7%)	5 (35.7%)	0
Koster	14 (100%)	6 (42.9%)	5 (35.7%)	2 (14.3%)	3 (21.4%)	0
Bloemhof	14 (100%)	5 (35.7%)	3 (21.4%)	6 (42.9%)	6 (42.9%)	0
Taung	14 (100%)	6 (42.9%)	3 (21.4%)	2 (14.3%)	0	1 (7.1%)
Swaartruggens	14 (100%)	8 (57.1%)	3 (21.4%)	2 (14.3%)	1 (7.1%)	1 (7.1%)
Klerksdorp	14 (100%)	10 (71.4%)	13 (92.9%)	2 (14.3%)	1 (7.1%)	0
Zeerust	14 (100%)	4 (28.6%)	3 (21.4%)	2 (14.3%)	2 (14.3%)	2 (14.3%)
Delaryville	14 (100%)	5 (35.7%)	0	2 (14.3%)	0	9 (64.3%)
Swarzrenerke	14 (100%)	13 (92.9%)	3 (21.4%)	2 (14.3%)	2 (14.3%)	0
Lichtenburg	14 (100%)	13 (92.9%)	2 (14.3%)	2 (14.3%)	1 (7.1%)	0
Potchefstroom	14 (100%)	13 (92.9%)	12 (85.7%)	5 (35.7%)	4 (28.6%)	0
Rustenburg	14 (100%)	14 (100%)	4 (28.6%)	2 (14.3%)	0	7 (50%)
Ventersdorp	14 (100%)	12 (85.7%)	4 (28.6%)	0	0	4 (28.6%)

Table 7: Proportion of beef and mince beef isolates that were positive for confirmatory biochemical tests

Areas of collection	API 20E	O157 serotyping	H7 serotyping	API 20E	O157 serotyping	H7 serotyping
	Beef isolates			Mince beef isolates		
Brits	3 (21.4%)	1 (7.1%)	1 (7.1%)	3 (21.4%)	2 (14.3%)	2 (14.3%)
Garankwa	2 (14.3%)	1 (7.1%)	1 (7.1%)	2 (14.3%)	2 (14.3%)	2 (14.3%)
Rustenburg	0	0	0	1 (7.1%)	0	0
Vryburg	1 (7.1%)	1 (7.1%)	1 (7.1%)	5 (35.7%)	3 (21.4%)	2 (14.3%)
Koster	0	0	0	4 (28.6%)	4 (28.6%)	4 (28.6%)
Bloemhof	3 (21.4%)	1 (7.1%)	1 (7.1%)	6 (42.9%)	6 (42.9%)	6 (42.9%)
Taung	0	0	0	0	0	0
Swaartruggens	0	0	0	1 (7.1%)	1 (7.1%)	1 (7.1%)
Klerksdorp	1 (7.1%)	1 (7.1%)	1 (7.1%)	2 (14.3%)	1 (7.1%)	1 (7.1%)
Ventersdorp	0	0	0	0	0	0
Zeerust	0	0	0	2 (14.3%)	1 (7.1%)	1 (7.1%)
Delareyville	4 (28.6%)	0	0	1 (7.1%)	0	0
Swarzrenerke	6 (42.9%)	1 (7.1%)	1 (7.1%)	2 (14.3%)	1 (7.1%)	1 (7.1%)
Lichtenburg	9 (64.3%)	1 (7.1%)	1 (7.1%)	3 (21.4%)	1 (7.1%)	1 (7.1%)
Potchefstroom	6 (42.9%)	1 (7.1%)	1 (7.1%)	5 (35.7%)	4 (28.6%)	4 (28.6%)
Mafikeng	4 (28.6%)	3 (21.4%)	3 (21.4%)	1 (7.1%)	1 (7.1%)	1 (7.1%)
Total	39	11	11	38	27	27

Table 8: Proportion of *E. coli* O157:H7 isolates and *E. coli* O157:H7 specific virulent determinants in isolates from beef samples

Sample areas	No of isolates screened	No positive for the <i>rfb</i> _{O157} gene fragment	No positive for the <i>fliC</i> _{H7} gene fragment	No of isolates positive for the <i>stx1</i> gene fragment	No of isolates positive for the <i>stx2</i>	No of isolates positive for the <i>theae</i>	No of isolates positive for the <i>hyla</i>
Brits	14	1	1	1	1	1	1
Garankwa	14	1	1	0	1	1	1
Mafikeng	14	2	2	1	2	2	2
Vryburg	14	1	1	1	1	0	0
Koster	14	0	0	0	0	0	0
Bloemhof	14	0	0	0	0	0	0
Taung	14	0	0	0	0	0	0
Swaartruggens	14	0	0	0	0	0	0
Klerksdorp	14	1	1	0	1	0	0
Zeerust	14	0	0	0	0	0	0
Delareyville	14	0	0	0	0	0	0
Swarzrenerke	14	0	0	0	0	0	0
Lichtenburg	14	1	1	0	1	1	1
Potchefstroom	14	1	1	0	1	1	1
Rustenburg	14	0	0	0	0	0	0
Ventersdorp	14	0	0	0	0	0	0
Total	224	8	8	3	8	6	6

Delareyville, Swartzrenerke, Rustenburg and Ventersdorp as well as mince beef samples from Taung, Zeerust, Delareyville, Rustenburg and Ventersdorp but the detection of the pathogen in some of the samples was a cause for concern. The findings of this paper revealed that beef and mince beef samples obtained from some supermarkets in the area may pose severe health

risks on consumers if not properly cooked before consumption.

DISCUSSION

Shiga-toxigenic *E. coli* particularly those belonging to the serotype O157:H7 have received remarkable attention recently even in countries

Table 9: Proportion of *E. coli* O157:H7 isolates and *E. coli* O157:H7 specific virulent determinants in isolates from mince beef samples

Sample areas	No of isolates screened	No positive for the <i>rfb</i> _{O157} gene fragment	No positive for the <i>fliC</i> _{H7} gene fragment	No of isolates positive for the <i>stx1</i> gene fragment	No of isolates positive for the <i>stx2</i>	No of isolates positive for the <i>eae</i>	No of isolates positive for the <i>hlyA</i>
Brits	14	2	2	1	2	2	2
Garankwa	14	2	2	0	1	2	2
Mafikeng	14	1	1	1	1	0	1
Vryburg	14	3	3	3	3	0	3
Koster	14	4	4	1	3	0	2
Bloemhof	14	5	5	0	0	0	0
Taung	14	0	0	0	0	0	0
Swaartruggens	14	1	1	0	0	0	0
Klerksdorp	14	1	1	0	1	0	0
Zeerust	14	0	0	0	0	0	0
Delareyville	14	0	0	0	0	0	0
Swarzrenerke	14	1	1	0	0	0	0
Lichtenburg	14	1	1	0	1	1	1
Potchefstroom	14	2	2	1	1	1	1
Rustenburg	14	0	0	0	0	0	0
Ventersdorp	14	0	0	0	0	0	0
Total	224	23	23	7	13	6	12

with advance more public health and health care facilities and this mainly due to their potential to cause a number of food-borne diseases in humans (Belongia et al. 1991; Meichtri et al. 2004; Harakeh et al. 2005; Amani et al. 2015; Kargar and Homayoon 2015). Sporadic cases of infections and outbreaks of *E. coli* O157:H7 infections usually result from the consumption of undercooked contaminated beef (Williams et al. 2000; Gordillo et al. 2014; Ahmed and Shimamoto 2015; Dong et al. 2015; Hessain et al. 2015; Xu et al. 2016). The primary aim of this paper was to determine the level of microbial contamination in beef and mince beef samples obtained from some supermarkets in the major cities in the North West Province. Based on colonial appearance large proportion (60%) of the samples was contaminated with potential heterotrophic bacteria as well as potential *E. coli* O157:H7.

A further objective of the paper was to identify *E. coli* O157:H7 using preliminary and confirmatory biochemical tests. Despite the fact that only a small proportion (8 and 11) of the isolates from beef and mince beef respectively were positively identified as *E. coli* O157:H7 the detection of the pathogen in some of the meat samples was a cause for concern. Similar observations have been reported (Ateba and Mbewe 2011). It has been reported that food products

of animals origin particularly beef has identified as a potential source for the transmission of these pathogens to humans (Meichtri et al. 2004; Ateba et al. 2008; Ojo et al. 2010; Ateba and Mbewe 2011; Evans et al. 2011; Ateba et al. 2013; Gordillo et al. 2014; Ahmed and Shimamoto 2015; Dong et al. 2015; Hessain et al. 2015; Xu et al. 2016). In most developing countries, proper hygiene practices are not always strictly implemented during the handling and processing (Harakeh et al. 2005). The equipment used for cutting meat and the environment in which the meat is stored as well as the hygiene status of employees in the meat processing plant and proper farm management techniques have been reported to contribute significantly to the level of *E. coli* O157:H7 contamination (Elder et al. 2000; Blanco et al. 2003; Conedera et al. 2004; Loneragan et al. 2005; Magwira et al. 2005; Dong et al. 2015; Hsu et al. 2015). In addition, strategies that limit person-to-person transmission of *E. coli* O157:H7 may greatly reduce both sporadic and outbreak cases of infections in humans (Seto et al. 2007). It is therefore important to constantly determine the occurrence of this pathogen in food products so as to reduce human infections. In the present paper *E. coli* O157:H7 isolates possessed virulence gene *stx1*, *stx2*, *eaeA* and *hlyA*. The presence of these genes in *E. coli* O157:H7 are re-

sponsible for their pathogenicity (Oswald et al. 2000; Brusa et al. 2012; Lorente et al. 2014; Claudia et al. 2015).

CONCLUSION

In conclusion, *E. coli* O157:H7 was successfully isolated in the study and the isolates possessed virulence gene determinants. Given that this pathogen can be introduced into the food chain if proper hygiene practices are not strictly implemented during the handling and processing of food. Against this background there is need to develop and implement control measures that will limit the transmission of this pathogen to humans.

REFERENCES

- Abraham S, Jordan D, Wong HS, Johnson JR, Toleman MA, Wakeham DL, Gordon DM, Turnidge JD, Mollinger JL, Gibson JS, Trott DJ 2015. First detection of extended-spectrum cephalosporin- and fluoroquinolone-resistant *Escherichia coli* in Australian food-producing animals. *Journal of Global Antimicrobial Resistance*, 3(4): 273-277.
- Ahmed AM, Shimamoto T 2015. Molecular analysis of multidrug resistance in Shiga toxin-producing *Escherichia coli* O157:H7 isolated from meat and dairy products. *International Journal of Food Microbiology*, 193: 68-73.
- Amani J, Ahmadpour A, Fooladi AAI, Nazarian S 2015. Detection of *E. coli* O157:H7 and *Shigella dysenteriae* toxins in clinical samples by PCR-ELISA. *The Brazilian Journal of Infectious Diseases*, 19(3): 278-284.
- Ateba CN, Mbewe M 2013. Determination of the genetic similarities of fingerprints from *Escherichia coli* O157:H7 isolated from different sources in the North West Province, South Africa using ISR, BOXAIR and REP-PCR analysis. *Microbiology Research*, 168: 438-446.
- Ateba CN, Mbewe M 2011. Detection of *Escherichia coli* O157:H7 virulence genes in isolates from beef, pork, water, human and animal species in the north-west province, South Africa: Public health implications. *Research in Microbiology*, 162(3): 240-248.
- Ateba CN, Mbewe M, Bezuidenhout CC 2008. The Prevalence of *Escherichia coli* O157 strains in cattle, pigs and humans in the North-West Province, South Africa. *South African Journal of Science*, 104: 7-8.
- Banatvala N, Griffin PM, Greene KD, Barrett TJ, Bibb WF, Green JH, Wells JG 2001. The United States national prospective haemolytic uremic syndrome study: Microbiologic, serologic, clinical, and epidemiologic findings. *Journal of Infectious Disease*, 183: 1063-1070.
- Belongia EA, MacDonald KL, Parham GL, White KE, Korlath MN, Lobato SM, Strand SM, Kasale KA, Osterholm MT 1991. An outbreak of *Escherichia coli* O157:H7 colitis associated with consumption of precooked meat patties. *Journal of Infectious Disease*, 164: 338-343.
- Bhakdi S, Muhly M, Korom S, Schmidt G 1990. Effects of *Escherichia coli* haemolysin on human monocytes. Cytocidal action and stimulation of interleukin 1 release. *Journal of Clinical Investigation*, 85: 1746-1753.
- Blanco M, Blanco JE, Mora A, Dahbi G, Alonso MP, Gonzalez EA, Bernardez MI, Blanco J 2004. Serotypes, virulence genes and intimin types of shiga toxin (Verotoxin) - Producing *Escherichia coli* isolates from cattle in Spain and identification of new intimin variant Gene. *Journal of Clinical Microbiology*, 42: 645 -651.
- Blanco J, Blanco M, Blanco EJ, Mora A, Gonzalez AE, Bernardez IM, Alonso MP, Coira A, Rodriguez A, Rey J, Alonso MJ, Usera AM 2003. Verotoxin - producing *Escherichia coli* in Spain: Prevalence, serotypes, and virulence genes of O157:H7. *Experimental Biology and Medicine*, 228: 345 -351.
- Brusa V, Aliverti V, Aliverti F, Ortega EE, de la Torre JH, Linares LH, Sanz ME, Etcheverría AI, Padola NL, Galli L, García PP, Copes J, Leotta GA 2012. Shiga toxin-producing *Escherichia coli* in beef retail markets from Argentina. *Frontiers of Cellular and Infection Microbiology*, 2: 171.
- Bugarel M, Beutin L, Martin A, Gill A, Fach P 2010. Micro-array for the identification of Shiga toxin-producing *Escherichia coli* (STEC) seropathotypes associated with Hemorrhagic Colitis and Hemolytic Uremic Syndrome in humans. *International Journal of Food Microbiology*, 142: 318-329.
- Chapman PA, Malo Cerdan AT, Ellin M, Ashton R, Harkin MA 2001. *Escherichia coli* O157:H7 in cattle and sheep at slaughter, on beef and lamb carcasses and in raw beef and lamb products in South Yorkshire, United Kingdom. *International Journal of Food Microbiology*, 64: 139-150.
- Claudia NV, AlejandroE, Markus MF, Argenis RG, Todd BM, Mueen A, Mindy BM 2015. Virulence characterization and molecular subtyping of typical and atypical *Escherichia coli* O157:H7 and O157:H(-) isolated from fecal samples and beef carcasses in Mexico. *Journal of Food Protection*, 78: 264-272.
- Conedera G, Dalvit P, Martini M, Galiero G, Gramaglia M, Goffredo E et al. 2004. Verocytotoxin-producing *Escherichia coli* O157 in minced beef and dairy products in Italy. *International Journal of Food Microbiology*, 96: 67-73.
- Díaz-Sánchez S, Sánchez S, Herrera-León S, Porrero C, Blanco J, Dahbi G, Blanco JE, Mora A, Mateo R, Hanning I, Vidal D 2013. Prevalence of Shiga toxin-producing *Escherichia coli*, *Salmonella* spp. and *Campylobacter* spp. in large game animals intended for consumption: Relationship with management practices and livestock influence. *Veterinary Microbiology*, 163(3-4): 274-281.
- Dong P, Zhu L, Mao Y, Liang R, Niu L, Zhang Y, Luo X 2015. Prevalence and characterization of *Escherichia coli* O157:H7 from samples along the production line in Chinese beef-processing plants. *Food Control*, 54: 39-46.

- Doyle JJ, Doyle JL 1990. A rapid total DNA preparation procedure for fresh plant tissue. *Focus*, 12: 13-15.
- Elder RO, James EK, Gregory RS, Gallagher-Barkocy GA, Koohmaraie M, Laegrid WW 2000. Correlation of Enterohemorrhagic *Escherichia coli* O157:H7 prevalence in feces, hides, and carcasses of beef cattle during processing. *Proceedings of the National Academy of Science*, 97: 2999-3003.
- Evans J, Knight H, McKendrick IJ, Stevenson H, Barbudo AV, Gunn GJ, Low JC 2011. Prevalence of *Escherichia coli* O157:H7 and serogroups O26, O103, O111 and O145 in sheep presented for slaughter in Scotland. *Journal of Medical Microbiology*, 60: 653-660.
- Gannon VP, D'Souza S, Graham T, King RK, Rahn K., Read S 1997. Use of Flagella H7 as a target in multiplex PCR assays and improved specificity in identification of enterohaemorrhagic *Escherichia coli* strains. *Journal of Clinical Microbiology*, 35: 656-662.
- Gordillo R, Rodríguez A, Werning ML, Bermúdez E, Rodríguez M 2014. Quantification of viable *Escherichia coli* O157:H7 in meat products by duplex real-time PCR assays. *Meat Science*, 96(2): 964-970.
- Harakeh S, Yassine H, Gharios M, Barbour E, Hajjar S, El-Fadel M, Toufeili I, Tannous R 2005. Isolation, molecular characterization and antimicrobial resistance patterns of *Salmonella* and *Escherichia coli* isolates from meat-based fast food in Lebanon. *Science of the Total Environment*, 134 : 33-44.
- Hessain AM, Al-Arfaj AA, Zakri AM, El-Jakee JK, Al-Zogibi OG, Hemeg HA, Ibrahim IM 2015. Molecular characterization of *Escherichia coli* O157:H7 recovered from meat and meat products relevant to human health in Riyadh, Saudi Arabia. *Saudi Journal of Biological Sciences*, 22(6): 725-729.
- Hsu HY, Sheen S, Sites J, Cassidy J, Scullen B, Sommers C 2015. Effect of high pressure processing on the survival of Shiga toxin-Producing *Escherichia coli* (Big Six vs. O157:H7) in ground beef. *Food Microbiology*, 48: 1-7.
- Kargar M, Homayoon M 2015. Prevalence of shiga toxins (*stx1*, *stx2*), *eaeA* and *hly* genes of *Escherichia coli* O157:H7 strains among children with acute gastroenteritis in southern of Iran. *Asian Pacific Journal of Tropical Medicine*, 8(1): 24-28.
- Law D 2000. Virulence factors of *Escherichia coli* O157:H7 and other Shiga toxin – producing *Escherichia coli*. *Journal of Applied Microbiology*, 88: 729-745.
- Liu Y, Wang Y, Walsh TR, Yi L, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, Yu L, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu J, Shen J 2016. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: A microbiological and molecular biological study. *The Lancet Infectious Diseases*, 16 (2): 161-168.
- Loneragan GH, Brashears MM 2005. Pre-harvest interventions to reduce carriage of *E. coli* O157 by harvest-ready feedlot cattle. *Meat Science*, 71: 72-78.
- Magwira CA, Gashe BA, Collison EK 2005. Prevalence and antibiotic resistance profiles of *Escherichia coli* O157:H7 in beef products from retail outlets in Gabone Botswana. *Journal of Food Protection*, 68: 403-406.
- McEvoy JM, Doherty AM, Sheridan JJ, Thompson-Carter FM, Garvey P, McGuire L, Blair IS, McDowell DA 2003. The prevalence and spread of *Escherichia coli* O157:H7 at a commercial beef Abattoir. *Journal of Applied Microbiology*, 95: 256-266.
- Meichtri L, Miliwebsky E, Gioffre I, Chinen A, Baschier G, Chillemi BEC, Guth MO, Masana A, Cataldi RH, Rodriguez M, Rivas 2004. Shiga Toxin –Producing *Escherichia Coli* in healthy young beef steers from Argentina: Prevalence and virulence properties. *International Journal of Food Microbiology*, 96: 189-198.
- Momba MNB, Abongo BO, Mwambakana JN 2008. Prevalence of enterohaemorrhagic *Escherichia coli* O157:H7 in drinking water and its predicted impact on diarrhoea HIV/AIDS patients in the Amathole District, Eastern Cape Province, South Africa. *Water SA*, 34: 365-372.
- Müller EE, Taylor MB, Grabow WOK, Ehlers MM 2002. Isolation and characterisation of *Escherichia coli* O157:H7 and shiga toxin-converting bacteriophages from strains of human, bovine and porcine origin. *Water Supplement*, 2: 29-38.
- Muyzer G, Teske A, Wirsén CO, Jannasch HW 1995. Phylogenetic relationships of *thiomicrospira* species and their identification in deep-sea hydrothermal vent by denaturing gradient electrophoresis of 16S rDNA fragments. *Archives of Microbiology*, 164: 165-172.
- Ojo OE, Ajuwape ATP, Otesile EB, Owoade AA, Oyekunle MA, Adetosoye AI 2010. Potentially zoonotic shiga toxin-producing *Escherichia coli* serogroups in the faeces and meat of food-producing animals in Ibadan, Nigeria. *International Journal of Food Microbiology*, 142(1-2): 214-221.
- Oswald E, Schmidt H, Morabito S, Karch H, Marches O, Capioli A 2000. Typing of intimin genes in human and animal enterohaemorrhagic and enteropathogenic *Escherichia coli*: Characterisation of a new intimin variant. *Infection and Immunity*, 68: 64-71.
- Paton AW, Paton JM 1998. Detection of Shiga toxin-producing *Escherichia coli* strains belonging to serogroups O111, O157, O113 by multiplex PCR. *Journal of Clinical Microbiology*, 37: 3362-3365.
- Rangel JM, Sparling PH, Crowe C, Griffin PM, Swerdlow DL 2005. Epidemiology of *Escherichia coli* O157:H7 outbreaks, United States, 1982–2002. *Emerging Infectious Diseases*, 11: 603-609.
- Sambrook J, Fritsch EF, Maniatis T 1989. *Molecular Cloning. A Laboratory Manual*. 2nd Edition. USA: Cold Spring Harbour Laboratory Press.
- Schouten JM, van de Giessen AW, Frankena K, De Jong MC, Graat EA 2005. *Escherichia coli* O157 prevalence in Dutch poultry, pig finishing and veal herds and risk factors in Dutch vealherds. *Preventative Veterinary Medicine*, 70: 1-15.
- Seto EYW, Soller JA, Colford Jr JM 2007. Strategies to reduce person-to-person transmission during widespread *Escherichia coli* O157:H7 outbreak. *Emerging Infectious Disease*, 13: 860-866.
- Stampi S, Caprioli A, Deluca G, Quaglio P, Sacchetti R, Zannetti F 2004. Detection of *Escherichia coli* O157

- in Bovine meat products in Northern Italy. *International Journal of Food Microbiology*, 90: 257-262.
- Tamang DM, Gurung M, Kang m, Nam H, Moon DC, Jang G, Jung S, Park YH, Lim S 2015. Characterization of plasmids encoding CTX-M β -lactamase and their addiction systems in *Escherichia coli* isolates from animals. *Veterinary Microbiology*, 174(3-4): 456-462.
- Vijay D, Dhaka P, Vergis J, Negi M, Mohan V, Kumar M, Doijad S, Poharkar K, Kumar A, Malik SS, Barbuddhe SB, Rawool DB 2015. Characterization and biofilm forming ability of diarrhoeagenic enteroaggregative *Escherichia coli* isolates recovered from human infants and young animals. *Comparative Immunology, Microbiology and Infectious Diseases*, 38: 21-31.
- Weir E 2000. *Escherichia coli* O157:H7. *Canadian Medical Association Journal*, 163: 205.
- Williams RC, Isaacs S, Decou ML, Richardson EA, Buffet MC, Slinger RW 2000. Illness outbreak associated with *Escherichia coli* O157:H7 in Genoa salami. *Canadian Medical Association Journal*, 162: 1409-1413.
- Wong CS, Jelacic S, Habeeb RL, Watkins SL, Tarr PI 2000. The risk of the haemolytic-uremic syndrome after antibiotic treatment of *E. coli* O157:H7 infections. *New England Journal Medicine*, 342: 1930-1936.
- Xu M, Wang R, Li Y 2016. Rapid detection of *Escherichia coli* O157:H7 and *Salmonella Typhimurium* in foods using an electrochemical immunosensor based on screen-printed interdigitated microelectrode and immunomagnetic separation. *Talanta*, 148: 200-208.
- Zheng H, Zeng Z, Chen S, Liu Y, Yao Q, Deng Y, Chen X, Lv L, Zhuo C, Chen Z, Liu J 2012. Prevalence and characterisation of CTX-M β -lactamases amongst *Escherichia coli* isolates from healthy food animals in China. *International Journal of Antimicrobial Agents*, 39(4): 305-310.