

Assessment of Nutritional Status among Muslim Adolescents of Deganga, North 24 Parganas, West Bengal

Argina Khatun¹, Ashish Mukhopadhyay² and Kaushik Bose¹

¹*Department of Anthropology, Vidyasagar University, Midnapore, West Bengal, India*

²*Department of Anthropology, A.P.C. College, New Barrackpore, West Bengal, India*

KEYWORDS Anthropometry. Body Mass Index. Thinness. Overweight. Undernutrition

ABSTRACT The purpose of this study was to examine the age and sex differences in nutritional status based on new age and sex specific international BMI cut-off points, among Muslim adolescents of North 24 Parganas, West Bengal. One thousand and sixty eight (1068) Bengali speaking Muslim school boys (n=522) and girls (n=546) aged 10-17 years participated in this study. The overall (combined age and sex) frequency of undernutrition was 54.8 percent. The frequencies (combined age) of undernutrition among boys and girls were 58.04 percent and 51.65 percent respectively. The rates of undernutrition of boys varied between 35.0 percent among 16 years old to 72.7 percent at age 13 years. The rates of undernutrition of girls varied between 35.7 percent among 15 years old to 71.2 percent at age 11 years. Significant age difference was observed in mean height, weight and BMI for boys (height: $F=234.37$, $p<0.05$, weight: $F=145.36$, $p<0.05$ and BMI: $F=27.47$, $p<0.05$) and as well as girls (height: $F=163.40$, $p<0.05$, weight: $F=107.82$, $p<0.05$ and BMI: $F=39.74$, $p<0.05$). There was significant age differences (boys, chi square= 62.80, $p=0.000$, girls, Chi-square = 76.21, $p=0.000$) in nutritional status based on BMI of studied children. In general, this study provided evidence that the Bengalee Muslim adolescents had very high rates of undernutrition. Most of the earlier Indian studies showed lower prevalence than present study.

INTRODUCTION

World Health Organization identifies adolescence as the period in human growth and development that occurs after childhood and before adulthood, from ages 10 to 19 years. The World Bank report says that the rate of malnutrition cases among children in India is almost five times more than in China, and twice than in Sub-Saharan Africa. Anthropometry has been used during adolescence in many contexts related to nutritional status (WHO 1995; Bose and Mukhopadhyay 2004). Malnutrition (undernutrition or overnutrition) which refers to an impairment of health either from a deficiency or excess or imbalance of nutrients, is of public health significance among adolescent all over the world. It creates lasting effect on the growth, development and physical fitness of a person. It is well recognized worldwide that anthropometric measurements are indispensable in diagnosing un-

dernutrition. It has now been well established that the body mass index (BMI) is the most appropriate variable for determining nutritional status among adolescents (WHO 1995; Himes and Bouchard 1989; Must et al. 1991; Rolland-Cachera 1993). Several recent studies have investigated nutritional status of children from different parts of India (Kanade et al. 1999; Singh and Mishra 2001; Venkaiah et al. 2002, Mondal et al. 2011; Bansal et al. 2013; Mondal N 2014; Das et al. 2015; Bharti et al. 2017). The researchers have already published two papers (Khatun et al. 2016a, b) on nutritional status comprising of the same sample using the World Health Organization (WHO 1995) recommended age and sex specific cut-off points of BMI based on the NHANES-I percentile values. However, there is scanty information on the nutritional status using new age and sex specific international BMI cut-off points (Cole et al. 2000, 2007) of Muslim adolescents from rural West Bengal. The present interpretation attempted to evaluate the nutritional status and to assess age-sex trends in the level of different categories of nutritional status on international BMI cut-off points among 10-

Address for correspondence:
Prof. Kaushik Bose
Department of Anthropology,
Vidyasagar University,
Midnapore, West Bengal, India

17 years old Bengalee Muslim adolescents of North 24 Parganas, West Bengal.

MATERIAL AND METHODS

The investigation was carried out among adolescent Muslim boys and girls in three secondary schools in Deganga block, North 24 Parganas, West Bengal. The schools are situated in a rural area approximately 38 km to the east from the heart of the city of Kolkata and approximately 17 km from the Barasat town. Barasat is the administrative headquarters of the district of North 24 Parganas, West Bengal, India.

One thousand and sixty eight school boys (n=522) and girls (n=546) participated in this study. The age range of the subjects was 10 to 17 years.

A total of twelve anthropometric measurements (height, weight, five circumferences and five skinfolds) were made in the present field survey but only two basic variables (height and weight) and a single derived variable (body mass index) have been used in the present report. All the anthropometric measurements were taken following the standard techniques recommended by Lohman et al. (1988) and body mass index (BMI) was computed using the standard equation: $BMI (kg / m^2) = Weight (kg) / Height^2 (m^2)$.

Nutritional status of the studied children was evaluated using new age and sex specific international BMI cut-off points (Cole et al. 2000 and

2007). Statistical Package for Social Sciences (SPSS) and Microsoft Office Excel (2010) were used to analyze the data. Statistical significance was set at $p < 0.05$ level. This cut-off point has been utilized by several recent studies worldwide on undernutrition among adolescents (Venkaiah et al. 2002; Woodruff and Duffield 2002).

Technical errors of measurements (TEM) were calculated and the results were found to be within reference values cited by Ulijaszek and Kerr (1999). Therefore, TEM was not incorporated in statistical analyses.

RESULTS

The means and standard deviations and F values of the anthropometric characteristics by age groups of the boys and girls are presented in Table 1. Height of boys and girls increased progressively (37.09 cm and 25.02 cm) from 10 to 16 years. Weight of boys showed a steady positive age trend from 13 (28.28 kg) to 16 (50.09 kg) years of age. Weight of girls showed a steady positive age trend from 10 (23.90 kg) to 15 (43.27 kg) years of age. Mean BMI of boys increased progressively by 3.85 kg/m² from 10 years to 17 years with a pause at age 14 and 17 years. Mean BMI of girls increased gradually by 4.42 kg/m² from 10 years to 15 years. Significant age difference was observed in mean height, weight and BMI for boys (height: $F=234.37$, $p < 0.05$, weight: $F=145.36$, $p < 0.05$ and BMI: $F=27.47$, $p < 0.05$) and as well as girls (height: $F=163.40$, $p < 0.05$,

Table 1: Age and sex variation in anthropometric characteristics of 10-17 years old rural Bengalee Muslim adolescents

Age groups (years)	Sex	Height (cm)		Weight (kg)		Body mass index (kg/m ²)
10.0-10.9	Boys (n=56)	127.58	(4.70)	23.66	(2.46)	14.51 (1.00)
	Girls (n=62)	127.71	(4.59)	23.90	(2.89)	14.63 (1.26)
11.0-11.9	Boys (n=63)	131.55	(6.00)	26.01	(4.26)	14.98 (1.87)
	Girls (n=73)	133.28	(5.34)	26.34	(4.89)	14.78 (2.28)
12.0-12.9	Boys (n=65)	135.77	(5.76)	28.98	(7.99)	15.56 (3.17)
	Girls (n=82)	140.95	(6.04)	30.54	(6.19)	15.30 (2.42)
13.0-13.9	Boys (n=77)	136.09	(6.30)	28.28	(5.11)	15.18 (1.81)
	Girls (n=79)	146.80	(7.45)	37.09	(6.84)	17.09 (2.19)
14.0-14.9	Boys (n=84)	145.83	(9.91)	34.95	(7.87)	16.27 (2.19)
	Girls (n=78)	149.45	(5.82)	39.59	(6.24)	17.68 (2.33)
15.0-15.9	Boys (n=71)	158.06	(11.05)	46.09	(10.77)	18.43 (4.15)
	Girls (n=70)	150.56	(5.46)	43.27	(7.88)	19.05 (3.07)
16.0-16.9	Boys (n=60)	164.67	(5.97)	50.09	(7.00)	18.45 (2.22)
	Girls (n=55)	152.73	(4.76)	42.65	(5.85)	18.27 (2.30)
17.0-17.9	Boys (n=46)	163.51	(5.10)	49.06	(5.47)	18.36 (1.97)
	Girls (n=47)	151.82	(5.98)	43.31	(6.21)	18.76 (2.22)
F- ratio	Boys (n=522)	234.37*		145.36*		27.47*
	Girls (n=546)	163.40*		107.82*		39.74*

Standard deviations are presented in parentheses

Table 2: Prevalence of nutritional status (based on BMI) of 10-17 years old rural Bengalee Muslim adolescents

Age groups (years)	Sex	Grade III orthinness 1	Grade II orthinness 2	Grade I orthinness 3	Overall CED	Normal	Overweight
10.0-10.9	Boys	-	8 (14.3)	27 (48.2)	35 (62.5)	21 (37.5)	-
	Girls	2 (3.2)	8 (12.9)	27 (43.5)	37 (59.7)	25 (40.3)	-
11.0-11.9	Boys	3 (4.8)	9 (14.3)	28 (44.4)	40 (63.5)	22 (34.9)	01 (1.6)
	Girls	10 (13.7)	15 (20.5)	27 (37.0)	52 (71.2)	19 (26.0)	02 (2.7)
12.0-12.9	Boys	3 (4.6)	16 (24.6)	25 (38.5)	44 (67.7)	19 (29.2)	02 (3.1)
	Girls	19 (23.2)	14 (17.1)	22 (26.8)	55 (67.1)	25 (30.5)	02 (2.4)
13.0-13.9	Boys	11 (14.3)	16 (20.8)	29 (37.7)	56 (72.7)	21 (27.8)	-
	Girls	5 (6.3)	7 (8.9)	18 (22.8)	30 (38.0)	48 (60.8)	01 (1.3)
14.0-14.9	Boys	10 (11.9)	15 (17.9)	25 (29.8)	50 (59.5)	32 (38.1)	02 (2.4)
	Girls	3 (3.8)	8 (10.3)	23 (29.5)	34 (43.6)	41 (52.6)	03 (3.8)
15.0-15.9	Boys	5 (7.0)	7 (9.9)	22 (31.0)	34 (47.9)	28 (39.4)	9 (12.7)
	Girls	2 (2.9)	6 (8.6)	17 (24.3)	25 (35.7)	41 (58.6)	04 (5.7)
16.0-16.9	Boys	3 (5.0)	4 (6.7)	14 (23.3)	21 (35.0)	37 (61.7)	27 (49.1)
	Girls	2 (3.6)	9 (16.4)	17 (30.9)	28 (50.9)	02 (3.3)	-
17.0-17.9	Boys	1 (2.2)	8 (17.4)	14 (30.4)	23 (50.0)	23 (50.0)	-
	Girls	2 (4.3)	7 (14.9)	12 (25.5)	21 (44.7)	26 (55.3)	-
All ages	Boys	36 (6.9)	83 (15.9)	184 (35.2)	303 (58.0)	203 (38.9)	16 (3.1)
	Girls	45 (8.2)	74 (13.6)	163 (29.9)	282 (51.7)	252 (46.2)	12 (2.2)
Overall	Boys+Girls	81 (7.6)	157 (14.7)	347 (32.5)	585 (54.8)	455 (42.6)	28 (2.6)
Chi-square		(Boys) Chi square (df=28) = 62.80, P= 0.000			(Girls) Chi-square (df=28) = 76.21, P= 0.000		

weight: $F=107.82$, $p<0.05$ and BMI: $F=39.74$, $p<0.05$).

The overall (combined age and sex) frequency of undernutrition was 54.8 percent (Table 2). The frequencies (combined age) of undernutri-

tion among boys and girls were 58.04 percent and 51.65 percent respectively. The rates of undernutrition of boys varied between 35.0 percent among 16 years old to 72.7 percent at age 13 years. The rates of undernutrition of girls


**Fig. 1. Prevalence of undernutrition among Muslim adolescents**

Table 3: Comparison of the prevalence of undernutrition among Indian children

<i>Studied children</i>	<i>Age groups</i>	<i>n</i>	<i>Methods</i>	<i>Prevalence (%)</i>	<i>References</i>
School children, Bankura District, West Bengal	6-14	454	BMI/age Z-score	23.1	Bose et al., 2008
School children, Paschim Medinipur and Purulia District, West Bengal	10-15	2016	Thinness by BMI	44.5	Bose and Bisai, 2008
School children, Dibrugarh district, Assam	6-14	304	Thinness by BMI	53.9	Medhi et al., 2006
Children, Dibrugarh district, Assam	10-18	605	Thinness by BMI	50.2	Medhi et al., 2006
Kora-Mudi tribal children, Paschim Medinipur, West Bengal	2-13	119	Thinness by BMI	67.2	Bisai et al., 2010
Muslim school children, Nadia, West Bengal	6-16	638	Thinness by BMI	56.7	Mondal et al., 2011
Paschim Medinipur, West Bengal	6-12	500	Thinness by BMI	77.0	Das et al., 2012
School children, Darjeeling, West Bengal	10-18	1165	Thinness by BMI	49.1	Mondal N, 2014
North 24 Parganas, West Bengal	10-17	1068	Thinness by BMI	54.8	Present study

varied between 35.7 percent among 15 years old to 71.2 percent at age 11 years. The overall (combined age and sex) frequency of undernutrition was 28 percent. A distinctive age variation in the change of the rate of undernutrition was observed in both sexes (Fig. 1). Boys demonstrated a steady increase in the rate of undernutrition from 10 to 13 years of age. Thereafter, the rate decreased from 14 to 16 years then increased at age 17 years. Contrarily, among girls there was a substantial increase in the rate of undernutrition from 10 to 11 years of age followed by a distinctive decrease from 12 to 13 years, and then it is increase at age 14 years and slightly decreases at age 15 years. Thereafter, there was an increase at the 16 years and a slight decrease at 17 years. There was significant age differences (boys, Chi-square= 62.80, $p = 0.000$, girls, Chi-square= 76.21, $p = 0.000$) in nutritional status based on BMI of studied children.

DISCUSSION

Adolescence is a period of increased nutritional requirements and adolescent anthropometry varies significantly worldwide (WHO 1995; Himes and Bouchard 1989; Bhadra et al. 2001). Undernutrition is a significant problem and continues to be a cause of morbidity and mortality among children in developing countries like India (UNICEF 2006). Cole et al. (2007) has stated that undernutrition is better assessed as thinness (low body mass index for age) than as wasting (low weight for height). Malnutrition (un-

der-nutrition and over-nutrition) continues to be a problem of considerable magnitude in most developing countries of the world (Som et al. 2006). The mean height, weight and BMI of the Bengalee Muslim boys and girls of this study were higher than those reported among rural adolescents in a recent study from India (ICMR 1996). The comparison of prevalence of undernutrition among Indian children with present study is presented in Table 3. In the present investigation more than half (54.8%) of the rural Bengalee Muslim adolescents was undernourished. The rate of undernutrition of children of present study was higher than those reported by six Indian studies, that is, 23.1 percent (Bankura District, Bose et al. 2008), 44.5 percent (Paschim Medinipur and Purulia Districts, Bose and Bisai 2008), 53.9 percent, 50.2 percent (Dibrugarh District, Assam, Medhi et al. 2006), and 49.1 percent (Darjeeling Districts, Mondal N. 2014). However, the rate of undernutrition of the present study is lower to those of two Paschim Medinipur studies that is, 67.2 percent (Bisai et al. 2010) and 77 percent (Das et al. 2012) and the Muslim school children of Nadia Districts that is, 56.7 percent (Mondal et al. 2011).

CONCLUSION

The present study revealed that both under and over nutrition is present study among Muslim school children.

This study provided three vital messages:

1. Rural Muslim adolescents of North 24 Parganas, West Bengal, had critical situation of undernutrition (overall = 54.8%, boys = 58.04% and girls = 51.65%).
2. There were straightforward age and sex variations in the rates of undernutrition prevalent among the studied samples.
3. These rates of undernutrition of the present study were higher than maximum earlier Indian findings (Bose et al. 2008, Bose and Bisai 2008, Medhi et al. 2006, Bansal et al. 2013 Mondal N. 2014).

ACKNOWLEDGEMENTS

The Head Masters and the school authorities of three schools are thankfully acknowledged for their help during data collection. Thanks are also due to the students who participated in this study.

REFERENCES

- Ahmed F, Zareen M, Khan MR, Banu CP, Haq MN, Jackson AA 1998. Dietary patterns, nutrient intake and growth of adolescent school girls in urban Bangladesh. *Pub Health Nutr*, 1: 83-92.
- Beghin I, Cap M, Dujardin B 1988. *A Guide to Nutritional Assessment*. Geneva, Switzerland: World Health Organization.
- Bhadra M, Mukhopadhyay A, Bose K 2001. Body mass index, regional adiposity and central body fat distribution among Bengalee Hindu girls: A comparative study of pre-menarcheal and menarcheal subjects. *Acta Med Auxol*, 33: 39-45.
- Bhadra M, Mukhopadhyay A, Bose K 2004. Sex differences in anthropometric characteristics among 11-14 year old urban Bengalees of North 24 Parganas, West Bengal, India. *Anthropologie*, In Press.
- Bharthi K, Ghritlahre M, Das S, Bose K 2017. Nutritional status among children and adolescents aged 6-18 years of Kolam tribe of Andhra Pradesh, India. *Anthropological Review*, 80(2): 13-23.
- Bisai S, Ghosh T, De GK, Bose K 2010. Very high prevalence of thinness among Kora-Mudi Tribal Children Paschim Medinipur District of West Bengal, India. *EJBS*, 3(1): 43-49.
- Bose K, Mukhopadhyay A 2004. Nutritional status of adolescent Bengalee Boys. *Indian Pediatrics*, 41: 633.
- Bose K, Bisai S 2008. Prevalence of undernutrition among rural adolescents of West Bengal, India. *J Trop Pediatr*, 54: 422-423.
- Bose K, Bisai S, Mukherjee S 2008. Anthropometric characteristics and nutritional status of rural school children. *Intern J Biol Anthropol*, 2(1):.
- Cole TJ, Bellizzi MC, Flegal KM, Dietz WH 2000. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*, 320: 1240-1243.
- Cole TJ, Flegal KM, Nicholls D, Jackson AA 2007. Body mass index cut offs to define thinness in children and adolescents: International survey. *BMJ*, 335: 194-198.
- Cookson ST, Woodruff BA, Slutsker L 1998. *Prevalence of Anemia and Low Body Mass Index among Adolescents 10-19 y of Age in Refugee Camps in Dadaab District, Kenya*. Atlanta, GA: Centers for Disease Control and Prevention.
- De Onis M, Dasgupta P, Saha S, Sengupta D, Blossner M 2001. The National Center for Health Statistics reference and the growth of Indian adolescent boys. *Am J Clin Nutr*, 74: 248-253.
- Himes JH, Bouchard C 1989. Validity of anthropometry in classifying youths as obese. *Int J Obes*, 13: 183-193.
- International Rescue Committee 1997. *Nutritional Status of School Aged Children in Kakuma Refugee Camp*. Nairobi: International Rescue Committee.
- Kanade AM, Joshi SB, Rao S 1999. Undernutrition and adolescent growth among rural Indian boys. *Indian Pediatrics*, 36: 145-156.
- Khatun A, Bhadra M, Mukhopadhyay A, Bose K 2016a. Anthropometric assessment of nutritional status of Bengalee Muslim adolescents of Deganga, North 24 Parganas, West Bengal, India. *Int J Exp Res Rev*, 4: 34-39.
- Khatun A, Bhadra M, Mukhopadhyay A, Bose K 2016b. Nutritional status and effect of physical activity on anthropometric characteristics of Bengalee Muslim adolescents boys of North 24 Parganas, West Bengal, India. *Int J Exp Res Rev*, 5: 8-14.
- Kurz KM 1996. Adolescent nutritional status in developing countries. *Proc Nutr Soc*, 55: 321-331.
- Lohman TG, Roche AF, Martorell R 1988. *Anthropometric Standardization Reference Manual*. Chicago: Human Kinetics Books.
- Medhi GK, Barua A, Mahanta J 2006. Growth and nutritional status of school age children (6-14 years) of tea garden worker of Assam. *Journal of Human Ecology*, 19: 83-85.
- Mondal N 2014. Thinness as major underlying problem among adolescents of north east India. *J Nepal Paediatr Soc*, 34(1): 39-47.
- Mondal PR, Biswas S, Bose K 2011. Prevalence of thinness among rural Bengalee Muslim school children from Eastern India. Jason E Lee (Ed.): *Undernutrition: Effects, Causes and Management*. New York: Nova Science Publishers, pp. 99-110.
- Must A, Dallal GE, Dietz WH 1991. Reference data for obesity: 85th and 95th percentiles of body mass index (wt/ht²) and triceps skinfold thickness. *Am J Clin Nutr*, 53: 839-846.
- Rahmathullah L, Underwood BA, Thulasiraj RD, Milton RC, Ramaswamy K, Rahmathullah R, Babu G 1990. Reduced mortality among children in Southern India receiving a small weekly dose of vitamin A. *N Engl J Med*, 323: 929-935.
- Rolland-Cachera MF 1993. Body composition during adolescence: methods, limitations and determinants. *Hormone Research*, 39: 25-40.
- Singh J, Mondal N 2014. Use of upper-arm anthropometry measure of body composition and nutritional assessment in children and adolescents (6-20 years) of Assam, Northeast India. *Ethiop J Health Sci*, 24(3): 243-252.

- Singh N, Mishra CP 2001. Nutritional status of adolescent girls of a slum community of Varanasi. *Indian J Public Health*, 45: 128-134.
- Som S, Pal M, Bhattacharya B, Bharati S, Bharati P 2006. Socioeconomic differentials in nutritional status of children in the states of West Bengal and Assam. *J Biosoc Sc*, 38: 625-642.
- Ulijaszek SJ, Kerr DA 1999. Anthropometric measurement error and the assessment of nutritional status. *Brit J Nutr*, 82: 165-177.
- UNICEF 2006. *Report Progress for Children – A Report Card on Nutrition*. New York: UNICEF.
- Venkaiah K, Damayanti K, Nayak MU, Vijayaraghavan K 2002. Diet and nutritional status of rural adolescents in India. *Eu J Clin Nutr*, 56: 1119-1125.
- Woodruff BA, Slutsker L, Cook ST 1998. *Prevalence of Anemia and Low Body Mass Index in Adolescents 10-19 y Age in Kakuma Camp, Kenya*. Atlanta, GA: Centers for Disease Control and Prevention..
- Woodruff BA, Duffield A, Blanck H, Larson MK, Pahari S, Bhatia R 1999. *Prevalence of Low Body Mass Index and Specific Micronutrient Deficiencies in Adolescents 10-19 y of Age in Bhutanese Refugee camps, Nepal, October 1999*. Atlanta, GA: Centers for Disease Control and Prevention.
- Woodruff BA, Duffield A 2002. Anthropometric assessment of nutritional status in adolescent populations in humanitarian emergencies. *Eu J Clin Nutr*, 56: 1108-1118.
- World Health Organization 1985. *Measuring Nutritional Status*. Geneva: World Health Organization.
- World Health Organization 1995. *Physical Status. The Use and Interpretation of Anthropometry*. Technical Report Series No. 854. Geneva: World Health Organization.
- World Bank Report. 2009. (Retrieved on 12 October 2014).

Paper received for publication on May 2017

Paper accepted for publication on October 2017