

Effects of Warm-up Exercises on Spirometric Measurements in Athletes and Sedentary Individuals

Aynur Otag¹ and İlhan Otag²

¹*Department of Physiotherapy and Rehabilitation, Health Sciences Faculty, Cumhuriyet University, Sivas, Turkey*

²*Vocational School of Health, Cumhuriyet University, Sivas, Turkey*

KEYWORDS Athlete. Exercise. Sedentary. Stretching. Spirometry

ABSTRACT Regular physical activity is important for health. The present study aimed to examine the effects of warm-up exercises involving stretching on Spiro metric measurements in 30 athletes and 30 sedentary individuals. This study was designed as a randomized, crossover study. Initially, Spiro metric measurements were conducted before and after treadmill exercise in accordance with the Bruce protocol. After 15 days, Spirometric measurements were conducted before and after a warm-up program and the same exercise test. With warm-up exercises, the forced expiratory volume in 1 second increased in the athlete group. Additionally, the peak expiratory flow, forced inspiratory flow (25-75%), and $V_{max_{25}}$ increased with warm-up exercises. However, warm-up exercises did not influence Spiro metric measurements in the sedentary group. Among athletes, ventilation volume values increased generally with warm-up exercises. For this reason, we can say that warm-up exercises increase muscle strength, resulting in optimization of aerobic function.

INTRODUCTION

Warm-up exercises typically include low-intensity aerobic exercises, stretching exercises, and special exercises (Fradkin et al. 2010; Neiva et al. 2014). Warm-up exercises increase the amount of blood flow to the muscles and tendons. The flexibility and temperature of muscles increase with warm-up; therefore, movement becomes easier (Gray et al. 2002; Bishop and Maxwell 2009; Fradkin et al. 2010; Neiva et al. 2014). In healthy individuals, the limiting factors for aerobic capacity include the cardiovascular system and skeletal muscles. In endurance athletes and those with pulmonary diseases, the respiratory system limits exercise capacity (Burtscher et al. 2011). Pulmonary function tests can provide information on the pulmonary system (Darby and Pohlman 1999; Albayrak et al. 2002; Basso et al. 2010; Layton et al. 2011; Atabek 2015; Burnett et al. 2016). Spirometry measurements are one of the most widely used methods (Layton et al. 2013; Özgül et al. 2015). These measurements

are forced vital capacity (FVC), forced expiratory volume in 1 second (FEV1), peak expiratory flow (PEF), which is frequently used to distinguish between airway obstruction and restrictive diseases (FEV1/FVC), peak value of forced inspiratory flow (PIF), forced expiration period with the condition of lasting less than 6 seconds (FET), and volume withdrawal speeds of forced vital capacities of twenty-five percent, fifty percent, and seventy-five percent ($V_{max_{25}}$, $V_{max_{50}}$, and $V_{max_{75}}$, respectively) (Miller et al. 2005; Evans et al. 2015). The exercise performance may be restricted by the functionality of the respiratory muscles. In this regard, there are many studies about the effects of specific warm-up exercises for respiratory muscles on Spiro metric measurements (Lin et al. 2007; Wilson et al. 2014; Taylor et al. 2015). However, there are no studies in the literature about the effects of general warm-up exercises on Spiro metric measurements. This present study aimed to determine the effects of warm-up exercises on Spiro metric measurements in athletes and sedentary individuals.

Address for correspondence:

Aynur Otag, PhD
Department of Physiotherapy and Rehabilitation,
Health Sciences Faculty, Cumhuriyet University,
Sivas, Turkey
E-mail: otgaynur@gmail.com

METHODOLOGY

This study was designed as a randomized, crossover study. The Research and Ethics Com-

mittee of the University of Cumhuriyet granted approval of the study after verifying that it fulfills the criteria laid out by the Helsinki Declaration. The subjects visited the laboratory twice. During their first visit, exercise test was performed with the Bruce protocol on a treadmill (Safrit 1990). In the exercise test, the speed of the treadmill was adjusted to 2.7 km/h and tilt was adjusted to 10 degrees in the first stage. A total of 7 stages were applied, with each lasting 3 minutes. At each stage, the speed and tilt were increased, according to the exercise protocol. When the subjects became tired, the test was terminated. The exercise lasted for approximately 21 minutes. Spirometric measurements were conducted before and after exercise. European Respiratory Society standards were used for the spirometric measurement methods (Miller et al. 2005). After 15 days, during their second visit, warm-up exercise was applied, the Bruce protocol was repeated, and spirometric measurements were performed before and after exercise (Table 1).

The study included 30 healthy male volunteers who were amateur football players for 5 years (athlete group) and 30 male volunteers who had no regular physical activity (sedentary group) (Table 2). Ethical approval was obtained from Cumhuriyet University Clinical Research Ethics Committee.

Table 2: The demographic data of the subjects

<i>N=30</i>	<i>Sedentary</i>	<i>Athlete</i>
Age	22.40± 2.66	23.06± 1.98
Height (cm)	176.20± 7.63	179.16± 4.69
Weight (kg)	66.33± 8.28	73.46± 8.96

All statistical analyses were performed using SPSS 22 Windows software. For comparisons between the athlete and sedentary groups, the t-test was used. For comparisons among the groups, the paired t-test was used in dependent subjects, p-value <0.05 was considered significant.

RESULTS

In the sedentary group, no significant changes were observed in the spirometric measure-

ments after exercise with warm-up, while in the athlete group, significant changes were observed in the Spiro metric measurements, particularly in expiratory volume flow measurements, after exercise with warm-up. The FVC ($p=0.001$) and FET ($p=0.031$) values exhibited a significant decline, while the FEV1 value increased ($p=0.002$). The PEF ($p=0.004$), forced inspiratory flow (FEF) twenty-five – seventy-five percent ($p=0.040$), and $V_{max_{25}}$ ($p=0.018$) values increased in the athlete group with warm-up (Table 3).

DISCUSSION

Warm-up exercises are used in almost every sports area to increase physical performance and prevent injuries prior to physical activity (O'Leary et al. 2014; Baklouti et al. 2015; McCrary et al. 2015). Many researchers emphasize that warm-up exercises increase muscle temperature and muscle contractions, increasing blood flow to a safe state that increases vasodilation (Guidetti et al. 2007; Fradkin et al. 2010; Neiva et al. 2014).

Warm-up exercises typically include aerobic exercises followed by stretching and specific exercises at low intensity (Fradkin et al. 2010). In the literature, there are many publications about the contribution of warm-up exercises to the prevention of injuries (Gray et al. 2002; Guidetti et al. 2007; Fradkin et al. 2010; Neiva et al. 2014; McCrary et al. 2015; Baklouti et al. 2015). Additionally, there are many publications in the literature in view of respiratory performance, regarding the application of warm-up exercises on specific respiratory muscles and the relation with performance (Lin et al. 2007; Cheng et al. 2013; Sales et al. 2016). However, there are no studies on the effects of general warm-up exercises on spirometric measurements. In this regard, the researchers' study appears to be important for demonstrating the effects of general warm-up exercises on spirometric measurements in sportsmen and sedentary individuals.

With regard to the relationship between warm-up exercises and the respiratory system, in 1980, De Bruyn-Prevost and Lefebvre showed

Table 1: The exercise protocol

1. Test	Spirometric measurements	Exercise test	Spirometric measurements		
2. Test	Spirometric measurements	Jog (10 minutes)	Active Stretching exercise	Exercise test	Spirometric measurements

Table 3: Spirometric values of the subjects

		<i>Sedentary (n=30)</i>		<i>Athletes (n=30)</i>	
		<i>Without warm-up</i>	<i>Warm-up</i>	<i>Without warm-up</i>	<i>Warm-up</i>
<i>FVC (L)</i>	Pre-exercise	3.31± 0.91	3.40± 0.72	4.11± 0.9 ^{***}	4.49± 0.91 ^{***}
	Post-exercise	3.09± 0.92	3.22± 0.85	3.85± 0.86 [*]	4.25± 0.87 ^{***}
<i>FEV1 (L)</i>	Pre-exercise	2.2± 0.82	2.92± 0.79	3.23± 0.79 ^{***}	3.77± 0.78
	Post-exercise	2.80± 0.81	2.94± 0.76	3.37± 0.77 ^{***}	3.79± 0.71 [#]
<i>PEF (L/min)</i>	Pre-exercise	4.08± 1.71	4.20± 1.68	4.28± 1.57	5.43± 2.01 ^{***}
	Post-exercise	4.17± 1.42	4.52± 1.47	4.67± 1.85	5.74± 2.25 ^{***}
<i>PIF (L/min)</i>	Pre-exercise	3.35± 1.48	3.62± 1.43	4.05± 1.48	4.1± 1.11
	Post-exercise	3.40± 1.51	3.85± 1.52	3.84± 1.43	4.09± 1.25
<i>FEV1/FVC</i>	Pre-exercise	86.41± 13.93	86.52± 14.97	81.23± 14.3	84.89± 13.5
	Post-exercise	91.84± 9.24 [*]	89.81± 20.59	88.28± 11.81 [*]	88.64± 11.31 [#]
<i>FEF25/75(L/min)</i>	Pre-exercise	3.29± 1.35	3.53± 1.36	3.61± 1.25	4.22± 1.48
	Post-exercise	3.52± 1.14	3.75± 1.19	4± 1.52	4.43± 1.5 [#]
<i>Vmax25</i>	Pre-exercise	3.86± 1.76	3.84± 1.55 [*]	4± 1.52	4.97± 1.94 ^{***}
	Post-exercise	3.99± 1.45	4.28± 1.52	4.54± 1.94	5.42± 2.35 ^{***}
<i>Vmax50</i>	Pre-exercise	3.48± 1.48	3.59± 1.46	3.79± 1.38	4.45± 1.64 ^{***}
	Post-exercise	3.72± 1.21	3.92± 1.25	4.26± 1.75	4.71± 1.62 ^{***}
<i>Vmax75</i>	Pre-exercise	2.48± 0.98	2.71± 0.96	2.81± 0.89	3.08± 0.98
	Post-exercise	2.68± 0.88	2.92± 0.95	2.94± 1.16	3.16± 1.04
<i>FET(min)</i>	Pre-exercise	1.51± 0.58	1.50± 0.68	1.73± 0.71	1.56± 0.54
	Post-exercise	1.35± 0.65 [*]	1.29± 0.6 [*]	1.37± 0.35	1.49± 0.45 [#]

^{*}p<0.05 compared with the previous exercise value

[#]p<0.05 compared between with and without warm-up

^{***}p<0.05 compared between the athlete and sedentary

that 5 minutes of mild warm-up exercises increased the heart rate and oxygen consumption, and reduced lactic acid buildup. In 2015, Banerjee et al. showed that the FVC and FEV1 values reduced after a bicycle ergometer test in high-school students.

In the researchers' study, no change was observed during the warm-up protocol in the sedentary group, while the FVC values decreased significantly in the athlete group after exercise, and the values were higher with warm-up than without warm-up. This finding shows that warm-up increases the temperature and contraction rate of muscles; thus, it appears to be consistent with previous findings indicating that ventilation is supported (Gray et al. 2002; Neiva et al. 2014).

In a previous study, it was stated that the functionality of the respiratory muscles limited exercise performance; therefore, these muscles should be warmed up before their use. In studies performed in different areas, it was shown that the performance increases when respiratory muscles are warmed up with special tools (Lin et al. 2007; Cheng et al. 2013; Wilson et al. 2014; Arend et al. 2015; Taylor et al. 2015; Sales et al. 2016).

In the researchers' study, routine warm-up methods were used, and the FEV1, PEF, FEF twenty-five – seventy-five percent, Vmax₂₅, Vmax₅₀,

and Vmax₇₅ values increased. These results indicate that expiratory flow volume increases in athletes, and forced expiratory volume, which is an important indicator of ventilation, improved with warm-up exercises. On the other hand, FET values reduced after warm-up exercises in both sedentary subjects and athletes, and the lower decrease in athletes supports the researchers' view that warm-up exercises have a therapeutic effect on ventilatory parameters. This view is consistent with the views of authors who mentioned that aerobic function is optimized with vasodilation and an increase in muscle blood flow that occur in veins as a result hyperthermia following warm-up exercises (Gray et al. 2002; Neiva et al. 2014).

CONCLUSION

There were differences in spirometry measurements between the athlete and sedentary groups, as well as between exercises performed with warm-up and those performed without warm-up. Warm-up exercises increase muscle strength, resulting in optimization of aerobic function. Additionally, the researchers' results may indicate that warm-up exercises accelerate the adaptation to changes in ventilation among athletes. The adaptation of the ventilatory con-

trol of the nervous system and the fact that skeletal muscles can contract better with warm-up and stretching exercises might explain this increase. In sedentary individuals, late adaptation to exercise might not change ventilation.

RECOMMENDATIONS

In the same work can be done in future studies of men and women. The effects of warm-up exercises on ventilation in women and men can be examined.

REFERENCES

- Albayrak SY, Kayserilioglu A, Dinçer C, Kasikcioglu E, Ünal M 2002. Profesyonel ve amatör futbolcuların hematolojik ve solunum parametrelerinin karşılaştırılması. *1 Tip Fak Mecmuası*, 65(1): 30-33.
- Arend M, Mäestu J, Kivastik J, Rämson R, Jürimäe J 2015. Effect of inspiratory muscle warm-up on sub-maximal rowing performance. *J Strength Cond Res*, 29(1): 213-218.
- Atabek ÇH 2015. Investigation of some pulmonary functions and bio-motoric properties in 15-17 aged group of students engaged in different sports. *Inonu University Journal of Physical Education and Sport Sciences*, 2(1): 1-16.
- Baklouti H, Chtourou H, Aloui A, Chaouachi A, Souissi N 2015. Effect of active warm-up duration on morning short-term maximal performance during Ramadan. *Libyan J Med*, 11(10): 26229.
- Banerjee S, Chaudhuri A, Roy PS 2015. Effect of exercise on undergraduate health care students. *Saudi Journal of Sport Medicine*, 15(1): 55.
- Basso RP, Jamami M, Pessoa BV, Labadessa IG, Regueiro EM, Di Lorenzo VA 2010. Assessment of exercise capacity among asthmatic and healthy adolescents. *Rev Bras Fisiote*, 14(3): 252-258.
- Bishop D, Maxwell NS 2009. Effects of active warm up on thermoregulation and intermittent-sprint performance in hot conditions. *Journal of Science and Medicine in Sport*, 12: 196-204.
- Burnett D, Burns S, Merritt S, Wick J, Sharpe M 2016. Prevalence of exercise-induced bronchoconstriction measured by standardized testing in healthy college athletes. *Respir Care*, 61(5): 571-576.
- Burtscher M, Schocke M, Koch R 2011. Ventilation-limited exercise capacity in a 59-year-old athlete. *Respir Physiol Neurobiol*, 175(1): 181-184.
- Cheng CF, Tong TK, Kuo YC, Chen PH, Huang HW, Lee CL 2013. Inspiratory muscle warm-up attenuates muscle deoxygenation during cycling exercise in women athletes. *Respir Physiol Neurobiol*, 186(3): 296-302.
- Darby LA, Pohlman RL 1999. Prediction of MAX-VO₂ for women: Adaptation of the Fox cycle ergometer protocol. *Journal of Exercise Physiology Online*, 2(4): 13-19.
- De Bruyn-Prevost P, Lefebvre F 1980. The effects of various warming up intensities and durations during a short maximal anaerobic exercise. *Eur J Appl Physiol Occup Physiol*, 43(2): 101-107.
- Evans HJ, Ferrar KE, Smith AE, Parfitt G, Eston RG 2015. A systematic review of methods to predict maximal oxygen uptake from submaximal, open circuit spirometry in healthy adults. *J Sci Med Sport*, 18(2): 183-188.
- Fradkin AJ, Zazryn TR, Smoliga JM 2010. Effects of warming-up on physical performance: A systematic review with meta-analysis. *J Strength Cond Res*, 24(1): 140-148.
- Gray SC, Devito G, Nimmo MA 2002. Effect of active warm-up on metabolism prior to and during intense dynamic exercise. *Med Sci Sports Exerc*, 34(12): 2091-2096.
- Guidetti L, Emerenziani, GP, Gallotta MC, Baldari C 2007. Effect of warm up on energy cost and energy sources of a ballet dance exercise. *Eur J Appl Physiol*, 99(3): 275-281.
- Layton AM, Garber CE, Thomashow BM, Gerardo RE, Emmert-Aronson BO, Armstrong HF et al. 2011. Exercise ventilatory kinematics in endurance trained and untrained men and women. *Respir Physiol Neurobiol*, 178(2): 223-229.
- Layton AM, Moran SL, Garber CE, Armstrong HF, Basner RC, Thomashow BM et al. 2013. Optoelectronic plethysmography compared to spirometry during maximal exercise. *Respir Physiol Neurobiol*, 185(2): 362-368.
- Lin H, Tong TK, Huang C, Nie J, Lu K, Quach B 2007. Specific inspiratory muscle warm-up enhances badminton footwork performance. *Appl Physiol Nutr Metab*, 32(6): 1082-1088.
- McCrary JM, Ackermann, BJ, Halaki M 2015. A systematic review of the effects of upper body warm-up on performance and injury. *Br J Sports Med*, 49(14): 935- 942.
- Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A et al. 2005. Standardisation of spirometry. *Eur Respir J*, 26(2): 319-338.
- Neiva HP, Marques MC, Barbosa TM, Izquierdo M, Marinho DA 2014. Warm-up and performance in competitive swimming. *Sports Med*, 44(3): 319-330.
- O'Leary JD, O'Sullivan O, Barach P, Shorten GD 2014. Improving clinical performance using rehearsal or warm-up: An advanced literature review of randomized and observational studies. *Academic Medicine*, 89(10): 1416-1422.
- Özgül F, Eliöz M, Otag A, Atan T 2015. Yüzme sporu yapan 10-14 yaş grubundaki çocukların solunum parametrelerinin karşılaştırılması. *Türkiye Klinikleri J Sports Sci*, 7(2): 35-40.
- Safrit MJ 1990. *Introduction to Measurement in Physical Education and Exercise Science*. Toronto, Boston: Times Mirror/Mosby College Publishing.
- Sales AT, Fregonezi, GA, Ramsook AH, Guenette JA, Lima IN, Reid WD 2016. Respiratory muscle endurance after training in athletes and non-athletes: A systematic review and meta-analysis. *Phys Ther Sport*, 17: 76-86.
- Taylor ST, Jared WC, Lee EB, Albert B, Tori LB, Karg P et al. 2015. Effects of respiratory muscle warm-up on high-intensity exercise performance. *Sports*, 3(4): 312-324.
- Wilson EE, McKeever TM, Lobb C, Sherriff T, Gupta L, Hearson G et al. 2014. Respiratory muscle specific warm-up and elite swimming performance. *Br J Sports Med*, 48(9): 789-791.